

Cytotoxic Effect of Some Commercial Neonicotinoids on Sheep Phagocytic Cells Activity *In vitro*

Ramadan AA^{1*} and Mansour UM²

¹Department of Animal and Poultry Sciences, Virginia Polytechnic Institute and State University, USA

²Department of veterinary Pharmacology, College of Veterinary Medicine, Zagazig University, Egypt

Research Article

Received date: 03/09/2018

Accepted date: 08/10/2018

Published date: 16/10/2018

*For Correspondence

Department of Animal and Poultry Sciences, Virginia Polytechnic Institute and State University, USA.

E-mail: ahmedibraheem60@gmail.com

Keywords: Neonicotinoids, Sheep, Polymorph Nuclear Leucocytic Cells, Lysozyme, Nitric Oxide, Phagocytosis

ABSTRACT

Oshin, Blanch and Lex are three commercial insecticides belong to neonicotinoid (a new class of agrochemicals). In the present study, sheep Polymorph nuclear leukocytes (PMNL) were exposed *in vitro* to different concentrations (100 µM, 50 µM, and 10 µM) of three insecticides (Clothianidin, Imidacloprid, and Thiamethoxam) to explore their effect on selective immune parameters such as phagocytic activity, lysozyme and nitric oxide concentrations. *In vitro* exposure of sheep PMNL to the three pesticides at different concentrations for 2 hours decreased phagocytic activity compared with control non treated cells, in dose-dependent manner. Oshin was the most powerful toxic neonicotinoid in this study. Unlike phagocytic activity, decrease of lysozyme activity was not dose-dependent. Oshin and blanch-treated cells showed significant decrease in lysozyme concentration at 100 µM dose compared with other concentrations. Lex-treated cells, at 100 µM and 50 µM there was a significant decrease in lysozyme concentration compared with 10 µM. Nitric oxide analysis showed significant decrease in Oshin-treated cells compared with both Lex and Blanch treated-cells at 100 µM. No significant difference between Oshin and Blanch treated cells at 50 and 10 µM concentration.

INTRODUCTION

Nowadays, there is increasing number of chronic cases of poisoning than acute as a consequence of environmental pollution. Diseases of the digestive, urinary, respiratory and reproductive systems caused by continuous exposure to contamination by toxic pesticides are nonspecific and manifested as a consequence of compromised immune system of animals [1,2].

Neonicotinoids represent one of the mostly used, quickest developing and promising broadly distributed class of insecticides [3-5]. Neonicotinoid pesticides have started supplanting pyrethroids, chlorinated hydrocarbons, organophosphates, carbamates, and other insecticides [6]. Neonicotinoids are an acetylcholine- insecticides that influence the central nervous system of insects resulting in paralysis and death [7] and are classified as N-nitroguanidines (imidacloprid, thiamethoxam, dinotefuran, and clothianidin) and N-cyano-aminides (acetamiprid and thiacloprid). They are utilized in a variety of venues ranging from veterinary medicine for livestock against external parasite, urban landscaping, and in many agricultural systems as agents of crop protection [8]. They have a broad spectrum insecticidal activity and a high level of efficacy at low concentrations [9]. Neonicotinoids are highly toxic to many classes of insects and exhibit relatively low vertebrate toxicity when compared with other insecticide classes currently in use [10].

Although, they are classified by the EPA [Environmental Protection Agency, 2003] as class II and III toxins of low mammalian toxicity and are labeled with a signal word: "Warning" or "Caution" [11] but in 2013, the European union restricted some neonicotinoids and its use came under increasing security over their environmental impacts [12] and in ban the outdoor use of certain neonicotinoids (Clothianidin, Imidacloprid, and Thiamethoxam). The previous studies revealed that Imidacloprid, one of the mostly used insecticide compared to organophosphate and carbamate induces significant immune-toxicity in male albino rats [13,14], mice and rat bone-marrow cells [15,16]. Furthermore, imidacloprid causes DNA strand breaks in human blood lymphocytes [17] and leukocytes in culture. There are missing information about different groups of neonicotinoids and its effect on ruminant immune system considering that neonicotinoid insecticides are a new class of agrochemicals.

In this study, we examined the effect of three different types of neonicotinoids which are Oshin (active principal is Dinotefuran),

Blanch (active principal is thiacloprid), and Lex active principal is thiamethoxam) on immune system parameters. Neonicotinoid exposure was performed in laboratory tissue culture experiments, including three concentrations as sub chronic doses (100 μ M, 50 μ M, and 10 μ M). We selected three established parameters of immunity to analyze, namely phagocytic activity, lysosome concentration, and nitric oxide concentrations of PMNL.

MATERIALS AND METHODS

Experimental Design

Sheep have been used in this experiment as a model of ruminants which are widely distributed and consumed in Egypt.

The experiment was conducted *in vitro* with multiple measurements. Each concentration of each neonicotinoid has assigned in triplicate tubes, i.e. for example the concentration of 10 μ l thiamethoxam repeated in 3 tubes and each tube was analyzed separately and data recorded. The whole experiment was repeated twice to conclude the final results.

Pesticides Preparation

Three types of pesticides were used in current study from the neonicotinoids; Thiamethoxam ($C_8H_{10}ClN_5O_3S$) molecular weight 291.7147 commercially known as Lex with active material 25%, Thiacloprid ($C_{10}H_9ClN_4S$) molecular weight 252.74 commercially known as Blanch with active material 48%, Dinotefuran ($C_7H_{14}N_4O_3$) molecular weight 202.211 commercially known as Oshin with active material 20%. Three concentrations of each pesticide were prepared 100 μ M, 50 μ M, and 10 μ M. Different concentrations of pesticides were prepared using sterile physiological saline and were adjusted to reach final concentration of 100, 50 and 10 μ M in the glass tubes after adding phagocytic cells and bacteria

Bacteria Preparation

Staphylococcus aureus bacteria were cultured onto nutrient agar. Cells harvested onto HBSS and opsonized using 10% homologous serum for 1 hour at 37°C with frequent shaking, washed twice, pellet was suspended in 1 ml RPMI without phenol and adjusted spectrophotometrically to give final concentration of 7.5×10^5 cfu/ml using RPMI.

Phagocytic Assay

The assay conducted after modification of the original assay^[18]. In sterilized glass tubes exactly 1 ml of each concentration of each pesticide was transferred and 1 ml of PMNL cells was added and incubated at 37°C with frequent shaking. After exactly one hour of incubation, one ml of *Staphylococcus aureus* bacteria was added to each tube so that PMNL: cfu is 1:15. Control tubes were prepared containing PMNL and *Staphylococcus aureus* only. Tubes were incubated for further 2 hours with frequent gentle shaking. At the end of the incubation period, samples were taken from each tube and laid on glass microscopic slides and spread over the slide, fixed with methanol and stained with Giemsa stain for microscopic examination. A total of 50 phagocytic cells engulfing *Staphylococcus aureus* were counted per each slide.

Nitric Oxide Concentration Assay

Measurement of nitric oxide was assessed according to the assay described by^[19]. Briefly, 100 μ l from each sample was transferred into flat-bottom 96-well ELISA plate and 100 μ l of Griess reagent (0.5% sulfanilamide; Sigma Chemical Co.) in 2.5% phosphoric acid (Mereck Co.) and 0.05% N-(1-naphthyl) ethylenediamine dihydrochloride (Sigma Co.). The mixture was incubated at 21°C for 10 minutes. Absorbency of the samples and standards was measured at 570 nm using ELISA reader (Dynatech MR7000; Dynatech Laboratories Inc.). Absorbency of test samples was converted to micromolar of nitrite by comparison with absorbance values of sodium nitrite (Sigma Co.) standard curve within a linear curve fit.

Lysozyme Concentration Assay

Assaying for lysozyme was done according to Peerters and Vantrape^[20]. Briefly, lysoplates were prepared by dissolving 1% agarose in 0.067 M PBS at pH 6.3 and heated to 100°C for complete dissolve, then cooled down to 60°C-70°C, 500 mg uniform suspension of *Micrococcus lysodeikticus* in 5 ml saline was added to 1 liter of agarose and mixed well. Plates were poured at thickness of 4 mm depth and left to cool down and wells 2 mm in diameter in 4 X 4 rows, 15 mm apart were cut in agarose. At time of assay, plates, serum samples, and standard lysozyme solutions were brought at room temperature. Exactly 25 μ l of each serum sample were put in 2 wells, the same was done with standard lysozyme solutions. Plates were incubated at room temperature for 12-18 hours. The diameter of clearance zone around each well was measured to the nearest 0.1 mm. A standard curve was developed for lysozyme standard solution at concentrations of 3, 15, 30, 60 and 120 μ g/ml. Log of concentrations were plotted against diameter of corresponding cleared zone on a semi-logarithmic graph.

RESULTS

Phagocytic activity of PMNL cells of sheep (**Figure 1**) has significantly affected by increasing concentration of each neonicotinoid used in this study. The decreased phagocytic activity showed dose-dependent manner. Oshin was the most inhibiting neonicotinoide in this study (data not showed).




Figure 1. Phagocytic activity of sheep PMNL in presence of different concentrations of 3 neocotinoids. Superscripts with different letters within same group are significantly different (P<0.05).

Significant difference in phagocytic activity detected between Lex and Blanch at 100 µM (**Figure 2**). Meanwhile, significant difference between Oshin and both Blanch and Lex at concentration 50 µM and 10 µM.




Figure 2. Comparison between 3 different concentrations of neonicotinoid on phagocytic activity of sheep PMNL. Oshin significantly decreased phagocytic activity at all concentration other than the other 2 neonicotinoids.

Significant difference observed between control and the three neonicotinoids used in this study. Unlike phagocytic activity, lysozyme activity was not dose-dependent (**Figure 3**). In Oshin and Blanch groups there was significant decrease in lysozyme concentration only at 100 µM dose compared with other concentrations. In Lex group, 100 µM and 50 µM significantly decreased lysozyme concentration.




Figure 3. Lysozyme activity of sheep PMNL in presence of different concentrations of 3 neonicotinoids. Superscripts with different letters within same group are significantly different (P<0.05).

Significant difference between Lex and both Oshin and Blanch at 100 µM concentration and at 50 µM concentration, has significant difference between only Blanch and Lex (**Figure 4**). No significant difference between the three neonicotinoids at 10 µM concentration.

Lysozyme activity of sheep PMNL cells in presence of 3 different concentrations of three neonicotinoids




Figure 4. Comparison between 3 different concentrations of neonicotinoids on lysozyme activity of sheep PMNL.

Nitric oxide activity of PMNL cells of sheep has significantly affected by increasing concentration of each neonicotinoids used in this study. The decreased nitric oxide activity showed dose-dependent manner (Figure 5).




Figure 5. Nitric oxide activity of sheep PMNL in presence of different concentrations of 3 neonicotinoids. Superscripts with different letters within same group are significantly different (P<0.05).

Significant difference between Oshin and both Lex and Blanch at 100 µM (Figure 6). Meanwhile, significant difference observed between Lex and both Blanch and Oshin at 50 µM and 10 µM. No significant difference between Oshin and Blanch at concentration 50 µM and 10 µM.

Nitric oxide activity of sheep PMNL cells in presence of 3 different concentrations of three neonicotinoids




Figure 6. Comparison between 3 different concentrations of neonicotinoids on Nitric oxide activity of sheep PMNL.

DISCUSSION

Healthy and compatible immune defense system is essential for animal's health. This defense can be weakened by environmental factors that may render animals more exposed to parasites and pathogens. Environmental pollution is a detrimental factor for normal immune system in animal's farm. Insecticides is required for crops and animals all over the world to combat plant and animal pathogens. Unfortunately these insecticides are also harmful for livestock. Vital systems in livestock are negatively affected by insecticides contaminated environment.

The immune system is a sensitive and complicated system that is responsible for defeating pathogens in healthy animals. Phagocytic activity is the first and crucial activity in the immune system that is responsible for engulfing and degrading invading pathogens and presenting it to the rest of the immune system to neutralize pathogens. Mainly, polymorph nuclear leukocytes (PMNL) are group of white blood cells that is responsible for phagocytic activity. Suppressing activity of PMNL could deprive the animal from proper combating of pathogens.

Neonicotinoids which is a new generation of pesticides are tested in current study. Three neonicotinoids (Oshin from Dinotefuran group, Blanch from thiacloprid group and Lex from thiamethoxam group) in current study manipulated polymorph nuclear leucocytic activity. Three concentrations of each pesticide were tested on sheep phagocytic activity *in vitro* showed dose-dependent significant decrease in this activity. The most negative effect was obtained in higher dose, 100 μM followed by 50 μM then the lowest effect was from 10 μM . The overall phagocytic activity of control non-treated cells was significantly higher than all treated cells with different concentration. Suppression of immune response to infection by exposure to neonicotinoids was evidenced in bees ^[21] and ^[22], fish ^[23] and birds ^[24]. Also in ^[25], reported that oral administration of imidacloprid (which has same chemical structure of Thiacloprid) to female mice significantly suppressed some cellular and humeral immune parameters in treated group in comparison with the control group. Similar results in honey bees, where increased exposure to neonicotinoids (Thiacloprid, Imidacloprid and Clothianidin) decreased immunity parameters were reported ^[26].

Oshin had the most powerful negative effect on sheep PMNL phagocytic activity especially at 50 μM and 10 μM . Dinotefuran, a third generation and new neonicotinoid insecticide commercialized by Mitsui Chemicals (Tokyo, Japan), could be a useful candidate in public health because it shows low mammalian toxicity and great insecticidal activity against a broad range of pests. Meanwhile it has adverse effect on immune system suppressing phagocytosis. The previous studies on Dinotefuran shows mammalian toxicity by affecting nervous and immune systems (decreased spleen and thymus weights) ^[27].

According to many researchers, lysozyme is antimicrobial enzyme and one of the most important components in the nonspecific humeral immune mechanism due to its effectiveness and wide-spread occurrence ^[28].

Mean lysozyme concentrations in the three groups were significantly lower than control non-treated cells. Significant difference between Lex and both Oshin and Blanch at 100 μM concentration. At 50 μM concentration, only significant difference between Blanch and Lex with no significant difference between the three neonicotinoids at 10 μM concentration. Kimura-Kuroda et al. ^[29] demonstrated that chemicals that bind irreversibly to specific receptors (neonicotinoids, genotoxic carcinogens and some metalloids) will produce toxic effects in a time-dependent manner, no matter how low the level of exposure. This effect of neonicotinoids may be evidenced from basic neuroscience research that the neonicotinoids have effects on mammalian neurons as well as on invertebrate ones, and that the effects on the nicotinic acetylcholine receptors (nAChRs) are more significant in vertebrates than was initially stated by the pesticide companies. Cai et al. ^[30-32] studies demonstrated a connection between the nAChRs and the immune system and suggest that neonicotinoid insecticide act a source of free radicals in exposed human cells where the ROS (reactive oxygen species) are highly reactive and produce damage of cells.

Nitric oxide (NO) is important as a toxic defense molecule against infectious organism. Cells of innate immune system, macrophage, neutrophil, and natural killer cells use pattern recognition receptors to recognize molecular patterns associated with pathogen activated macrophage then inhibit pathogen replication by releasing a variety of molecules, including NO.

Neonicotinoid application (Oshin, Thiamethoxam and Thiacloprid) in this study caused a significant decrease in nitric oxide production in tissue culture compared with control ($p < 0.05$). The decrease in nitric oxide concentration was dose dependent. This results was inconsistent with the study ^[33] confirmed the finding that imidacloprid induced the mRNA transcriptions of the three isoforms of nitric oxide synthases (iNOS, eNOS, nNOS) in brain and two isoforms (iNOS, eNOS) in the liver. Exposure to imidacloprid caused significant increase in nitric oxide production and lipid peroxidation in plasma, brain, and liver. While the superoxide-generating enzyme xanthine oxidase activity was elevated in both tissues, myeloperoxidase activity was increased only in the liver. Antioxidant enzyme activities showed various alterations following exposure, but a significantly depleted antioxidant glutathione level was detected in brain. Evidence of chronic inflammation by imidacloprid was observed as induction of pro-inflammatory cytokines such as TNF- α , IL-1 β , IL-6, IL-12 and IFN- γ in the liver and brain. In conclusion, chronic imidacloprid exposure causes oxidative stress and inflammation by altering antioxidant systems and inducing pro-inflammatory cytokine production in the liver and central nervous system of non-target organisms. The same results of increased nitric oxide by using neonicotinoids were observed in carp fish.

CONCLUSION

The results showed the presence of immune system damage at low concentrations of the three studied neonicotinoid insecticides (Oshin, Blanch and Lex). The damage of the immune system observed in the impaired phagocytic activity of the PMNL cells and in the decrease of the killing molecules released from the affected cells (namely lysozyme and nitric oxide). Oshin was the most negatively affected the phagocytic activity. The results contribute to the toxicology of these insecticides on environment and are a risk factor on animal health. The evidence of an immune system hazard related to these insecticides stresses the need for educational programs for agricultural workers to reduce the use of these chemicals in agriculture.

CONFLICT OF INTEREST

The authors declare on conflict of interest.

REFERENCES

1. Prarabdh C, et al. Immunotoxic effects of imidacloprid following 28 days of oral exposure in BALB/c mice. *Environ Toxicol Pharmacol.* 2013;35:408-418.
2. Pittis JS, et al. Effects of Coumpos on queen rearing in the honey bee, *Apis mellifera*. *Apidologie* 2004;35:605-610.
3. Jesctrike P, et al. Overview of the status and global strategy for neonicotinoids. *J Agric Food Chem.* 2011;59:2897-2908.
4. Van der Sluis JP, et al. Conclusions of worldwide integrated assessment on the risks of neonicotinoids and Fipronil to biodiversity and ecosystem functioning, *Environ Sci Pollut Res.* 2014;22:148-154.
5. Simon N, et al. Systemic insecticides neonicotinoids Fipronil trend, uses, mode of action and metabolites . *Environ Sci Pollut Res.* 2014;22:5-34.
6. Official Journal of EU (European Union).2018.L132/30.
7. Nauen R and Denholm I. Resistance of insect pests to neonicotinoid insecticides. current status and future prospects. *Arch Insect Biochem Physiol.* 2005;58:200-215.
8. Meyer K, et al. Discovery of the nitromethylene heterocycle insecticides. *Nicotinoid Insecticides and the Nicotinic Acetylcholine Receptor.* 1999;71-89.
9. Yamamoto I and Casida JE. Nicotinoid insecticides and the nicotinic acetylcholine receptor. 1999;30:37.
10. EPA, Environmental Protection Agency. 2003. Clothianidin; pesticide tolerance. *Federal Register.*68:32390-32400.
11. Fishel FM. The EPA conventional reduced risk pesticide program. Gainesville: University of Florida Institute of Food and Agricultural Sciences. 2010;224.
12. Feng S, et al. Assessing the genotoxicity of imidacloprid and RH-5849 in human peripheral blood lymphocytes in vitro with comet assay and cytogenetic tests. *Ecotoxicol Environ Saf.* 2005;61:239-246.
13. Costa C, et al. Genotoxicity of imidacloprid in relation to metabolic activation and composition of the commercial product. *Mutat Res Genet Toxicol Environ Mutagen.* 2009;672:40-44.
14. Kocaman AY and Topaktas M. In vitro evaluation of the genotoxicity of acetamiprid in human peripheral blood lymphocytes. *Environ Mol Mutagen.*2007;48:483-490.
15. Demisa G, et al. Assessment of the genotoxicity of Imidacloprid and metalaxyl in cultured human lymphocytes and rat bone marrow. *Mutat Res.* 2007;63432-63439.
16. Thea Lu, et al. Phagocytosis and killing of staphylococcus aureus by human neutrophils. *J Innate Immun.* 2014;6:639-649.
17. Rajaraman V, et al. Effect of vitamins A and E on nitric oxide production by blood mononuclear leukocytes from neonatal calves fed milk replacer. *J Dairy Sci.* 1998;81:3278-3285.
18. Peerters TL and Vantrapen GR. Factors influencing lysozyme determinations by lysoplate method. *Cli Chim Acta.* 1997;74:217-255.
19. Alaux C, et al. Interactions between *Nosema* microspores and a neonicotinoid weaken honeybees (*Apis mellifera*). *Environ Microbiol.* 2010';12:774-782.
20. Vidau C, et al. Exposure to sub-lethal doses of fipronil and thiacloprid highly increases mortality of honeybees previously infected by *Nosema ceranae*. *PLoS One.* 2011;6:e21550.
21. Sánchez-Bayo F and Goka K. Unexpected effects of zinc pyrethrin and imidacloprid on Japanese medaka fish (*Oryzias latipes*). *Aquat Toxicol.* 2005;74:285-293.
22. Cutler GC, et al. Exposure to clothianidin seed-treated canola has no long-term impact on honey bees. *J Econ Entomol.* 2007;100:765-772.

23. Prarabdh C, et al. Immunotoxic effects of imidacloprid following 28 days of oral exposure in BALB/c mice. *Environ Toxicol Pharmacol.* 2013;35:408-418.
24. Annely B, et al. The neonicotinoids thiacloprid, imidacloprid, and clothianidin affect the immunocompetence of honey bees (*Apis mellifera* L.). *J Insect Physiol.* 2016;86:40-47.
25. Lois Rossi. Environmental protection. Administrative practice and procedure, Agricultural commodities, Pesticides and pests, Reporting and record keeping requirements. 2005;78.
26. Sakurai T, et al. Enhancement of murine alveolar macrophage functions by orally administered beta-glucan. *Int J Immunopharmacol.* 1992;14:821-830.
27. Tennekes HA and Sánchez-Bayo F. Time-dependent toxicity of neonicotinoids and other toxicants: implications for a new approach to risk assessment. *J Environ Anal Toxicol.* 2012;S4-001.
28. Duzguner V and Edogaan S. Acute oxidant and inflammatory effects of imidacloprid on the mammalian central nervous system and liver in rats. *Pestic Biochem Physiol.* 2010;97:8-13.
29. Kimura-Kuroda J, et al. Nicotine-like effects of neonicotinoids on rat cerebellar neurons. *Neurosci Res.* 2011;71:399.
30. Cai B, et al. Novel insights for systemic inflammation in sepsis and haemorrhage. *Mediators Inflamm.* 2010;1-10.
31. Tripathi P. Nitric oxide and immune response. *Indian J Biochem Biophys.* 2007;25;13-15.
32. Duzguner V and Edogaan S. Chronic exposure to imidacloprid induces inflammation and oxidative stress in the liver and central nervous system of rats. *Pestic Biochem Physiol.* 2012;104:58-64.
33. Selqok O. et al. Imidacloprid exposure cause the histopathological changes activation of TNF- α , 8-OHdG biomarkers and alteration of capsizes INOs, CyplA, MTI level gene expression. *Toxicol Rep.* 2017;5:125-133.