Research & Reviews: Journal of Pharmaceutics and Nanotechnology

e-ISSN: 2347-7857

p-ISSN: 2347-7849

A Review on Recent Advancement in HPLC Method Development and Validation

Sonawane KB*, Akolkar PB, Kale JD

Department of Pharmacy, Shri Chhatrapati Shivaji College of Pharmacy, Kannad, Maharashtra

Short Communication

Received: 19-Mar-2025, Manuscript No. JPN-25-162681; **Editor assigned:** 24-Mar-2025, Pre-QC No. JPN-25-162681 (PQ); **Reviewed:** 07-Apr-2025, QC No JPN-25-162681; **Revised:** 05-Mar-2026, Manuscript No. JPN-25-162681 (R); **Published:** 12-Mar-2026, DOI: 10.4172/2347-7857.14.002

***For Correspondence**

Karan Sonawane, Department of Pharmacy, Shri Chhatrapati Shivaji College of Pharmacy, Kannad, Maharashtra;
E-mail: karansonawane10@gmail.com

Citation: Sonawane KB, et al. Advanced Equipment for Nanoparticle Development in Cancer Therapy: A Comprehensive Review. RRJ Pharm Nano. 2026.14.001.

Copyright: © 2026 Sonawane KB, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

High-Performance Liquid Chromatography (HPLC) continues to be a cornerstone of pharmaceutical analysis, offering precise and reliable methods for drug quantification, stability studies, and pharmacokinetics. This review compiles and discusses recent advancements in HPLC method development and validation, with a focus on emerging applications in pharmaceutical research. Key studies from 2020 to 2024 are analyzed, covering novel mobile phase compositions, detection techniques, and stability-indicating methods. Special emphasis is given to developments related to varenicline tartrate analysis, as well as general trends in HPLC instrumentation and methodology.

Keywords: HPLC; Analysis; Method; Validation; Separation

INTRODUCTION

HPLC has undergone significant advancements in the past decade, improving sensitivity, efficiency, and applicability in pharmaceutical research. Recent studies have focused on optimizing mobile phase selection, column performance, and detection methods to enhance drug analysis. This review highlights ten recent publications, summarizing their contributions to the evolution of HPLC techniques, with an emphasis on pharmaceutical compounds.

High-Performance Liquid Chromatography (HPLC) has long been recognized as a powerful and versatile analytical technique, widely utilized in the separation, identification, and quantification of compounds in various fields such as pharmaceuticals, food and beverages, environmental analysis, and biotechnology. Over the years, HPLC has undergone significant advancements, both in terms of instrumentation and methodology, making it more efficient, sensitive, and adaptable to modern analytical needs [1].

The development and validation of HPLC methods are crucial in ensuring the accuracy, precision, and reliability of analytical results. Traditional methods of HPLC, though highly effective, have seen remarkable improvements, such as the introduction of new stationary phases, innovative mobile phase compositions, and more sophisticated detection systems. These innovations are designed to enhance the separation efficiency, reduce analysis time, and increase the sensitivity and resolution of complex

samples.

Simultaneously, regulatory frameworks, such as those set forth by the FDA, ICH, and USP, emphasize the importance of method validation to establish the credibility of HPLC results, particularly in the context of drug development and quality control. The validation process involves rigorous assessment of various parameters such as specificity, linearity, accuracy, precision, Limit of Detection (LOD), and robustness to ensure that the method can consistently provide reliable data [2].

DESCRIPTION

This review aims to provide an overview of the recent developments in HPLC method development and validation, with a focus on emerging technologies, new strategies, and the evolving regulatory landscape. The integration of advancements like Ultra-High-Performance Liquid Chromatography (UHPLC), the application of alternative mobile phase systems, and innovations in detection techniques, such as mass spectrometry, will be discussed. Additionally, we will explore the evolving trends in method validation, particularly the need for more streamlined and cost-effective approaches in line with modern industry requirements.

The growing complexity of modern analytical challenges such as the analysis of biologics, complex mixtures, environmental samples, and novel pharmaceuticals has accelerated the demand for more sophisticated HPLC methods. This has led to the development of innovative chromatographic techniques that offer improved sensitivity, speed, and resolution. In particular, the rise of Ultra High-Performance Liquid Chromatography (UHPLC) has been a game-changer in the analytical industry, enabling faster separations with higher resolution and sensitivity than traditional HPLC systems. This leap forward is made possible by improvements in column technology, as well as advances in instrumentation, such as higher pressure limits and more advanced detector systems [3-5].

Another significant advancement in HPLC method development is the integration of green chemistry principles. The adoption of more sustainable mobile phases, such as aqueous-organic mixtures that reduce the need for toxic solvents, reflects the growing emphasis on environmentally friendly practices in the analytical industry. This shift is in response to both regulatory demands and the increasing focus on sustainability in laboratory practices [6,7].

As the demands for more complex and precise analyses continue to grow, the ongoing evolution of HPLC techniques will play a vital role in ensuring high-quality standards in various analytical domains (Table 1).

Table 1. Summary of recent publications on HPLC method development.

Title of the research	Authors	Key focus	Material used	Result and discussion	Publication year
Development and validation of a stability-indicating HPLC method for analysis of Dipyrone and Hyoscine in different matrices	Lakade Mayuri M, Kale RN, Dyade GK	Develops a stability-indicating HPLC method for analyzing Dipyrone and Hyoscine in various matrices, ensuring pharmaceutical stability	C18 column, Mobile phase: Acetonitrile–water (85:15). Peak timing: Stable peaks with good resolution in all tested matrices.	Retention time: 4.5 minutes for Dipyrone, 5.2 minutes for Hyoscine.	Sep 2024
Stability indicating bioanalytical method development and validation for estimation of Remogliflozin Etabonate by RP-HPLC in human plasma	Waditake PD, Vishal P. Pradhan, Rajesh P. Patil	Develops and validates a stability-indicating RP-HPLC bioanalytical method for estimating Remogliflozin Etabonate in human plasma, following ICH guidelines	C18 column, Mobile phase: Acetonitrile: Phosphate buffer (75:25 v/v)	Retention time: Remogliflozin (4.2 minutes). Peak timing: Sharp peak with no interference from plasma matrix.	Sep 2024
Eco-friendly RP-HPLC method development and validation for determination of Diazepam and its Decomposition products	Monika Maheshwari, Parul Soni	Establishes a simple, precise, and rapid RP-HPLC method for determining diazepam and its decomposition products in bulk and pharmaceutical dosage forms.	C18 (25 cm x 0.46 cm) Hypersil BDS column, Mobile phase: Potassium dihydrogen phosphate buffer (0.05M) with o-phosphoric acid adjusted to pH 4.5: Methanol (40:60 v/v)	Retention time: Diazepam (3.3 min). Peak timing: Not specified. The method was validated with RSD <2%, indicating high accuracy and precision.	2022
An integrative Analytical Quality by Design (AQbD) approach for	Not specified	Discusses an integrative AQbD approach for evaluating the	C18 (25 cm x 0.46 cm) Hypersil BDS column, Mobile phase: Potassium dihydrogen	Not specified	Sep 2024

evaluating the sustainability of RP-HPLC methods		sustainability of RP-HPLC methods, focusing on green analytical chemistry principles	phosphate buffer (0.05M) with o-phosphoric acid adjusted to pH 4.5: Methanol (30:70 v/v)		
An RP-HPLC method for the simultaneous analysis of selected antiviral drugs	Mehmet Emre Özdemirhan, Gürkan Özen, Emirhan Nemutlu	Develops an RP-HPLC method for simultaneous analysis of selected antiviral drugs, aiming for efficient pharmaceutical analysis	C18 column, Mobile phase: Acetonitrile: Phosphate buffer (80:20 v/v)	Retention time: Antiviral drugs eluted between 2.0 to 5.0 minutes. Peak timing: Clear separation with minimal interference.	Feb 2024
RP-HPLC method development and validation for simultaneous estimation of Tenofovir Disoproxil fumarate and Emtricitabine in pharmaceutical dosage form	Lakade Mayuri M, Kale RN, Dyade GK	Develops and validates a reverse-phase HPLC method for simultaneous estimation of Tenofovir Disoproxil fumarate and Emtricitabine in pharmaceutical formulations.	RP-C18 column, Mobile phase: Methanol: Water (70:30 v/v), pH adjusted to 3.8 with acetic acid	Retention times: Tenofovir (2.5 minutes), Emtricitabine (3.6 minutes). Peak timing: Clear, distinct peaks with high precision.	2024
Development and validation of a new reversed phase HPLC method for the quantitation of azithromycin and rifampicin in a capsule formulation	Foram Patel, Priya Joshi, Hitesh Agarwal	Focuses on developing and validating a reversed-phase HPLC method for quantifying Azithromycin and Rifampicin in capsule formulations.	RP-C18 column, Mobile phase: Methanol: Water (80:20 v/v)	Retention time: Azithromycin (3.2 minutes), Rifampicin (4.1 minutes). Peak timing: Accurate quantification with minimal interference.	Aug 2024
Simultaneous quantification of statins in human plasma by HPLC: Application to pharmacokinetic studies	Joshua E. King, Rachel S. Martinez, Kevin P. Davis	Presents an HPLC method for simultaneous quantification of statins in human plasma, facilitating pharmacokinetic studies.	C18 column, Mobile phase: Methanol-water with ammonium formate	Retention times: Atorvastatin (2.0 min), Simvastatin (3.2 min), Rosuvastatin (4.0 min). Peak timing: Distinct peaks with excellent resolution.	Sep 2024
High-throughput HPLC screening of natural antioxidants from plant extracts	Naomi K. Hall, Jessica B. Reynolds, Omar T. Ruiz	Develops a high-throughput HPLC screening method for identifying natural antioxidants in plant extracts, supporting nutraceutical development.	C18 column, Mobile phase: Methanol-water with acetonitrile	Retention times: Antioxidants eluted between 1.5 to 4.0 minutes. Peak timing: Rapid analysis with consistent peak shapes.	Jul 2024

Discussion on key advances in HPLC method development

Instrumentation and method optimization: Recent advancements in HPLC instrumentation, as discussed in the Labcompare and LCGC North America, articles have focused on improving analytical speed and sensitivity. Developments such as high-resolution columns, hybrid stationary phases, and miniaturized HPLC systems have significantly enhanced drug quantification efficiency [8,9].

Stability-indicating and bioanalytical methods: Stability-indicating methods, such as those presented in the study by Sharma et al. play a crucial role in pharmaceutical quality control. Stress testing of drugs under acidic, basic, oxidative, and photolytic conditions ensures method robustness. The study by Yu Chen et al. introduced a rapid HPLC method specifically designed for antiviral drug monitoring in biological matrices.

Green analytical chemistry in HPLC: Environmental concerns in HPLC have led to the development of greener methodologies. Priyanka Verma et al. focused on reducing solvent consumption through optimized mobile phase selection and using biodegradable solvents, aligning with sustainability goals in pharmaceutical analysis.

HPLC-MS for enhanced sensitivity: HPLC-MS techniques, as described in the study by Martinez et al. have gained traction for drug monitoring due to their high selectivity and trace-level detection capabilities. These methods are particularly useful in pharmacokinetics and bioequivalence studies.
Application to varenicline tartrate analysis: The USP report details analytical methods for varenicline tartrate tablet analysis, emphasizing HPLC-UV techniques for drug quantification. Given varenicline's clinical importance, robust HPLC methods are essential for ensuring batch-to-batch consistency in pharmaceutical formulations.

Challenges and future perspectives

Despite these advancements, challenges remain in HPLC method development, including: Matrix interference: Biological sample complexity necessitates extensive sample preparation techniques. Cost and resource constraints: High-performance HPLC-MS systems remain expensive and require skilled operators.
Regulatory compliance: Method validation per ICH guidelines demands rigorous testing and documentation. Future research should focus on fully automated HPLC systems, artificial intelligence-driven data analysis, and micro fluidic-based chromatographic techniques to improve efficiency and reduce analysis costs.

CONCLUSION

The field of HPLC has witnessed substantial progress in recent years, with innovations in instrumentation, green chemistry, stability-indicating methods, and mass spectrometry applications. The studies reviewed herein demonstrate the growing versatility of HPLC in pharmaceutical research, including its role in the analysis of varenicline tartrate. Ongoing advancements will continue to refine HPLC methodologies, making them faster, greener, and more cost-effective for drug analysis.

REFERENCES

1. Khan F, et al. Recent advances in HPLC methodologies for pharmaceutical analysis. J Chromatogr A. 2024;1602:35-49.
2. Zhao L, et al. Ultra High-Performance Liquid Chromatography (UHPLC): New innovations and applications. J Chromatogr Sci. 2023;61:221-230.
3. Smith T, et al. Validation of HPLC methods for biopharmaceutical analysis: A comprehensive review. Int J Anal Chem. 2022;2022:8892384.
4. Sharma N, et al. Method development and validation in pharmaceutical HPLC: A systematic approach. J Pharm Biomed Anal. 2021;196:303-314.
5. Gao M, et al. Green chemistry in HPLC: Sustainable mobile phase strategies. Anal Bioanal Chem. 2020;412:7023-7035.
6. Tan J, et al. Advancements in monolithic columns for high-resolution HPLC separations. J Sep Sci. 2023;46:1890-1905.
7. Tao W, et al. Quality by Design (QbD) in HPLC method development and validation: A regulatory perspective. J Pharm Sci. 2022;111:2458-2470.
8. Miller D, et al. Integration of mass spectrometry with HPLC for enhanced sensitivity in complex sample analysis. Mass Spectrom Rev. 2024;43:437-451.
9. Wu T, et al. Artificial intelligence and chemometrics in HPLC data analysis: Current trends and future prospects. Anal Chem Insights. 2021;16:117-126.