

Activity of Acetylcholinesterase (AChE) in *OxycarenusHyalinipennis*Costa (Hemiptera: Lygaeidae, 1847) After Exposure to *Chrysanthemum Indicum* I. ExtractRupture

Muhammad Abubaker Hassan*, Shoab Freed

Department of Entomology, Faculty of Agricultural Sciences and Technology, Bahaudinzakariya University, Multan, Pakistan

Research Article

Received date: 04/09/2020

Accepted date: 19/09/2020

Published date: 24/09/2020

*For Correspondence

Muhammad Abubaker Hassan,
Department of Entomology, Faculty of
agricultural sciences and technology,
Bahaudinzakariya university, Multan,
Pakistan

E-mail: ahawan97@outlook.com

Keywords: Hyalinipennis; Chrysanthemum
Indicum; Botanical Extract; Diptera; True Bug

ABSTRACT

OxycarenusHyalinipennis Costa (Hemiptera: Lygaeidae) is serious pest of *GossypiumHirsutum* L. which is commercially grown to generate foreign exchange reserves for Pakistan. Both adult and nymph causes reduction of oil content, weight loss of seeds, germination rate and stains cotton with pink color during ginning. The main objective of the study was to check the effect of *Chrysanthemum Indicum* flower extract on the activity of acetylcholinesterase (AChE) in *O. Hyalinipennis*. The adults of *O. Hyalinipennis* were exposed to different concentration (8, 10, 12 and 14 mg/20 ml) by leave dip method prior to biochemical analysis of surviving bug samples. The results demonstrated that *C. Indicum* extract affect the activity of AChE in *O. Hyalinipennis* adult. Maximum enzymatic activity was 0.544 α -Na μ mol/min/ mg protein at day 1 in T3 (12 mg/20 ml) treatment while in control maximum activity was 0.160 α -Na μ mol/min/ mg protein at day 4 whereas highest and lowest mortality was (66.66 and 0%) at 14 and 8 mg/20 ml concentrations respectively. The rise in activity of AChE is one of the main reasons to develop resistance in insect. This can be managed by adaptation of alternative control methods.

INTRODUCTION

GossypiumHirsutum L is commercially grown as a source to generate foreign exchange reserves for Pakistan^[1]. The average yield of cotton in Pakistan is lower as compared to other cotton growing countries^[2]. About 30-40% yield loss is associated with insect pests which are also considered as vital limiting factors^[3,4].

*OxycarenusHyalinipennis*Costa (Hemiptera: Lygaeidae) also known as Dusky cotton bug (DCB) is an economic pest of family malvaceae^[5,6]. Both adult and nymph feeds on seeds and leaves which results in both quantitative and qualitative losses to various plants including cotton in most countries of the world^[7-9]. *O. Hyalinipennis* Causes economical damage by decreasing oil contents, weight loss of seeds, germination rate and also stains cotton lint with pink color during the process of ginning due to crushing of bugs^[8,10]. Resistance has been developed by *O. Hyalinipennis* against several insecticides like imidacloprid, spirotetramat, triazophos, fipronil and nitenpyram but resistance level varies from low to very high which is due to the misuse of insecticides in Pakistan^[11-13]. The occurrence of *O. Hyalinipennis* increasing in *Bt*Cotton genotype due to lesser number of insecticide applications for the management of bollworms. With the growing awareness about the damages caused by DCB the necessity for its management is gaining more attention^[14]. Nowadays, the only solution available for the farmer is to use chemical insecticides and almost 90% of farmer use chemical insecticides to protect their crops and regular insecticide exposer over a long course of time lays the foundation of resistance. Insect develop resistance by changing target site, decrease infiltration and metabolic resistance arbitrate by detoxifying enzymes such as acetylcholinesterase (AChE), glutathione transferases (GST), and esterases (EST) support to improve resistance against pesticides^[15].

Awareness about hazards related to use of synthetic pesticide resulted in need to discover alternate methods to control pest. Products of biological nature in their raw form or oil extract offers limitless opportunities as bio-pesticides. Ecofriendly nature of bio-pesticides make them best control method against agriculture pest^[16]. Most plants used for medicinal purposes contains compounds which restrict the growth of insect and feeding activity or behave like repellents^[17]. *Chrysanthemum Indicum* L has anti-inflammatory, immunomodulatory and anti-microbial compounds like 1,8-cineole, camphor, borneol and bornyl acetate^[18]. Looking at the important cases of pesticide resistance and their ecological effects. The present experiment was conducted to evaluate the effect of acetone extract of *C. Indicum* flowers on activity of AChE in *O. Hyalinipennis*.

MATERIALS AND METHODS

Plant collection and extraction

Chrysanthemum Indicum flowers were collected from Bahauddin Zakariya University, Multan and its adjacent areas. The collected flowers were washed 3 times using tap water and rinsed by distilled water and shade dried. *C. Indicum* petals were milled using a domestic grinder and kept in air tight jar for future use. 50 gram of *C. Indicum* powder was taken in a beaker with 100 ml of acetone solvent and stirred with magnetic stirrer and afterwards left for 24 hours.

Insect culture

O. Hyalinipennis nymphs and adults were collected from cotton fields of District Multan. The collected insect population were maintained in the laboratory condition with $27 \pm 2^\circ\text{C}$ of temperature and 60%-70% relative humidity within plastic jar and feed on fresh leaves of *Hibiscus rosa-sinensis* L. The insects from F_1 generation were utilized for the examination.

Bioassay

Botanical extract of *C. Indicum* was tested to find out its lethal and sub-lethal effect against *O. Hyalinipennis*. Different concentrations (8,10,12,14 ml/20 ml solution) were formed for *C. Indicum*. Fresh leaves of *H. rosa-sinensis* were used for the treatment by leaf dip-method and treated leaves were dried by placing them on tissue paper than shifted in polystyrene petri dish separately having five newly emerged adults of *O. Hyalinipennis*. For each treatment four replication were made. The leaves in control group were treated with distilled water. The mortality data was collected after every 24 hours for next 96 hours.

Enzyme assay

After treating *O. Hyalinipennis* with *C. Indicum* extract than activity of acetylcholinesterase was noticed following the procedure of SEREBROV with minor alterations^[19]. The treated *O. Hyalinipennis* was crushed within 1.5 ml Eppendorf tubes containing 50 ml of NaCl (0.15 M). After crushing, the samples were centrifuged at 5000 rpm for 10 minutes in centrifuge machine and only supernatant were used. Samples were made for four sequential days for treated adults with four replications were used for each concentration of *C. Indicum* extract for AChE evaluation.

Assay of acetylcholinesterase

ELLAMAN method was used to check the activity of ACh by using acetylcholine iodide (0.075M) as substrate and changes in absorbance at 412nm wavelength were noticed at 20 sec interval^[20].

Data analysis

Experimental data and values were applied to analysis variance using Microsoft Excel 2013 and Statistix 8.1 software were used for statistical analysis and making graphs to explain the detoxification activities of AChE enzyme.

RESULTS

Animals

Extract of *C. Indicum* flowers doses of 8, 10, 12 and 14 mg per 20 ml were applied against adults of *Oxycarenus Hyalinipennis* Costa. Among these doses 10 ml give maximum mortality of *O. Hyalinipennis* followed by 8, 12, 14 ml, respectively (**Figure 1**).




Figure 1. Percentage motility of *O. Hyalinipennis* exposed to different doses of *C. Indicum* flower extract.

The lowest mortality percentage (0%) was observed in T2, T3, T4 and control after 24 hours while the highest mortality percentage (66.66) was observed in T2 after 96 hours.

Activity of acetylcholinesterase (AChE)

After treating *O. hyalinipennis* with different doses of extract, activity of AChE (acetylcholinesterase) was observed. In treated samples, maximum enzymatic activity was 0.544153 α -Na μ mol/min/ mg protein at day 1 in T3 (12 mg/20 ml) treatment while in control maximum activity was 0.160311 α -Na μ mol/min/ mg protein at day 4 (Figure 2 and Table 1).

Table 1. AChE Activity (α -Na μ mol/min/ mg protein) comparison at high, low and control doses on 4 consecutive days.

		Days			
		1	2	3	4
α -Na μ mol/ min/protein	High dose	0.075066	0.045663	0.147424	0.864187
	Low dose	0.158137	0.13046	0.271435	0.098294
	Control	0.05309	0.08402	0.160311	0.163331




Figure 2. Acetylcholinesterase (AChE) activity in adult of *O. Hyalinipennis* after exposing with *C. Indicum* flower extract on 4 consecutive days.

DISCUSSION

Biochemical and bioecological reasons are believed to be the source of insect resistance mechanisms against several insecticides^[21]. Pyrethroids were widely used against most agricultural pests, including *O. Hyalinipennis*^[22]. *Chrysanthemum Indicum* plant had insecticidal properties against many serious pests because of natural source of pyrethrin. Rajalakshmi justified the anti-microbial activity against plant pathogen by *C. indicum* leaf extract^[23]. KAMARAJ observed highest toxic effects of acetone and methanol *C. Indicum* leaf extract against mosquitolarvae^[24]. Outcome of present analysis support results of previous analysis on detox enzymes of insects. When construction of detox enzyme increased in insect body they caused the insensitivity of target sites or by detoxification of foreign elements^[25]. Insects detoxify insecticides by various enzymes like acetylcholinesterase (AChE), acid and alkaline phosphatases and excrete them from their body^[26]. Acetylcholine (AChE) is a neurotransmitter which is present in the nervous system which is hydrolyzed by acetylcholinesterase (AChE) and high resistance was detected due to increase activity of AChE activity in *Tribolium Castaneum* after exposing with permethrin and diflubenzuron which shows direct relationship of resistance development with enzymes in insects^[27,28]. In recent study similar results were observed after exposing *O. hyalinipennis* with various doses of *C. indicum* flower extract. Among all doses the maximum AChE activity was observed in 12 mg/20 ml sample of day 1 while lowest was 8 mg/20 ml sample of day 4.

CONCLUSION

The overall results suggest the potential of *C. Indicum* flower extract as a bio pesticide for crop protection against *O. Hyalinipennis* but further investigation is needed on behalf of active ingredients accountable for insecticidal properties to conclude some absolute recommendations.

REFERENCES

1. Mallah G, et al. A review of varietal resistance and crop management to control whitefly in cotton. Pak Cott. 1997;41:46-51.
2. Bakhsh K, et al. Factors affecting cotton yield: A case study of Sargodha (Pakistan). J Agric Soc Sci. 2005;1:332-334.
3. Haque H, et al. Insect pests of fiber crops in: AA Hashmi (Ed.) Insect pest management, Cereal and cash crops. Pak Agri Res Coun. 1994;193-206.
4. Kannan M, et al. Impact of insecticides on sucking pests and natural enemy complex of transgenic cotton. Curr Sci. 2004;86:726-729.
5. EL-RAHIM A, et al. Population fluctuations of *Oxycarenus Hyalinipennis* and effect of certain compounds on its population on okra in Asyut Governorate. Egypt. J Agric Res. 2015;9:25-34.
6. Sanghi AH, et al. Efficacy of different insecticides against dusky cotton bug *Oxycarenus Hyalinipennis* (Hemiptera: Lygaeidae) in ecological zone of Rahim Yar Khan. Int J Comprehen Res Biol Sci. 2014;1:49-54.
7. Atta B, et al. Toxicity of some insect growth regulators (IGRS) against different life stages of dusky cotton bugs *Oxycarenus Hyalinipennis* Costa (Hemiptera: Lygaeidae: Oxycareninae). Bulg J Agric Sci. 2015;21:367-371.
8. Khan M. et al. Toxicity of neem fruit extract and seed oil against *Oxycarenus* (Heteroptera) of cotton crop. Acta Biol Cracoviensia Ser Zool. 2000;42.
9. Rajashekhargouda R, et al. New record of insect pests infesting kasturibhendi, *Hibiscus abelomoschus* Linnaeus, a medicinal plant. J Bombay Nat Hist Soc. 1984;81:212-213.
10. Henry T. Pests not known to occur in the United States or of limited distribution. USDA- APHIS-PPQ. APHIS. 1989;43:1-6.
11. Ullah S, et al. Study of synergism, antagonism, and resistance mechanisms in insecticide-resistant *Oxycarenus Hyalinipennis* (Hemiptera: Lygaeidae). J Econ Entomol. 2017;110:615-623.
12. Ullah S, et al. Toxicity of insecticides, cross-resistance and stability of chlorfenapyr resistance in different strains of *Oxycarenus Hyalinipennis* Costa (Hemiptera: Lygaeidae). Crop prot. 2017;99:132-136.
13. Ullah S, et al. Resistance of dusky cotton bug, *Oxycarenus Hyalinipennis* Costa (Lygaeidae: Hemiptera), to conventional and novel chemistry insecticides. J Econ Entomol. 2015;109:345-351.
14. Prayogo Y, et al. Prospect of entomo-pathogenic fungus *Metarhizium Anisopliae* to control *Spodopteralitura* on soybean. J Litbang Pertanian. 2005;24:19-26.
15. Ahmad M, et al. Penetration and metabolism of trans-cypermethrin in a susceptible and a pyrethroid-resistant strain of *Helicoverpa armigera*. Pestic Biochem Physiol. 1999;65:6-14.
16. Abbad M, et al. Bioassay of the botanical insecticide, Tondexir, on two natural enemies of the common pistachio psyllid. Int J Agron Plant Prod. 2013;4:1191-1196.
17. WU LY, et al. Analysis of chemical composition of *Chrysanthemum indicum* flowers by GC/MS and HPLC. J Med Plant Res. 2010;4:421-426.
18. Shunying Z, et al. Chemical composition and antimicrobial activity of the essential oils of *Chrysanthemum indicum*. J Ethnopharmacol. 2005;96:151-158.
19. Serebrov V, et al. Effect of entomopathogenic fungi on detoxification enzyme activity in greater wax moth *Galleria mellonella* L. (Lepidoptera, Pyralidae) and role of detoxification enzymes in development of insect resistance to entomopathogenic fungi. Biol Bull. 2006;33:581.
20. Ellman GL, et al. A new and rapid colorimetric determination of acetylcholinesterase activity. Biochem Pharmacol. 1961;7:88-95.
21. Damayanthi B, et al. Biochemical characterization of insecticide resistance in insect pests of vegetables and predatory ladybird beetles. J Nat Sci Found Sri Lanka. 2005;33:115-122.
22. Akram M, et al. Bioefficacy of organophosphates, pyrethroids and new chemistry insecticides against a field population of dusky cotton bug, *Oxycarenus* spp. (Hemiptera: Oxycarenidae) in Bt cotton ecosystem. Pak J Life Soc Sci. 2013;11:48-52.
23. Rajalakshmi G, et al. In vitro micropropagation and antimicrobial activity of *Chrysanthemum indicum*. Sch Acad J Pharm. 2013;4:285-288.

24. KAMARAJ C, et al. Larvicidal potential of medicinal plant extracts against *Anopheles subpictus* Grassi and *Culex tritaeniorhynchus* Giles (Diptera: Culicidae). *Parasitol Res.* 2009;104:1163.
25. Zhang L, et al. Quantitative and qualitative changes of the carboxylesterase associated with beta-cypermethrin resistance in the housefly, *Musca domestica* (Diptera: Muscidae). *Biochem. Mol Biol.* 2010;156:6-11.
26. Zibae A, et al. The effect of diazinon on some biochemical characteristics of *Chilo suppressalis* Walker (Lepidoptera: Pyralidae), rice striped stem borer. *MunEntomol Zool.* 2008;3:255-265.
27. Saleem MA, et al. Joint effects of Dimilin and Ambush on enzyme activities of *Tribolium castaneum* larvae. *Pest Biochem Physiol.* 1987;29:127-137.
28. Wang JJ, et al. The effect of the insecticide dichlorvos on esterase activity extracted from the psocids, *Liposcelis bostrychophila* and *L. entomophila*. *J Insect Sci.* 2004;4:23.

