

## Biological Potential and Phytopharmacological Screening of *Gomphrena* Species

Y Arsia Tarnam<sup>1</sup>, \*M H Muhammad Ilyas<sup>2</sup>, T Nargis Begum<sup>1</sup>

1. Department of Biotechnology, Jamal Mohamed College, 7, Race Course Road, Khajanagar, Tiruchirapalli-620 020, India.
2. Department of Botany, Jamal Mohamed College, 7, Race Course Road, Khajanagar, Tiruchirapalli-620 020, India.

### ABSTRACT

The traditional system of medicine consists of large number of plants with various medicinal and pharmacological importances and hence considered as a priceless tank of new bioactive molecules. The *Amaranthaceae* family comprises of many species, which are used in nutrition and in traditional folk medicine for the treatment of several diseases such as infections, inflammation and fever. *Gomphrena* species found all over the world. It has been recognized in different traditional system of medicines for the treatment of various diseases of human being. *Gomphrena* species is an edible, commercial ornamental and medicinal plant commonly known as Globe Amaranth or Bachelor Button, belongs to the family Amaranthaceae. It comprises approximately 120 species found in the America, Antartica and Indo-Malaysia. 46 species found in Brazil. The phytoconstituents present in it are flavonoids, phytosterols, phenolics and terpenoids. The different parts of this plant are traditionally claimed to be used for the treatment of broad spectrum of ailment including baby gripe, oligouria, body sore, malaria, bacterial infections, jaundice, urinary problems, high cholesterol, cough, fever, diarrhea, liver disorders, kidney disorders and cooling. Due to changes in the lifestyle and associated decline of the use of plants, the ethnobotanical knowledge is continuing to decline. Hence further work to isolate and characterise the active compounds responsible for this activities in the plant is recommended.

**Keywords:** Alkaloid, ethnobotany, flavanoid, gomphrena, phytochemistry, pharmacology

Received 11 Nov 2013

Received in revised form 29 Nov 2013

Accepted 01 Dec 2013

### \*Address for correspondence:

**Muhammad Ilyas MH**

Jamal Mohamed College, P.Box.No.808, 7, Race Course Road, Khajanagar, Tiruchirapalli-620 020, India  
E-mail: jinjian31@126.com

### INTRODUCTION

Medicinal plants have been playing a vital role on the health and healing of man since down of the human civilization. In spite of tremendous development in the field of allopathic medicines during the 20<sup>th</sup> century, plants still remain one of the major sources of drugs in modern as well as in traditional system of medicine [1, 2]. The green plants synthesise and they preserve a variety of biochemical products, many of which are extractable and used as a chemical feed stocks or as raw material for various scientific investigations. Many secondary metabolites of plant origin are commercially important and find use in a number of pharmaceutical compounds [3]. Nations with highly developed pharmaceutical industries are mainly interested in plants as a source of biologically active and medicinal important

compounds which might be lead to discovery of new and better drugs with pharmacological potency [4]. The chemical constituents present in the herbal medicine or plants are the major part of the physiological functions of living flora and hence they are believed to have better compatibility with the human body.

Natural products from plants are a rich resource used for centuries to cure various ailments. The use of bioactive plant-derived compounds is on the rise, because the main preoccupation with the use of synthetic drugs is the side effects which can be even more dangerous than the diseases they claim to cure. In contrast, plant derived medicines are based upon the premise that they contain natural substances that can promote health, alleviate illness and proved to be safe, better patient tolerance,

relatively less expensive and globally competitive. So, in respect of the healing power of plants and a return to natural remedies is an absolute requirement of our time [5, 6].

In this present review paper, an attempt has been made to summarize all the information so far available on the floristic composition, ethnobotanical practices on the edible and ornamental plant *Gomphrena* species. *Gomphrena* species is an edible, commercial ornamental and medicinal plant commonly known as Globe Amaranth or Bachelor Button, belongs to the family Amaranthaceae. It comprises approximately 120 species found in the America, Antartica and Indo-Malaysia. 46 species found in Brazil.

The Amaranthaceae is a cosmopolitan family which occurs at disturbed, arid or saline areas; one of the characteristics that ensure

its survival in adverse environments is the operation of C4 pathway of photosynthesis [7]. It is erect or ascending herbs to 50cm. leaves spiral, deltoid to elliptic ovate. Inflorescence is axillary, slender, thyriform spike. Flowering mostly occurs in June to December. Globe amaranth [*Gomphrena globosa* L.] might be one of the plants that can use atmospheric sulfides for its growth [8, 9].

**TAXONOMY**

The *Amaranthaceae* family, class Magnoliopsida and order Caryophyllales, was established by A. L. Jussieu in 1789 and includes approximately 176 genera and 2400 species [6]. The members of *Amaranthaceae* family are divided in four tribes: Celosieae, Achyrantheae, Braylineae and Gomphreneae. The species in Brazil are distributed in Celosieae, Achyrantheae and Gomphreneae tribes [10].

Kingdom	Plantae
Subkingdom	Tracheobionta
Superdivision	Spermatophyta
Division	Magnoliophyta
Class	Magnoliopsida
Subclass	Caryophyllidae
Order	Caryophyllales
Family	Amaranthaceae
Genus	<i>Gomphrena</i> L.

**PHYTOCHEMISTRY**

The previous chemical studies with species of this genus were related to the isolation of hydrocarbons, alcohols, steroids, terpenoids, ecdysteroids, flavonoids and protoalkaloids. Despite of some phytochemical and biological activity studies have been performed in certain species, overall the genus *Gomphrena* still remains poorly studied [10].

Some species of *Amaranthaceae* have been studied seeking the characterization of inorganic elements and heavy metals in

their tissues. Some of these works [Table 1] suggest that these vegetables can bioconcentrate inorganic elements and heavy metals. And using the X-ray fluorescence analysis, the results showed that several inorganic elements like Al, Si, P, S, Cl, K, Ca, Ti, Cr, Mn, Fe, Co, Ni, Cu, Zn, As, Br, Rb, Sr, Mo, Cd, Sn, Sb, Ba and Pb could be detected in the extracts and powder of species of Amaranthaceae [10].

**Table 1: Inorganic Elements Detected in Two *Amaranthaceae* Species [10].**

Species	Inorganic elements detected
<i>Gomphrena glabrata</i> <sup>a</sup>	P, S, Cl, K, Ca, Cr, Mn, Fe, Co, Ni, Cu, Zn, Br, Sr, Cd, Ba and Pb
<i>Gomphrena globosa</i> <sup>d</sup>	S, K, Ca, Sc, Ti, V, Mn, Fe, Co, Cu and Zn
<i>Gomphrena globosa</i> <sup>b</sup>	S, K, Ca, Fe, Co, Cu and Zn

Part used: whole plant<sup>a</sup>, aerial part<sup>b</sup>, roots<sup>d</sup> [P- phosphorus, S- sulphur, Cl- chlorine, K- potassium, Ca- calcium, Cr - chromium , Mn- manganese, Fe- Iron, Co- cobalt, Ni- Nickel, *Gomphrena* species are employed in folk medicine for the treatment of several diseases and for their nutritive value. The species with biological activities [Table 2],

Cu- copper, Zn- Zinc, Br- bromine, Sr- strontium, Cd- cadmium, Ba- barium, Pb- Lead, Sc- scandium, Ti- Titanium, V- vanadium].


including antimicrobial activity, used to treat gastrointestinal and respiratory disorders as well as infectious diseases were listed [10].

**Table 2: Main Biological Activities Reported for Species from *Gomphrena* [10].**

Species	Biological Activity
<i>Gomphrena boliviana</i>	Antimicrobial activity; Anticancer properties; cytotoxic activity
<i>Gomphrena celosioides</i>	Antimicrobial activity (Bacteria, <i>Plasmodium falciparum</i> ); Anticancer properties; Analgesic activity
<i>Gomphrena globosa</i>	Antimicrobial activity; Anticancer properties; cytotoxic activity
<i>Gomphrena haenkeana</i>	Antimicrobial activity; Anticancer properties; cytotoxic activity
<i>Gomphrena macrocephala</i>	Molluscicidal ( <i>Biomphalaria glabrata</i> ) and cytotoxic against <i>Artemia salina</i>
<i>Gomphrena martiana</i>	Antimicrobial activity; Anticancer properties; cytotoxic activity
<i>Gomphrena meyeniana</i>	Antimicrobial activity; Anticancer properties; cytotoxic activity
<i>Gomphrena perennis</i>	Antimicrobial activity; Anticancer properties; cytotoxic activity
<i>Gomphrena pulchella</i>	Antimicrobial activity

The phytochemical screening of *G. globosa* ethanol extracts [Gg-EE] detected the presence of saponins, alkaloids, reducing sugars and coumarins [11]. Dosumu *et al.* [12] Isolated the bioactive compounds like saponins, steroids, amino acids, non-reducing sugars, phenols and flavonoids from the methanol extract of *G. celosioides*. Hamiduzzaman and Azam [13] isolated three compounds stigmasterol [1],  $\beta$ -

sitosterol [2] and isochavicine acid [3, 16] from the crude methanol extract of whole plant of *Gomphrena globosa* by a combination of column chromatography and preparative thin layer chromatography over silica gel. The structures of these compounds were determined by H NMR spectroscopic [Fig. 1].




**Figure 1: Compounds Extracted from *G. globosa*.**

The ethyl acetate and methanol extracts of *Gomphrena celosioides* was phytochemically examined by Dosumu *et al.* [4] and revealed the presence of secondary metabolites. These metabolites include alkaloids,

tannins, saponins, steroids, glycosides and reducing sugars. Babu *et al.* [17] isolated the compound oleuropein from *Gomphrena serrata* by subjecting it to exhaustive hot extraction with chloroform. Arginine-

betaxanthin as a novel betaxanthin in *Gomphrena globosa* inflorescences was detected by Biswas *et al.* [18] and yellow Ulluco, which has not been reported so far as a pigment that occurs naturally and he performed characterization based on UV-Vis and mass spectrophotometry [Fig. 2].



**Figure 2: Chemical structures and identification of Betalains [Betacyanins and Betaxanthins] from *G.globosa***

Dosumu *et al.* [12] tested the bioactivities of the *G. celosioides* extracts against organisms. The ethyl acetate and methanol extracts of the plant displayed inhibition activities on *Staphylococcus aureus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Escherichia coli* and *Salmonella typhi*. The methanol extract was active against *Candida albicans*, *Aspergillus niger* and *Trichophyton species* with diameter zones of inhibition between 14 and 20 mm. The fractionated methanol extract produced 3-[4-hydroxyphenyl] methylpropenoate with mild antimicrobial activity against the test microorganisms. The antihelmintic assay showed ethyl acetate and methanol extracts to be active against *Fasciola gigantica*, *Taenia solium* and *Pheretima pasthuma*. Ethyl acetate was the most toxic of the extracts causing paralysis of *Taenia solium* within 15 minutes and the death of *Fasciola gigantica* within 20 minutes of application. The brine shrimp assay gave an LC50 of 52.15 and 77.98  $\mu\text{g/ml}$  on hexane and methanol extracts respectively.

Ferreira *et al.* [20] found that the aerial parts contain sterol glucoside like

**BIOLOGICAL AND MEDICINAL USES OF Genus *Gomphrena***

The leaves and flowers of *G. globosa* is a folk remedy for oliguria, heat and empacho, hypertension, antimicrobial, antioxidant [1], cough, diabetes, hypertension, kidney problems, hoarseness, cough, bronchitis and other respiratory diseases, mainly as expectorant, reproductive problems. It also has significant cytotoxic and estrogenic activity [13, 11, and 16]. The different

Corke *et al.* [19] identified the betacyanins for eight amaranthine-types, six gomphrenin types, and two betanin-types from Amaranthaceae by HPLC characterization of Reversed - Phase [RP] High-Performance Liquid Chromatography [HPLC].

Gomphsterol  $\beta$ -D-glucoside along with  $\beta$ -sitosterol, stigmasterol, campesterol, stigmasterol -  $\beta$ -glucoside, friedelin, 3-epi-friedelinol, allantoin and chrysoeriol - 7-O- $\beta$ -D-glucoside, gomphrenoside, hopane-7 $\beta$ -ol,  $\beta$ -sitosterol-3-O- $\beta$ -D-glucoside and 1-triacontanol [31]. Betalain, betacyanin and flavones were also isolated from the whole plant [21] and the flowers [22]. Kuroda *et al.* [23] screened the phytochemical constituent of *Gomphrena macrocephala* and has resulted in the isolation of two new oleananeglycosides (1 and 2) and a new taraxerane glycoside (3). The structures of 1-3 were determined as 11  $\alpha$ ,12  $\alpha$ -epoxy-3  $\beta$ -[[O- $\beta$ -D-glucuronopyranosyl]oxy]olean-28,13-olide [1], 11  $\alpha$ ,12  $\alpha$ -epoxy-3  $\beta$ -[[O- $\beta$ -D-galactopyranosyl-[1 $\rightarrow$ 3]-O- $\beta$ -D-glucopyranosyl-[1 $\rightarrow$ 2]]- $\beta$ -D-glucuronopyranosyl]-oxy]olean-28,13-olide [2], and 11  $\alpha$ ,12  $\alpha$ -epoxy-3  $\beta$ -[[O- $\beta$ -D-glucuronopyranosyl]oxy]taraxer-14-en-28-oic acid  $\beta$ -D-glucopyranosyl ester (3), respectively, on the basis of their spectroscopic data and the results of hydrolysis.

species of this genus also showed antibacterial, antimalarial, diuretic activities [24]. Besides as an ornamental plant *G.globosa* is also commonly used for the treatment of jaundice, high cholesterol and urinary problems in Latin America and Caribbean [25, 26]. Extracts of aerial parts of some species of the *Gomphrena* genus have the following biological activities reported: larvicide for *G. globosa*; antimicrobial for *G. martiana* and *G.*

*boliviana*, antitumor to *G. martiana* and estrogenic to *G. demissa* Mart. *Gomphrena celosioides* is used in ethnomedical practice in Nigeria for treatment of various skin diseases, worms' infections and infectious diseases.

In South America, the plant is utilized as an abortifacient. A decoction of the whole plant and a related species *Gomphrena globosa* is applied to gangrenous wound. *G. martiana* [27] and *G. boliviana* are employed as antimicrobial agents by the natives [12]. *Gomphrena globosa* Linn is applied to gangrenous wounds [4]. As a treatment of body sore, the leaves are crushed as a paste which is applied to the affected part and covered with the leaf of the same plant [28]. A number of Brazilian *Gomphrena* species are employed in the treatment of bronchial asthma, and fever, and as an analgesic, tonic, or carminative. This species show antimalarial and diuretic activities [17]. *Gomphrena globosa* flowers contain betacyanins which have potential as food colorant, antioxidants and in prostate problems [29]. Ethnic tribes, farmers, fishers and hunters observe small size of lower leaves with dark colour of *G. globosa* to predict the weather forecasting which indicates the adverse weather condition (typhoon or flood) [30, 31].

#### **Pharmacological Actions of *Gomphrena* Species**

##### **Anti-Inflammation**

Andrade *et al.* [8] determined the 24 phenolic compounds and eight betacyanins by HPLC-DAD in three different extracts of *G. globosa* inflorescences. The decoction presented the highest amount of phenolic compounds, kaempferol-3-O-[6-rhamnosyl] hexoside and kaempferol-3-O-hexoside being the main compounds. The rich betacyanins extract showed isogomphrenin III and gomphrenin III as major metabolites. The data provided the evidence that *G. globosa* inflorescences has the potential as a source of anti-inflammatory compounds, with relevance for the treatment of acute or chronic inflammatory conditions, and health-promoting antioxidants for use by both food and pharmaceutical industries. *Gomphrena vaga* Mart. entire plant is used as anti-inflammatory of ovaries. And it is drunk "as water" until symptoms disappear.

Its root is used as analgesic for toothache and leaves is used as an infusion against asthmas and allergies [15].

##### **Antimicrobial, Antioxidant and Cytotoxic Activity**

Hamiduzzaman and Azam [13] subjected the crude methanol extract and its hexane, carbon tetrachloride, chloroform and aqueous soluble fractions of *Gomphrena globosa* to antimicrobial, antioxidant [32] and brine shrimp lethality bioassays. Among all extractives the hexane soluble fraction exhibited the highest antioxidant activity having  $IC_{50}$  of  $13.17 \pm 0.308 \mu\text{g/ml}$  and the crude methanol extract showed significant antioxidant activity with  $IC_{50}$  of  $20.35 \pm 0.360 \mu\text{g/ml}$ . The chloroform soluble materials revealed significant cytotoxicity with  $IC_{50}$  of  $0.331 \pm 0.029 \mu\text{g/ml}$ . The carbon tetrachloride and chloroform soluble partitionates demonstrated mild to moderate antimicrobial activity with zone of inhibition  $8 \pm 0.208$  to  $14 \pm 0.069$  mm.

The methanolic extracts of leaf, stem and root of *G. celosioides* was investigated by Sharma and Vijayrergia [24] for anti-bacterial activity and was found to be most effective against the three tested bacteria [*Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Escherichia coli*]. The stem and leaf extract were found to have maximum zone of inhibition against *Escherichia coli* with 15.5mm and 16mm, while the minimum zone of inhibition against *Staphylococcus aureus* was 10mm and 12.5 mm. The root extract was found maximum zone of inhibition against *Pseudomonas aeruginosa* with 12mm. The maximum activity index for the stem and leaf extracts was 1.19 and 1.23 for *Escherichia coli* and the root and leaf extracts showed minimum activity index of 0.72 against *Pseudomonas aeruginosa*. Abalaka and Damisa [14] carried out the antifungal activities for *Gomphrena celosioides* extracts at different concentrations on *Aspergillus niger*, *Candida albicans* and *Trichophyton rubrum*.

The methanol extract of *G. celosioides* had a fungicidal effect on the selected fungal isolates at a concentration of  $2000 \mu\text{g/ml}$ . The minimum inhibitory concentration [fungistatic] was ranged from  $2000 \mu\text{g/ml}$  to  $1500 \mu\text{g/ml}$ . The cytotoxic activity of 65

tested extracts was studied by Mahmoud *et al.* [33] from eight plant species [*Tabebuia heptaphylla* [Vell.] Toledo, Bignoniaceae, *Tapirira guianensis* Aubl., Anacardiaceae, *Myracrodruon urundeuva* Allemão, Anacardiaceae, *Schinus terebinthifolius* Raddi, Anacardiaceae, *Gomphrena elegans* Mart., Amaranthaceae, *Attalea phalerata* Mart. ex Spreng., Arecaceae, *Eugenia uniflora* L., Myrtaceae, and *Annona dioica* A. St.-Hil., Annonaceae], *Gomphrena elegans* Mart., : n-hexane and n-butanol fraction of the leaves showed considerable activity.

Pomilio *et al.* [34] determined the antimicrobial activity of extracts and constituents of *Gomphrena martiana* [35] and *Gomphrena boliviana* [Amaranthaceae] in order to identify the compounds responsible for the folk-medicinal use of these plants. Each extract was evaluated against 20 microorganisms, including Gram-positive and Gram-negative bacteria, spore-forming Gram-positive bacteria, an acid-fast bacterium, a fungus and two yeasts. Fractionation of each with petroleum ether [PE] extract yielded five 5, 6, 7-trisubstituted flavones that were separately tested showing highest activity against *M. phlei* with minimum inhibitory concentration of 15, 20 and 75 mg/ml, approaching that of commercial bactericides.

#### **Natural Blood Coagulant**

The leaves of *G. globosa* act as a natural blood coagulatory. It stops bleeding due to cut injury is of immense importance among the ethnic people. The ethanolic extract of four medicinal plants *Polygonum microcephalum*, *Moringa oleifera*, *Croton tiglium* and *Gomphrena globosa* were examined for antioxidant activity, phenol and flavonoid content. Total phenol and flavonoid content and DPPH radical scavenging activity of the extracts were spectrophotometrically determined. The antioxidant activity was found to be higher in *Polygonum microcephalum*, *Moringa oleifera* and *Gomphrena globosa* [1].

#### **Cardiovascular Effects**

*Gomphrena globosa* is used in the treatment of high blood pressure. It promoted a hypotensive activity by significant reduction in arterial blood pressure without change in heart rate, confirming the

therapeutic use as antihypertensive for this plant [11, 15]. They also prepared the ethanol extract from leaves of *G. globosa* L. by maceration and analyzed the phytochemical screening and biological assays. The phytochemical screening detected the presence of saponins, alkaloids, reducing sugars and coumarins.

#### **Cooling and Cough**

In Trinidad, the flowers of *Gomphrena* are boiled to make a tea which is used for baby gripe, oliguria, cough, diabetes and cooling.

#### **Food Colourant**

The fresh crude extract samples *Gomphrena globosa* without any purification was used by Corke *et al.* [19] for MALDI-QIT-TOF MS analysis using 2, 5-dihydroxybenzoic acid as a matrix. Fourteen free and acylated betacyanins, belonging to amaranthin-type, betanin-type and gomphrenin-type betacyanins, respectively, were identified. Lans [29] confirmed that *Gomphrena globosa* flowers contain betacyanins which have potential as food colorants and antioxidants.

#### **Anti-Microbial Activity**

Dias *et al.* [6] screened the ethanolic extract and pure compounds of *Gomphrena celosioides* for antimicrobial activity by Kirby-Bauer method. Quantitative determination of 4-hydroxy-3-methoxybenzoic acid in stems, leaves, flowers and roots was established by TLC-densitometry. Their results showed significant activity against *Staphylococcus aureus* and *Salmonella typhi*. Dosumu *et al.* [4] found that ethyl acetate and methanol extracts of *G. celosioides* exhibited anthelmintic activities against *Pheretima posthuma*, *Fasciola gigantica* and *Taenia solium*. Higher anthelmintic and antibacterial activities were displayed in ethylacetate extract. Methanol extract exhibited pronounced antifungal activity. Preliminary phytochemical analysis of the extracts indicated the presence of steroids, glycosides, alkaloids, saponins and tannins. The ethyl acetate and methanol extracts of *G. celosioides* was tested by Dosumu *et al.* [12] which showed inhibition activities on *Staphylococcus aureus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Escherichia coli* and *Salmonella typhi*. Methanol extract was active against *Candida albicans*, *Aspergillus*

*niger* and *Trichophyton species* with diameter zones of inhibition between 14 and 20 mm.

#### **Anti-Cancer Agent**

The phytochemical analysis of chloroform extract of *Gomphrena serrata* was studied by Babu *et al.* [17] and reported the presence of carbohydrates, glycosides, aminoacids, phytosterols, flavonoids, phenolics and terpenoids. They isolated the compound oleuropein from *Gomphrena serrata*. The development of novel oleuropein as an anti cancer agents and in-silico docking or computational studies are in progress.

Latha *et al.* [37] performed a study to investigate the anticancer activity of chloroform extract of aerial parts of *Gomphrena globosa* against Ehrlich Ascites Carcinoma [EAC] induced solid tumor. This study showed that, the extract was non-toxic up to 2000 mg / kg body wt. The histological study indicated that the damaged tissue was recovered by extract and solid tumor volume was reduced significantly. Thus, the findings revealed a significant anti-cancer activity of the extract

#### **Reproductive Problems**

*Gomphrena globosa* are used for prostate and reproductive problems [29]. The root decoction of *Gomphrena demissa* Mart. in a liter of water is used for the treatment of female sterility, amenorrhea, inflammations and ovarian diseases [15].

#### **Organic Sulfur Fertilizer**

Wang *et al.* [38] explored that *G. globosa* absorb atmospheric sulfides, which may be of great importance for ameliorating the environment and for farming as a green organic sulphur fertilizer used to balance insufficient soil sulphur content for intensive cultivation in China; H<sub>2</sub>S and mainly SO<sub>2</sub> are emitted to air as a result of the rapid industrialized and economic development.

#### **Urinary Problems**

A non-experimental validation was conducted by Lans [26] on the plants used for urinary problems and diabetes mellitus. The *G.globosa* plant is used for hypertension, jaundice that may be safe. [39] Conducted the survey of ethnomedicinal plants in the muzaffarnagar district of Uttarpradesh to assess the

potentiality of plant resources. The study revealed that 15 plant species belonging to 13 families are used as anti-urolithiatic agents in local remedies. *G.celosioides* Mart. is most effective and commonly used in treatment of urinary tract and kidney stones.

#### **Anti-Malarial Activity**

*Gomphrena arborescens* Mart. leaves, flowers and tuberous root is used as anti-malarial activity which is introduced by Brazilian priests that learned it from Indians [40].

#### **Analgesic Activity**

Hamiduzzaman [41] investigated the central and peripheral analgesic activity of *Gomphrena globosa* for the wound healing activity to the tribal people of Bangladesh. The fractions like n-hexane soluble fraction, carbon tetrachloride soluble fraction, chloroform soluble fraction, aqueous soluble fraction were investigated for analgesic activity. The crude methanolic extract and n-hexane soluble fraction have significant analgesic activity at 400mg/kg dose. The central analgesic activity was highest after 30 minutes. After 60 minutes there was no central analgesic activity in the plant extracts.

The anti-inflammatory and analgesic properties of aqueous leaf extracts of *Gomphrena celosioides* and *Momordica charantia* in rats and mice were reported by Oladele *et al.* [42]. The result of the study revealed that the leaf extracts of the two plants possess anti-inflammatory property as they were found to significantly [ $p<0.05$ ] inhibit oedema induced by carrageenan in the rat paws. The results of the study suggest the anti-inflammatory and analgesic effects of the aqueous leaf extracts of the two plants.

#### **CONCLUSION**

The studies on *Gomphrena* species elaborate the biological and medicinal applications in various ways from the ancient time have been discussed in the present review. It has been used throughout India not only as an ornamental source but also as a natural remedy for treatment of kidney failure and reproductive problems, cough, etc., On the basis of the above biological screenings and folk medicinal uses it can be concluded that this plant has

significant medicinal properties and hence it can be carried out to isolate and characterise the active compounds responsible for this activities in the plant is recommended. The present knowledge on medicinal uses of plants needs scientific

#### REFERENCES

1. Upadhyaya S and Saikia LR. Antioxidant Activity, Phenol and Flavonoid Content of Some less Known Medicinal Plants of Assam. International Journal of Pharma and Bio Sciences 2011; 2(2): 383-388.
2. Ganesan S, PonnuChamy M, Kesavan L and Selvaraj A. Floristic composition and practices on the sacred groves of pallapatty village (Reserved forest), Tamilnadu. Indian Journal of Traditional Knowledge 2009; 8(2): 154-162.
3. Joy PP, Thomas J, Mathew S and Skaria BP. Medicinal Plants. Tropical Horticulture Vol. 2. (Eds. Bose, T.K., Kabir, J., Das, P. and Joy, P.P.). Naya Prokash, Calcutta, pp. 449-632. 2001.
4. Dosumu OO, Onocha PA, Ajaiyeoba EA and Ekundayo O. Phytochemical Screening and Biological Activities of *Gomphrena celosioides* (C. Mart) Extracts. Nigerian Society for Experimental Biology 2005; 5(2): 61-67.
5. Ignacimuthu S and Ayyanar M. Plants used for non-medicinal purposes by the tribal people in kalakad Mundanthurai Tiger reserve, Southern India. Indian Journal Traditional of Knowledge 2009; 9(3): 515-518.
6. Dias DA, de Moura RMX, Pereira PS, Januário AH and França SC. Antimicrobial Screening and Quantitative Determination of Benzoic Acid Derivative of *Gomphrena celosioides* by TLC-Densitometry. Chem. Pharm. Bull. 2004; 52(11): 1342—1344.
7. Bão SN, Fank-de-Carvalho SM, Gomes MRA and Silva PIT. Leaf surfaces of *Gomphrena* species (Amaranthaceae) from *Cerrado biome*. Biocell 2010; 34(1): 23-35.
8. Andrade PB, Silva LR, Valentão P, Faria J, Ferreres F, Sousa C, Gil-Izquierdo and Pinho BR. Phytochemical investigations and biological potential screening with cellular and non-cellular models of globe amaranth (*Gomphrena globosa* L.) inflorescences. Food Chem. 2012; 135(2): 756-63.
9. Wang MY, Wu LH and Zhang J. Impacts of root sulfate deprivation on growth and elements concentration of globe amaranth (*Gomphrena globosa* L.) under hydroponic condition. Plant Soil Environ 2009; 55(11): 484-493.

investigation to confirm their medicinal values.

#### COMPETING INTERESTS

This review article has no competing interests.

10. Salvador MJ, Andrezza NL, Pascoal ACRF, Pereira PS, França SC, Zucchi and Dias DA. OLAD Bioactive Chemical Constituents and Biotechnological Production of Secondary Metabolites in Amaranthaceae Plants, Gomphreneae Tribe. Biotechnological Production of Plant Secondary Metabolites. 2012; 124-158.
11. Arcanjo DDR, Albuquerque ACM, Neto BM, Santana LCLR, Silva NCB, Moita MM, Medeiros MGF, Soares MJS and Cito AMGL. Phytochemical screening and evaluation of cytotoxic, antimicrobial and cardiovascular effects of *Gomphrena globosa* L. (Amaranthaceae). Journal of Medicinal Plants Research 2011; 5(10): 2006-2010.
12. Dosumu AA, Idowu PA, Onocha PA and Ekundayo O. Isolation Of 3-(4-Hydroxyphenyl) Methylpropenoate and Bioactivity Evaluation of *Gomphrena celosioides* Extracts. EXCLI Journal 2010; 9: 173-180.
13. Hamiduzzaman M and Azam ATMZ. Antimicrobial, Antioxidant and Cytotoxic Activities of *Gomphrena globosa* (L.). Bangladesh Pharmaceutical Journal 2012; 15(2): 183-185.
14. Abalaka ME and Damisa D. Antifungal activity of *Gomphrena celosioides* (soft khaki weed) on selected fungal isolates. J of Current research in science 2013; 1(2): 66-70.
15. Agra FM, Freitas PF and Barbosa-Filho JM. Synopsis of the plants known as medicinal and poisonous in Northeast of Brazil. Brazilian Journal of Pharmacognosy 2007; 17(1): 114-140.
16. Azam ATMZ, Hamiduzzaman M and Hasan CM. Isochavacinic Acid and Steroids from *Gomphrena globosa* L. Dhaka Univ. J. Pharm. Sci. 2012; 11(1): 79-81.
17. Babu G, Anju P, Biju CR and Rajapandi R. Phytochemical screening of *Gomphrena serrata* L. Journal of Chemical and Pharmaceutical Research 2012; 4(7) 3396-3399.
18. Biswas M, Dey S and Sen R. Betalains from *Amaranthus tricolor* L. J of Pharmacognosy and Phytochemistry 2013; 1(5): 88-96.
19. Corke H, Cai Y and Sun M. Characterisation of betalains from plants in the



- Amaranthaceae. J. of Chromatographic Science 2005; 43: 454-460.
20. Ferreira EQ, Salvadov MJ, Pral EMF, Alfieri SC, Ito IY and Dias DA. A new hwppta substituted (E)-Aurone glucoside and other aromatic compounds of *Gomphrena agrestis* with biological activity. Z. Naturforsch. 2004; 59c: 499-505.
  21. Liu Chang T and Chu C. Expectorant principle from *Gomphrena globosa* - structure determination of a new flavone glycoside. Yao Hsush Tung Pao 1981; 1(16): 55 - 6.
  22. Heuer S, Wray V, Metzger JW and Strack D. Betacyanins from flowers of *Gomphrena globosa*. Phytochemistry 1992; 31(5): 1801-1807.
  23. Kuroda M, Aoshima T, Haraguchi M, Young MC, Sakagani H and Mimaki Y. Oleanane and taraxerane glycosides from the roots of *Gomphrena macrocephala*. J of Nat. Prod 2006; 69(11): 1606-1610.
  24. Sharma N and Vijayrergia R. Study of primary metabolites and antimicrobial activity of *Gomphrena celosioides* Mart. Intern J of Pharma and Bioscience 2011; 2(4): 581-586.
  25. Hasnain S, Ghaffar I and Ali B. Effect of different hormonal combinations on regeneration of callus of *Gomphrena globosa* L. Pakistan Journal of Biological Science 2007; 10(20): 3708-3712.
  26. Lans CA. Ethnomedicines used in Trinidad and Tobago for urinary problems and diabetes mellitus. Journal of Ethnobiology and Ethnomedicine 2006; 2: 45.
  27. Buschi CA, Pomilio AB and Gros EG. New methylated flavones from *Gomphrena martiana*. Phytochemistry 1980; n19(5): 903-904.
  28. Bhat RB, Adeloye AA and Etejere EO. Some medicinal plants of Nigeria. J. Econ. Tax. Bot. 1985; 6(1): 161-165.
  29. Lans C. Ethnomedicines used in Trinidad and Tobago for reproductive problems. Journal of Ethnobiology and Ethnomedicine 2007; 3: 13.
  30. Archarya S. Presage Biology: Lessons from nature in weather forecasting. Indian Journal of Traditional Knowledge 2011; 10(1): 114-124.
  31. Biswanath D, Ghosh B, Achari B, Arima S, Sato N and Harigaya Y. Chemical Constituents of *Gomphrena globosa*. Natural Product Sciences 2006; 2(12): 89 - 93.
  32. Kyung-Soo C, Yu-Su S, Hyun-Jung J, Sang-Won L, Young-Ock K and Yoon-Pyo H. Analysis of anti-oxidant activity of medicinal plants according to the extracted parts. Journal of Medicinal Plants Research 2012; 6(31): 615-4624.
  33. Mahmoud TS, Marques MR, do O Pessoa, lo tufo LVC, Magalhaes HIF, de Moraes MO, de Lima DP, Tininis AG and de Oliveira JE. *In vitro* cytotoxic activity of Brazilian Middle West plant extracts. Brazilian J of Pharmacognosy 2011; 21(3): 456-464.
  34. Pomilio AB, Buschi CA, Tomes CN and Viale AA. Antimicrobial constituent of *Gomphrena martiana* and *Gomphrena boliviana*. J. Ethnopharmacology 1992; 36(2): 155-161.
  35. Pomilio AB, Sola GA, Mayer AM and Rumi LS. Antitumor and cytotoxic screening of 5, 6, 7-trisubstituted flavones from *Gomphrena martiana*. Z. Naturforsch 2004; 59(7-8): 499-505.
  36. Corke H, Yi-Zhong C, Xing J and Sun M. Rapid Identification of Betacyanins from *Amaranthus tricolor*, *Gomphrena globosa*, and *Hylocereus polyrhizus* by Matrix-Assisted Laser Desorption/Ionization Quadrupole Ion Trap Time-of-Flight Mass Spectrometry (MALDI-QIT-TOF MS). J. Agric. Food Chem. 2006; 54 (18): 6520-6526.
  37. Latha ST, Rajendran NN and Babu G. Anticancer screening of *Gomphrena globosa* against Ehrlich ascites carcinoma in Swiss albino mice. J of Chemical and Pharmaceutical Research. 2013; 5(2): 283-289.
  38. Wang MY, Wu LH and Zhang J. Impacts of root sulfate deprivation on growth and elements concentration of globe amaranth (*Gomphrena globosa* L.) under hydroponic condition. Plant Soil Environ 2009; 55(11): 484-493.
  39. Kasana MS, Prachi, Chauhan N and Kumar D. Medicinal plants of Muzaffarnagar District used in treatment of urinary tract and kidney stones. Indian J of Traditional Knowledge. 2009; 8(2): 191-195.
  40. Botsaris AS. Plants used traditionally to treat malaria in Brazil: the archives of Flora Medicinal. Journal of Ethnobiology and Ethnomedicine 2007; 3(18): 1-8.
  41. Hamiduzzaman M. Evaluation of central and peripheral analgesic activity of whole plant *Gomphrena globosa* (L) (Family: Amaranthaceae). International research Journal of Pharmacy 2013; 4(6): 54-57.
  42. Oladele GM, Abatan MO, Olukunle JO and Okediran BS. Anti-inflammatory and analgesic effects of Aqueous leaf extracts of *Gomphrena celosioides* and *Momordica charantia*. J. of Natural Sciences, Engineering and Technology 2009; 8(2): 1-8