



## Diabetes and Depression

## Faraz Farishta

Endocrine & Diabetes Center, India

## **Abstract:**

In Nepal, postpartum depression affects 1 out of 10 women, and suicide is the leading cause of death among women of reproductive age. Passive sensing technology is a way to collect data on the behaviors and activities of depressed mothers in order to better tailor psychological treatment and improve outcomes for postpartum depression. This study investigated (a) the feasibility and acceptability of wearable digital sensors with adolescent mothers and their families in rural Nepal and (b) the feasibility and utility of implementing this data into a phonebased application used by non-specialists to provide personalized psychological treatment. This study used a mobile phone and Bluetooth device to generate passive sensing data on aspects of a mother's life, such as amount of time spent with and away from the baby, movements and activities both inside and outside of the house, social interactions experienced, and physical activity. We then interviewed both depressed and non-depressed adolescent mothers who used these wearable digital devices and analyzed the ethicality, safety, social acceptability, utility, and feasibility of these technologies. This data was then used to develop StandStrong, a platform that collects passive sensing data to implement personalized depression treatment. Our results showed that both depressed and non-depressed mothers found it acceptable and feasible to collect passive sensing data. Depressed and non-depressed mothers expressed utility in having knowledge of their own movements and activities, as well as having information about their proximity to and interactions with their child. The non-specialized community counselors expressed utility in using the data collected from



the wearable digital sensors to encourage behavioral changes during sessions and also to track the progress of patients between sessions. Barriers to using wearable digital devices included difficulty carrying the phone around throughout the day, privacy concerns, fear of loss or damage to the device, and concern about possible adverse health effects of the device. In summary, it is feasible and acceptable to use passive sensing data to tailor psychological treatment for depressed adolescent mothers in low-resource settings. This research demonstrates the effectiveness of mobile health technology in improving treatment and outcomes for postpartum depression in rural areas.

## Biography:

Faraz Farishta is working at Endocrine & Diabetes Center, India.

Webinar on Stress and Depression Management; Barcelona, Spain; October 15, 2020

Citation: Faraz Farishta; Diabetes and Depression; Webinar on Stress Management; Barcelona, Spain; October 15, 2020

Res & Rev: Neurosci 2020 Volume: and Issue: S(1)