Research and Reviews: Journal of Agriculture and Allied Sciences
e-ISSN: 2347-226X
p-ISSN: 2319-9857

Effect of Sulphur and Sowing Date on Soil Fertility, Growth, and Yield of Rapeseeds (*Brassica napus* L.)

Kushal Shrestha*

Department of Medicine, Nepalese Army Institute of Health Sciences, Kathmandu, Nepal

Research Article

Received: 20-Feb-2025, Manuscript No. JAAS-25-161529; **Editor assigned:** 25-Feb-2025, PreQC No. JAAS-25-161529 (PQ); **Reviewed:** 11-Mar-2025, QC No. JAAS-25-161529; **Revised:** 20-Feb-2026, Manuscript No. JAAS-25-161529 (R); **Published:** 27-Feb-2026, DOI: 10.4172/jaas.15.01

***For Correspondence:**

Kushal Shrestha, Department of Agriculture, Kathmandu University, Dhulikhel, Nepal,

E-mail: kushalshrestha851@gmail.com

Citation: Shrestha K. Effect of Sulphur and Sowing Date on Soil Fertility, Growth, and Yield of Rapeseeds (*Brassica napus* L.). J Agri Allied Sci. 2026;15:001.

Copyright: © 2026 Shrestha K. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Rapeseed (*Brassica napus* L.) is one of the most important oil seed crops globally, contributing significantly to edible oil production and animal fodder. The present study, conducted during the Rabi season of 2022-23 at the Institute of Agriculture Training & Research, Prem Nagar, Dehradun, Uttarakhand, aimed to assess the impact of different sulfur doses and sowing dates on soil fertility, growth, and yield of rapeseed. The research evaluated key agronomic parameters, including plant height, number of leaves, seed yield, and soil nutrient status. Findings revealed that sulphur application positively influenced crop performance, enhancing biomass production and oil content. Similarly, the optimal sowing date was crucial in ensuring maximum yield by mitigating temperature stress and synchronizing the reproductive phase with favorable climatic conditions. The study underscores the significance of sulphur nutrition and appropriate sowing schedules in improving rapeseed productivity, thus contributing to sustainable agricultural practices.

Keywords: Rapeseed; Seed yield; Soil nutrient; Sulphur

INTRODUCTION

India is a leading global producer of various agricultural commodities, ranking first in the production of spices, milk, and pulses, and holding significant positions in paddy, wheat, and cotton cultivation. The country benefits from diverse climatic conditions, encompassing 15 agro-climatic zones and 20 agro-ecological zones. Among oilseed crops, rapeseed is the third most important oilseed globally, following palm oil and soybean, with major producers including India, Canada, China, and Nepal. Rapeseed (*Brassica napus* L.), commonly referred to as sarson, tori, or bird rape, belongs to the mustard family *Brassicaceae* and is valued for its edible oil, leafy vegetables, and use as animal feed. The crop is a significant component of India's agricultural economy, with its oil widely used in cooking, pickling, medicinal applications, and industrial processes such as leather tanning and grease production. Post-extraction, the residual oil cake is a nutrient-rich animal feed and organic fertilizer [1].

The growth and productivity of rapeseed are highly influenced by climatic conditions, soil type, and agronomic practices. The crop thrives in loam soils with a neutral pH and requires well-drained conditions for optimal growth. It is susceptible to water stress before flowering, necessitating careful water management. Additionally, temperature fluctuations, sowing time, and nutrient availability, particularly sulphur, play a crucial role in determining yield and oil content. Sulfur is an essential nutrient for rapeseed, contributing to enzyme activation, protein synthesis, and oil formation. Both organic and inorganic sulphur sources impact plant metabolism differently, with organic sulphur improving soil health over time and inorganic sulphur offering rapid nutrient availability. Given the varying agroecological conditions across India, the requirement for sulfur application is location-specific [2].

Time of sowing is another critical factor influencing rapeseed yield. Delayed sowing exposes the crop to unfavorable weather conditions, reducing seed yield due to shorter silique filling periods and increased drought stress. Selecting an optimal sowing date can enhance growth parameters, including plant height, leaf area, and biomass accumulation, thereby improving overall productivity.

Objectives

- To evaluate the effect of sulphur application on rapeseed growth and yield.
- To assess the impact of different sowing dates on crop performance.
- To analyze the interaction between sulphur levels and sowing dates on yield and soil fertility.
- To determine changes in soil nutrient status under varied sulphur and sowing date treatments.

MATERIALS AND METHODS

The present study, entitled "Effect of sulphur and sowing dates on soil fertility, growth, and yield of rapeseed (*Brassica napus*)," was conducted during the 2022-23 growing season at the experimental field of the Institute of Agriculture Training & Research, Prem Nagar, Dehradun, Uttarakhand, India. This section outlines the experimental site, climatic conditions, soil properties, field history, and experimental design [3].

Experimental site

The experiment was carried out at the Institute of Agriculture Training & Research, Prem Nagar, Dehradun, Uttarakhand (248197), India, situated at 30.33° N latitude, 77.97° E longitude, and an altitude of 410 meters above sea level.

Weather and climatic conditions

The region experiences a sub-tropical climate, characterized by hot summers, cold winters, and monsoonal rainfall. The average annual rainfall is approximately 1200 mm, with about 80% occurring between mid-June and September. Meteorological data, including temperature, relative humidity, and rainfall, were recorded at the Meteorological Observatory in Selaqui, Dehradun, during the cropping season [4].

Soil properties

The soil of the experimental field was sandy loam in texture. Composite soil samples were collected from a 30 cm depth before field preparation and analyzed for organic carbon, available nitrogen, phosphorus, potassium, pH, and electrical conductivity using standard procedures. The soil had an organic carbon content of 2.25%, available nitrogen of 348 kg/ha, phosphorus of 144 kg/ha, and potassium of 212 kg/ha. The pH was recorded at 7.40 with an electrical conductivity of 0.138 ds/m.

Previous cropping history

The field followed a rotational cropping sequence:

- **2020-21:** Maize (Kharif) - Pea (Rabi)
- **2021-22:** Maize (Kharif) - Rapeseed (Rabi)
- **2022-23:** Rice (Kharif) - Rapeseed (Rabi)

Experimental design and treatments

The experiment was laid out in a Randomized Block Design (RBD) with three replications and 20 treatment combinations. The treatments consisted of four sowing dates (S1: 25 August, S2: 5 September, S3: 15 September, S4: 25 September) and five Sulphur levels (F1: Control, F2: NPK only, F3: NPK + 20 kg S/ha, F4: NPK + 40 kg S/ha, F5: NPK + 60 kg S/ha) [5].

Field preparation and crop management

The experimental field was plowed, leveled, and divided into plots according to the treatment plan. The rapeseed variety Pant Toria-508 was used. Fertilizers were applied as per the treatments, and standard agronomic practices were followed for crop management, including irrigation, weeding, and pest control.

Data collection and analysis

Observations were recorded on soil fertility parameters, growth attributes (plant height, number of branches), and yield components (pods per plant, seed yield, and oil content), and analyzed statistically using ANOVA to determine treatment effects [6].

This methodology ensured a systematic approach to studying the influence of sulphur doses and sowing dates on rapeseed growth and yield.


RESULTS

Pre-harvest observations

Germination count:

- **Sulphur management:** Maximum germination count was observed with F2 (NPK only) and F3 (NPK + 20 kg S/ha). F1 (control) had the lowest germination.
- **Sowing date:** S4 (25th September) recorded the highest germination count, followed by S3 (15th September). S1 (25th August) had the lowest germination.
- **Interaction:** F2S4 (NPK + 25th September sowing) showed the highest germination count (7.39).

Figure 1. Germination count (No.) observed under different treatment combinations (F1S1–F5S4) (Figure 1).



Plant population:


- **Sulphur management:** F2 (NPK only) had the highest plant population, while F1 (control) had the lowest.
- **Sowing date:** S4 (25th September) recorded the highest plant population, with no significant variation among

e-ISSN: 2347-226X
p-ISSN: 2319-9857

other dates [7].

- **Interaction:** F2S4 (NPK + 25th September sowing) showed the highest plant population (7.45) (Figure 2).


Figure 2. Plant population recorded under different treatment combinations (F1S1–F5S4).



Plant height:

- **Sulphur management:** F2 (NPK only) resulted in the tallest plants, while F1 (control) had the shortest.
- **Sowing Date:** S4 (25th September) recorded the tallest plants at 25, 50, and 75 DAS, followed by S3 (15th September) [8].
- **Interaction:** F2S4 (NPK + 25th September sowing) showed the maximum plant height (Figure 3).


Figure 3. Plant height (cm) recorded at 25, 50, and 75 Days after Sowing (DAS) under different treatment combinations.



Root length:

- **Sulphur management:** F2 (NPK only) resulted in the longest roots, while F1 (control) had the shortest.
- **Sowing date:** S4 (25th September) recorded the longest roots at 25, 50, and 75 DAS.
- **Interaction:** F2S4 (NPK + 25th September sowing) showed the maximum root length (Figure 4).


Figure 4. Effect of different treatment combinations on root length (cm) at 25, 50, and 75 Days after Sowing (DAS).



Dry matter accumulation:

- **Sulphur management:** F2 (NPK only) resulted in the highest dry matter accumulation, while F1 (control) had the lowest.
- **Sowing date:** S4 (25th September) recorded the highest dry matter accumulation at 25, 50, and 75 DAS (Figure 5).
- **Interaction:** F2S4 (NPK + 25th September sowing) showed the maximum dry matter accumulation [9].

Figure 5. Dry matter accumulation (g) under different treatment combinations (F1–F5) at 25, 50, and 75 Days after Sowing (DAS).



Yield attributes

Number of seeds per silique and test weight:

- **Sulphur management:** F2 (NPK only) resulted in the highest number of seeds per silique and test weight, while F1 (control) had the lowest.

- **Sowing date:** S4 (25th September) recorded the highest number of seeds per silique and test weight.
- **Interaction:** F2S4 (NPK + 25th September sowing) showed the maximum number of seeds per silique and test weight.

Grain yield, stover yield, harvest index, and biological yield:

- **Sulphur management:** F2 (NPK only) resulted in the highest grain yield, stover yield, harvest index, and biological yield, while F1 (control) had the lowest.
- **Sowing date:** S4 (25th September) recorded the highest grain yield, stover yield, harvest index, and biological yield.
- **Interaction:** F2S4 (NPK + 25th September sowing) showed the maximum grain yield, stover yield, harvest index, and biological yield.

Nutrient uptake

Nitrogen, phosphorus, and sulphur uptake:

- **Sulphur management:** F2 (NPK only) resulted in the highest nutrient uptake, while F1 (control) had the lowest.
- **Sowing date:** S4 (25th September) recorded the highest nutrient uptake.
- **Interaction:** F2S4 (NPK + 25th September sowing) showed the maximum nutrient uptake.

NPK fertilization (F2) and sowing on 25th September (S4) resulted in the best growth, yield, and nutrient uptake for rapeseeds. The F2S4 interaction (NPK + 25th Sept sowing) showed maximum germination, plant population, height, root length, dry matter accumulation, yield attributes, and nutrient uptake. This combination consistently outperformed others across all parameters.

DISCUSSION

The study titled "Effect of Sulphur and Sowing Date on Soil Fertility, Growth, and Yield of Rapeseeds (*Brassica napus* L.)" conducted during the 2022-23 Rabi season in Dehradun, India, explored the impact of various sulphur management strategies and sowing dates on rapeseed growth and yield. The findings indicated that the weather conditions, particularly temperature, relative humidity, and rainfall, were favorable for rapeseed growth, with no significant deviations from normal. Among the sulphur management treatments, the application of NPK fertilizers (F2) led to the highest germination count, plant population, and growth attributes, including plant height, root length, and dry weight. This could be attributed to better sulphur uptake, which promoted better plant establishment and growth. The sowing date also had a substantial effect, with the 25th September sowing (S4) showing the best results in terms of germination, plant height, yield attributes, and nutrient uptake.

CONCLUSION

The study further revealed that the optimal sulphur management and sowing date significantly influenced rapeseed yield. The highest grain yield, straw yield, and harvest index were observed under the F2 (NPK fertilizers) treatment, with the 25th September sowing (S4) also showing the highest yield outcomes. This can be attributed to improved nutrient uptake, better growth during the vegetative and reproductive stages, and optimal environmental conditions during the sowing period. Delayed or early sowing reduced crop performance, with early sowing being prone to temperature stress, while late sowing affected maturity and yield. The results highlighted the importance of both sulphur management and sowing date in optimizing rapeseed yield and nutrient uptake, with F2 and S4 being the most effective treatments.

REFERENCES

1. Alam M, et al. Effect of planting dates on the yield of mustard seed. Int J Appl Sci Biotechnol. 2015;3:651-654.
2. Aziz M, et al. Effect of sowing dates on the growth, development and yield of mustard in the hilly areas. J Exp Biosci. 2011;2:33-36.
3. Charak AS, et al. Growth and productivity of rapeseeds (*Brassica rapa* var. dichotoma) as influenced by sowing time, row spacing and nitrogen levels. Indian J Agric Sci. 2006;76:685-687.

e-ISSN: 2347-226X
p-ISSN: 2319-9857

4. Kumar A, et al. Effect of different sowing dates on yield and yield attributes of Indian mustard (*Brassica juncea* L.) genotypes. Int J Pure Appl Biosci. 2018;6:848-853.
5. Mitiku W, et al. Effect of integrated sulphur management on yield and yield components of barley (*Hordium vulgare* L.) in Keffa Zone, South Western Ethiopia. Sci Technol Arts Res J. 2014;3:34-42.
6. Yadav A, et al. Effect of planting geometry on growth and yield of mustard (*Brassica juncea* (L.)) varieties. J Pharmacog Phytochem. 2018;7:2624-2627.
7. Rai A, et al. Sulphur in soils and plants: An overview. Int Res J Pure Appl Chem. 2020;21:66-70.
8. Jamal A, et al. Sulphur-a general overview and interaction with nitrogen. Australian J Crop Sci. 2010;4:523-529.
9. Ram A, et al. Effect of sulphur on soil biological properties, residual fertility and yield of aerobic rice grown under aerobic rice-wheat cropping system in Inceptisols. J Environ Biol. 2017;38:587.