

Neuroprotective Effect of Barley Plant (*Hordeum Vulgare*) Against the Changes in MAO Induced By Lead and Cadmium Administration in Different CNS Regions of Male Guinea Pig.

Somaya M Ismail*, Hayat A Ismail² and Ghidaa M Al-Sharif².

¹Department of Zoology, Faculty of Science, Cairo University.

²Biology Department, faculty of science, King Abd El Aziz University.

Research Article

Received: 15/02/2014

Revised: 16/03/2014

Accepted: 25/03/2014

*For Correspondence

Department of Zoology,
Faculty of Science, Cairo
University.

Keywords: lead, cadmium,
MAO, barley, CNS

ABSTRACT

Lead and cadmium are highly toxic metals that can be ingested or inhaled from a variety of industrial and dietary sources. The purpose of this study was to determine the impact of chronic administration of lead and cadmium on the activity of monoamine oxidase (MAO) in different central nervous system regions (CNS) of male guinea pigs and the protective effect of barley (*Hordeum Vulgare*) plant. Five groups of animals were used in this experiment, first group of animals was subcutaneous injected at dose level (30mg/kg. body weight) lead acetate, the second group was injected with (1mg/kg. body weight) cadmium chloride, third group of animals received a mixed dose of the two heavy metals with the same previous dose, the fourth dose was received an aqueous solution of barley (5mg/kg body weight) for two weeks followed by a mixed dose of lead and cadmium (30mg/kg. body weight), (1mg/kg.). Animals were decapitated after two, four, six and eight weeks after injection. The activity of monoamine oxidase (MAO) was determined in the cerebral cortex, caudate putamen, thalamus, hypothalamus, superior colliculus, inferior colliculus, cerebellum, pons, medulla and spinal cord in male guinea pig CNS. Lead and cadmium induced a general change in the level of the membrane bound enzyme (MAO) in most CNS regions and in turn affecting both cholinergic and adrenergic neurotransmitters. These results suggest that lead and cadmium may exert neurotoxic effect by altering certain membrane bound enzymes and may cause oxidative stress that could lead to neurodegenerative diseases. Supplemented groups with barley exhibited a similar MAO to the control group, suggesting that barley protected the CNS from functional damage resulted from the heavy metals. Accordingly, these results indicate that barley supplementation in a mixed dose of lead and cadmium injected pigs normalized MAO in most CNS regions, i.e. is lead to a general improvement in MAO activity.

INTRODUCTION

An important aspect in the study of mammalian central nervous system (CNS) is measurement of the activity of enzymes associated with the CNS functions. From such enzymes the degrading enzyme MAO which is a monoamine metabolizing enzyme located in the outer mitochondrial membrane of many cells and responsible for oxidation deamination of number of biogenic amines (Gu et al, 2013).

Although a number of reports have appeared in the literature regarding the neurotoxic effect of lead and cadmium, relatively little is known about the effect on membrane bound enzyme and distribution of both metals in the central nervous system (Kim et al., 2013).

Lead and cadmium are the most studied neurotoxins and its effects on the nervous system are well documented (wright and Baccavell, 2007). Lead (Pb) is considered to be a multi-target toxicant, Pb-acetate induced toxic manifestation in blood, liver, kidney, brain and heart of Wistar rats (Ranabir et al., 2013).

Now it is widely accepted that even small quantities of lead and cadmium are harmful to human and other organisms (Sandra et al, 2008). Lead poisoning is most commonly caused by occupational exposure (Gupta et al., 2011).Neurotoxicity of lead and other heavy metals known to have multiple mechanisms of actions but these mechanisms are still not clearly understood. According to(Silbergeld ,1992) one of these mechanisms may be neurodevelopmental toxicity which includes interference with cell adhesion molecule, resulting in the misfiring of the CNS during early development and destruction of the blood –brain barrier which lead to permanent dysfunction ,edema loss of neurons and gliosis (Zheng et al., 2003). The other possible mechanism could be neuropharmacological toxicity that might involve interaction between the studied heavy metals and calcium or zinc, this may result in interference with neurotransmission at the synapse.Lead and cadmium has the capacity to induce synergistic toxicity in astrocytes that may compromise the BBB and may cause behavioral dysfunction in developing rats (Rai et al., 2010). Cadmium exposure and accumulation in the body start at young age, one of the heavy metals, is an important environmental pollutant and a potent toxicant to organism. It poses a severe threat to the growth of the organism, and also has been recognized as a human carcinogen (liu, 2014) . Also oxidative stress has been proposed as a possible mechanism involved in Cd toxicity. Thus, prolonged exposure to Cd will cause toxic effect due to its accumulation over time in a variety of tissues, including kidneys, liver, central nervous system (CNS), and peripheral neuronal systems. Cd can be uptaken from the nasal mucosa or olfactory pathways into the peripheral and central neurons; for the latter and can increase the blood brain barrier (BBB) permeability (Moumita et al., 2013) .

Exposure routes in children are mainly via food, environmental tobacco smoke and house dust. Excretion from the body is limited. Cadmium accumulation in the kidney is responsible for effects such as nephrotoxicity and osteoporosis which are observed at adult age (Jerez, 2010). Cadmium exposure through inhalation is also associated with lung cancer in adulthood. Although transfer to the neonate through the placenta and through breast milk is limited, teratogenic and developmental effects were observed in experimental animals (Schaeters et al., 2006)

In spite of several publications addressing the action of lead and cadmium on neurotransmitter systems and the associated enzymes, such as monoamine oxidase (MAO), which is a mitochondrial enzyme responsible for the oxidative deamination of a variety of biogenic amines (Abdel-Rahman et al.,1992) many of these studies are highly variable and lack confirmation (Antonio et al., 2008). The effect of lead and cadmium depends on various aspects, such as exposure level, duration of exposure, the species of used animals and the developmental stage at which the animals were exposed. (Devi et al., 2005). The alterations in functioning of catecholamine could lead to brain dysfunction involving over activation of biogenic amine system (Szczerbak et al., 2007) .This is in agreement with(Baksi and Hughes,1982) who studied the effect of high lead level doses on the activity of catecholamine, nor epinephrine or dopamine and amino acid precursor, tyrosine.Cadmium may also induced cellular damage and lipid peroxidation in rat brain which lead to destruction in most brain enzymes including MAO (Sheril et al., 2003) . The present work aimed to study the alteration in MAO activity in different CNS regions of adult male guinea pigs injected with lead acetate and cadmium chloride with the study of barley plant as diet benefactor against the influence of heavy metals.

MATERIALS AND METHODS

Experimental animals

Adult males guinea pigs weighing (600-800g) were purchased from King Abd El Aziz University Research Institute were used in this study. They were housed under normal environmental conditions of temperature ($22\pm2^{\circ}\text{C}$), relative humidity (50 ± 10), and 12 photoperiod (12hr. light and dark cycle) . Animals were allowed free excess of food and water. Guinea pigs were divided randomly into five groups with five animals in each group. The first group served as control i.e was injected with saline solution .Animals in the second group were subcutaneous injected with lead acetate (30mg/kg body weight) parallel with the

third group which injected with cadmium chloride (1mg/kg body weight.) , fourth group was injected with mixed dose of lead and cadmium. And the last group was feed with aquas solution of barley (5 ml/kg body weight) by gastric tube for two weeks ,followed by injection with mixed dose of lead acetate (30mg/kg body weight) and cadmium chloride (1mg/kg body weight) . The injection with hevy metals was dialy and the decapitation was after 2,4,6 and 8 weeks of injection (lead acetate and cadmium chloride was purchased from Aldrich Chemical co). Approved by the Committee of King Abd El Aziz University for care and use of laboratory animals.

Handling of tissue samples:

Animals were decapitated, the brain and spinal cord of each animal were immediately removed and the brain was dissected in dry ice into the following regions: cerebral cortex, caudate nucleus - putamen, Thalamus, hypothalamus, superior colliculus, inferior colliculus, cerebellum, pons and medulla.

Extraction and tissue samples:

Tissue samples were weighed and homogenized in 0.2M.phosphate buffer pH 7.4 and monoamine oxidase was assayed by the method of (Krajl , 1965)as described by (OLcese - Devlaming 1979) Aliquots of 0.1ml of tissue homogenate were added to 0.25 ml of substrate (1Mm kynuramine)and 1.4 ml of the phosphate buffer 0.2M(Ph 7.4). The samples were incubated for 1 hr at 37°C in a shaker bath .Tissue blank for each samples were subjected to the same treatment , but the substrate was added after the incubation period .The reaction was stopped by the addition of 0.25ml of a 10%ZnSo₄ solution followed by 0.25ml of 1N NaOH. After boiling for 2min. the assay tubes were centrifuged for 10 min.at 3000 r.p.m.The supernatant was then decanted into another tube to which 2ml of 1N.

NaOH were added. The fluorescent products, 4-hydroxyquinoline (4-HQ) was measured in a spectrophotofluorometer, at 315nm (excitation) and 380(emission).Standard curve of 4-HQ were run in parallel with each assay. Monoamine oxidase activity is expressed as $\mu\text{M}4\text{-HQ}/\text{hr/g}$ fresh tissue.

Statistical analysis has been carried out by student t-test to compare between paired groups, results are expressed as mean $\pm\text{SD}$ and the statistical test was considered significant at $p<0.05$ and $p<0.01$ levels (Armitage, 1974).

RESULTS

The changes in MAO activity in various CNS regions of guinea pigs following subcutaneous injection with lead acetate (30 mg./ kg) ,cadmium chloride(1mg/kg),mixed dose of lead and cadmium ,barley plant and mixed dose are presented in (figs 1-10)

As shown in (Fig. 1) the subcutaneous injection with lead and mixed dose provoked an outstanding increase in the MAO activity in cerebral cortex while cadmium induced general decrease. However, the enzyme activity of the caudate putamen (fig. 2) exhibited a marked increase after all times of injection, in the thalamus and hypothalamus (fig.3 and 4) the decrease was also noticed. In (Fig. 5) MAO was affected by lead and cadmium administration, activity of this enzyme was decreased when compared to the controls. However a very highly decline occurred in the most time interval. In (fig6) the activity of MAO of guinea pig inferior colliculus was significantly increased in all times of experimental groups when compared with the control. A significant decrease in MAO activity occurred also in cerebellum in all times of experiments (fig. 7).Variable changes was occurred in pons, medulla and spinal cord(fig.8,9 and 10), improvement was noticed in all groups which received barley plant.

DISCUSSION

Neurotoxicity is a term used to describe neurophysiological changes caused by exposure to toxic agents. Such exposure can result in neurocognitive symptoms and/or psychiatric disturbances. Common toxic agents include heavy metals, drugs, organophosphates, bacterial, and animal neurotoxins. Among heavy metal exposures, lead and cadmium exposure is one of the most common exposures that can lead to significant neuropsychological and functional decline in humans. MAO have the ability to limit the action of many important neurotransmitters, oxidatively deaminate neurotransmitter and xenobiotic amines thereby setting up the basis of rapid repetitive response (Nehru and Dua , 1998). MAO plays an important role in catabolizing the neuroactive amines (Shaban etal., 2014). The general decrease of the activity of these enzyme observed after the treatment with lead and cadmium suggested that lead and cadmium is able to reach the CNS and impair its function .Lead and cadmium has been found to be a causative

agents of various sorts of disorders, including neurological, nephrological, immunological, cardiac, motor, reproductive and even genetic (Zahir et al., 2005). The transport of essential metals and other nutrients across tight membrane barriers such as the gastrointestinal tract and blood-brain barrier is mediated by specific transport mechanisms. Specific transporters take up metals at the apical surface and export them at the basolateral surface, and are involved in their intracellular distribution (Bressler et al., 2007).

(Teresa et al., 2003) determined the level of acid phosphatase, alkaline phosphatase, catalase, acetylcholinesterase and ATPase after lead and cadmium administration and suggested that both metals exert an additive effect, either competing for the same inhibitory binding sites or increasing oxidative damage in the CNS.

The data of the present work showed that, lead and cadmium administration provoked variable changes in MAO in the CNS regions of guinea pigs, which may be due to that lead resulted alterations in membrane bound enzymes, alteration in the level of neurotransmitters receptors which in turn affects on membrane bound enzymes and these data agreement with (Reddy et al., 2003). The results of the present study showed that lead-injection altered the aminergic system with a decrease in mitochondrial MAO activity and this converging agreement with studies of (Zhu et al., 2005). Another evidence suggested that lead can also enter the neuronal intracellular environment, and may activates the calcium mediated synaptic vesicle release mechanisms (Bressler and Goldstein 1991).

(Karadas et al., 2014) were found the ability of lead and or cadmium to penetrate and damage the blood-brain barrier very rapidly, leading to a dysfunction of the blood-brain barrier system .

(Gneurre et al., 2001) and (Swaran et al., 2007) were studied numbers of chemical compounds which considered as a potent inhibitor of MAO activity in vertebrate tissues and they were found that all experimental animals were depressed under lead injection which may be due decreased of the neurotransmitters and this agreement with Ieret et al., (2003) which examined the effects of early simultaneous exposure to low level of lead and cadmium on anxiety - like behavior in the rat and on monoamine levels in different CNS regions and showed an increased in indices of anxiety and change in behavior which related to alteration in serotonergic and dopaminergic receptors.

In the present study, a significant decrease in MAO activity was noticed in most CNS regions. The decrease in both MAO in CNS regions may be due to cellular damage induced by cadmium and this assumed by studies of Alfano et al., (1983). The high affinity of lead for sulphhydryl groups in enzymes may led to decrease in MAO and catecholamines. This may not be a direct inhibitory action of lead but a consequence of the inhibitory effects on cholinergic system (Reddy et al., 2003). Lead also crosses blood brain barrier in developing animals, disrupts the main structural components of blood-brain barrier by injuring the brain glial cells. Also it competes with calcium for common binding sites and is incorporated into calcium transport systems in nervous system, where it is important for neurotransmitter release and regulation (kambiz et al., 2003) .A significant depletion of MAO observed in the present study also reported by Saxena and Flora (2006). According to Shaffi (1995) the alterations in the activity of MAO may produce marked changes in the CNS function as responses to the contamination of the brain with the metal in different manner. The general decrease of the activity of enzymes observed in the experimented animals after treatment with lead and cadmium suggest that these metals able to reach the CNS and impair its function by neurochemical changes such as increasing oxidative damage in the CNS (Antonio et al ., 2003 and Elsayed., et al., 2006) .Experiments carried out by Sabbioni and Marafante (1976) with fractional liver mitochondria from rats injected with a tracer dose of lead revealed that most of the radioactivity was associated with inner membrane mitochondria and its fractions . MAO, which is also considered to be interneuronal enzymes, located largely in the outer membrane of mitochondria or in the exoplasm.

The specific localization of MAO and the accumulation of lead in the mitochondria could act as an inhibitor of the enzyme (Concepción et al., 2012) . Lead may led to alterations in membrane fluidity, which in turn result inactivation of membrane bound enzymes, changed ion permeability and alteration in the levels of neurotransmitter receptors, which in turn may result into alteration of membrane function (Kumar et al., 1996) and (Govinder and Prahald 2000). The available finding indicates the neurotoxic effects of Cd that was associated with both biochemical changes of the cell and functional changes of central nervous system, suggesting that neurotoxic effects may play a role in the systemic toxic effects of the exposure to Cd, particularly the long-term exposure..The brain is also considered as a sensitive organ prone to oxidative damage because of its low levels of protective enzymes to eliminate free radicals resulted from lead intoxication ,may led to brain damage and change in the activity of the related enzymes(Chalter ,2008). An

impairment of the cholinergic function was described in lead and cadmium treated animals, including alterations in neuorotransmitters, Suszkiw (2004).

Figure 1: Effect of lead ,cadmium and mixed dose of lead and cadmium on the activity of MAO in cerebral cortex of male guinea pig and treatment of the mixed dose with Hardium valgara




Figure 2: Effect of lead ,cadmium and mixed dose of lead and cadmium on the activity of MAO in caudate-putamen of male guinea pig and treatment of the mixed dose with Hardium valgara




Figure 3: Effect of lead ,cadmium and mixed dose of lead and cadmium on the activity of MAO in thalamus of male guinea pig and treatment of the mixed dose with Hardium valgara




Figure 4: Effect of lead ,cadmium and mixed dose of lead and cadmium on the activity of MAO in hypothalamus of male guinea pig and treatment of the mixed dose with Hardium valgara




Figure 5: Effect of lead ,cadmium and mixed dose of lead and cadmium on the activity of MAO in superior colliculus of male guinea pig and treatment of the mixed dose with Hardium valgara




Figure 6: Effect of lead ,cadmium and mixed dose of lead and cadmium on the activity of MAO in inferior colliculus of male guinea pig and treatment of the mixed dose with Hardium valgara




Figure 7: Effect of lead ,cadmium and mixed dose of lead and cadmium on the activity of MAO in cerebellum of male guinea pig and treatment of the mixed dose with Hardium valgara




Figure 8: Effect of lead ,cadmium and mixed dose of lead and cadmium on the activity of MAO in pones of male guinea pig and treatment of the mixed dose with Hardium valgara





Figure 9: Effect of lead ,cadmium and mixed dose of lead and cadmium on the activity of MAO in medulla of male guinea pig and treatment of the mixed dose with Hardium valgara



The results obtained in the present work suggest that the decrease of degrading enzyme activities as in the case of MAO following lead and cadmium administration in thalamus and hypothalamus area may be led to intoxication and neurobehavioral changes such as cognitive and attention deficit as well as hyperactivity, which is commonly observed both in lead and cadmium intoxication and perturbed nonaminergic neurotransmission. Sanah et al., 2012 explain the decrease of MAO in most CNS regions under the effect of Cadmium administration as, cadmium have neurotoxic effects, via damages cells of the cerebellar cortices of young rats as well as rabbits. Simultaneous injection of lead and cadmium in a mixed dose increased monoamine oxidase activity all ten parts of rat CNS when compared to treated group. Several reviews have been published recently on the toxicity of heavy metals, but few of these have made any reference to the influence of the nutritional state of animals upon their tolerance of heavy metals. In overall, it has been found that barley may have a protective-like ability to hinder lead and cadmium toxicity in the central nervous system. Barley grass is rich in vitamins and minerals, has antioxidant properties, and has been shown to reduce neurotoxicity of the heavy metals and also used as a cancer preventive, detoxification of pollutants, protection against solar and other forms of radiation, and boosting energy and immunity (Yu.etal,2002). Treatment with barley increases monoamine oxidase activity and contents of all central nervous system regions .Our results agree with a study by (Karabulut-Bulan et al., 2008) which determines whether vitamin C, vitamin E and selenium have the ability and preventive effects of the toxicity of the cadmium in mice. When male guinea pig were feed aqueous barley solution dose (5 mg) for two weeks and then treated with a mixture of lead and cadmium occurred a significant improvement of the enzyme monoamine oxidase in each of the cerebral cortex, caudate nucleus and putamen, thalamus, hypothalamus, superior colliculus, inferior, cerebellum cerebellum, pones and spinal cord when compared to the group exposed to a mixture of lead and cadmium. This results was agree with the study by (Bawazir, 2010) which found that the use of barley (oral) in albino mice has led to an improvement in the level of neurotransmitters (DA, GABA and 5-HT) in different.

REFERENCES

1. Abdel-Rahman,T.; Aly, M.S.; Mohamed, m.I. and Ismail,S.M.,1992:combined effect of ethanol administration and magnetic field exposure on the activities of MAO and ATPase in different CNS regions of two mammals. Proc.Zool.Soc.23(1):35-42 .
2. Alfano, D.P.; Petil, T.L. and Le Boutillier, 1983: Development and plasticity of the hippocampalcholinergic system in normal and early lead-exposed rats. Brain Res. 312:117-124.
3. Antonio, M.T.; Corridor,L. and Ieret, M.L., 2003:Study of the activity of several brain enzymes like markers of the neurotoxicity induced by perinatal exposure to lead and or cadmium. Toxicol. Lett. 143(3):331-340.
4. Antonio, M.T.and Gonzalez, E.L., 2008: Toxic effects of prenatal lead exposure on the brain of rats involvement of oxidative stress and the beneficial role Of antioxidants. Food. Chem. Toxicol 46(6):2089 – 2095.

5. Armitage, P.(1974). Statical Methods in medical research. Paried Student "t" test. 3rd ed. Blackwell Scientific publ., London, pp: 116-120.
6. Baksi, S.N. Hughes, M.J.,1982: Regional alterations of brain catecholamines by injection in adult Lead rat. *J. Arch. Toxicol.* 50(1) : 11 – 18.
7. Bawazir, A.E.,2010 :Investigations on the Chronic Effect of Talbina (Barly Water) on Hormone(Cortisol and Testosterone), Reproductive System and Some Neurotransmitter Contents in Different Brain Areas of Male Albino Rats
8. American-Eurasian Journal of Scientific Research 5 (2): 134-142. Bressler JP1, Olivi L, Cheong JH, Kim Y, Maerten A, Bannon D:2007
9. Metal transporters in intestine and brain: their involvement in metal-associated neurotoxicities. *Hum Exp Toxicol.* 26(3):221-9.
10. Bressler, J.P.and Goldstein,G.W.,1991: Mechanism of lead neurotoxicity *Biochem. Pharmacol.* 41 :479 – 484.
11. Chatler, J.R. ,2008 : lead. exposure, brain damage and adult behavior *Environ Sci. Technol.* . 42 (14) :5036.
12. Concepción ,N., Marisela, M. ,Armenta, C.2012: Lead neurotoxicity: effects on brain nitric oxide synthase. *J. Molecular Histology* : 43(5): 553-563
13. Devi, C. B. ; Reddy, G. H.; Prasanthi, R.P.; Chetty, C.S. and Reddy, G, R., 2005: Developmental lead exposure alters mitochondrial monoamine oxidase and synaptosomal catecholamine levels in rat brain. *Int. J. Dev. Neuro Sci.* 23 (1):375 – 81 .
14. Elsayed, I. H.; Lotfy, O. A.; El Kawaga, Y. A.; Uasif , W. A. and Elshahat. M., 2006 : Prominent free radicals scavenging activity of tannic acid in lead induced oxidative stress in experimental mice. *Toxicology and industrial health* 22(4):157-163.
15. Gnerre., C.; Kosel, M.; Baumann, P.; Carrupt, P . A. and Testa, B., 2001 Interaction with psychiatric drugs with monoamine oxidase in rat Brain. *J. Pharmacol.* 53(8):1125-1130.
16. Govinder, J. S. and Prahalad, K. S.,2000 :Alterations in some membrane Properties in rat brain following exposure to lead. *Cytobios.* 103:103-109.
17. Gui H., Qing-L, Ming J.and Mei S. ,2013: Protective effect of puerarin on lead-induced mouse cognitive impairment via altering activities of acetyl cholinesterase, monoamine oxidase and nitric oxide synthase 35(3) : 502-510
18. Gupta N., Goswami B., Singh N', Koner B.and Garg R .,2011:
19. Lead poisoning associated with Ayurvedic drug presenting as intestinal obstruction: a case report. *Clin. Chim. Acta.* 14;412(1-2):213-4
20. Lead poisoning associated with Ayurvedic drug presenting as intestinal obstruction: a case report. HYPERLINK "javascript:AL_get(this,%20'jour',%20'Clin%20Chim%20Acta.');" \o "Clinica chimica acta; international journal of clinical chemistry." *Clin. Chim. Acta.* 14;412(1-2):213-4
21. Jerez S1, Motas M, Cánovas RA, Talavera J, Almela RM, Del Río AB.2010: Accumulation and tissue distribution of heavy metals and essential elements in loggerhead turtles (*Caretta caretta*) from Spanish Mediterranean coastline of Murcia. *Chemosphere.* 78(3):256-64.
22. Kambiz, S.; Kebrree, A.; Minaee, Bstand,N; Azizi, E. and Abdalla, M., 2003: Biochemical and ultrestructural evidences for toxicity of lead through free radicals in rat brain. *Human & Experimental toxicology* 22(6):417-423.
23. Karadas,S. ; Refah S. M.; Aslan H. G.; Recep D.N. ; Latif D.P.and Recep H. D.,2014 : Serum Levels of Trace Elements and Heavy Metals in Patients with Acute Hemorrhagic Stroke . *The Journal of Membrane Biology* 247(2): 175-180
24. Karadas,S. ; Refah S. M.; Aslan H. G.; Recep D.N. Latif D.P.and Recep H. D.,2014 : Serum Levels of Trace Elements and Heavy Metals in Patients with Acute Hemorrhagic Stroke . *The Journal of Membrane Biology* 247(2): 175-180
25. Kim Y1, Ha EH, Park H, Ha M, Kim Y, Hong YC, Kim EJ, Kim BN.,2013:
26. Prenatal lead and cadmium co-exposure and infant neurodevelopment at 6 months of age: the Mothers and Children's Environmental Health (MOCEH) study. *Neurotoxicology.* 35:15-22.
27. Krajl, N. ,1965 : A rapid microflurometric determination of monoamine Oxidase. *Biochem Pharmacol.* 14 : 1684-1685.
28. Kumar, R.; Srivastava, S.; Agrawal, A. K. and seth, P. K., 1996 Alteration in some membrane properties in rat brain following exposure to manganese. *Pharmacol.Toxicol.* 79:47-8.
29. Leret. M. L.; Millan, J. A. and Antonio, M. T.,(2003): Perinatal exposure to lead and cadmium effects anxiety – Like behaviors. *Toxicology* 186(1-2):125-130.
30. Liu Y1, Li Y2, Liu K1, Shen J1.,2014: Exposing to cadmium stress cause profound toxic effect on microbiota of the mice intestinal tract): *PLoS One* 9(2): 85
31. Modak, A. T.; Weintraub, S. T. and Stavinoba, W. B., 1975: Effect of chronic ingestion of lead on the central cholinergic system in rat brain regions. *Toxicol. App. Pharm.* 34 : 340 -347.

32. Moumita, G., Sarmila,k. and Ranabir,S.,2013: Cadmium induced pathophysiology: Prophylactic role of edible jute (*Corchorus olitorius*) leaves with special emphasis on oxidative stress and mitochondrial involvement. *Food and Chemical Toxicology* 0:188–198
33. Nehru, B. and Dua, R, 1998: The effect of dietary selenium on lead neurotoxicity. *J. Environ. Path, Toxicol. and Oncology*.16(1) :47-50.
34. Olcese, J. M. and Vlaming, V. L., 1979 Characteristic of monoamine oxidase activity in The hypothalamus of *Carassius auratus* as assayed by a Fluorometric technique. *Comp. Biochem. Physiol.* 62C :213 – 215.
35. Rai A. , Maurya SK., Khare P., Srivastava A.and Bandyopadhyay S., 2010 :Characterization of developmental neurotoxicity of As, Cd, and Pb mixture: synergistic action of metal mixture in glial and neuronal functions .*Toxicol .Sci.* 2010 Dec;118(2):586-601
36. Ranabir, S., Moumita,G.and Sarmila,k:2013: Toxic effects of lead exposure in Wistar rats: Involvement of oxidative stress and the beneficial role of edible jute (*Corchorus olitorius*) leaves. *Food and Chemical Toxicology*. 55: 78–91
37. Reddy, G. R.; Basha ,M. R.; Devi, C. B.; Surash, A.; Baker, J. L.; Shafeek, A. and Cheetty, C.S. 2003 : lead induced effects on acetylcholinesterase activity in cerebellum and hippocampus of developing rat. *Int. J. Dev. Neurosci.* 21 (6) : 347- 352.
38. Sabbioni, E. and Marafanta , E., 1976 : Identification of lead binding components in rat liver. In vivo study. *Chem. Biol. Interact.* 15: 271 – 276.
39. Sandra V., Lucila A and Patricia I.,2008: Aluminium and lead: molecular mechanisms of brain toxicity. *Archives of Toxicology* 82(11): 789-802
40. Sanah ,S., Zena, G., Arnab, C., and Dietrich, B.,2012: Metal Toxicity at the Synapse: Presynaptic, Postsynaptic, and Long-Term Effects.
41. Journal of Toxicology 13:42
42. Saxena, S. and Flora, S.J., 2006 Changes in brain biogenic amines and haembiosyntheis and their response to combined administration of succimers and centella asiatica in lead poisoned rats. *J, Pharm. Pharmacol.* 58(4): 547-55.
43. Shaban ,N.Z. ;Masoud,M.S. ;Mawafi,M.A.and Sadek.O.M.,2014: Effect of Cd, Zn and Hg complexes of barbituric acid and thiouracil on rat brain monoamine oxidase-B (in vitro).*J. Chemico-Biological Interactions.* 208 :37-46.
44. Schoeters G., Den Hond E., Zuurbier M., Naginiene R., van den Hazel P., Stilianakis N., Ronchetti R. and Koppe JG.,2006: Cadmium and children: exposure and health effects. *Acta Paediatr Suppl.* 95(453):50-4.
45. Shaffi, S. A., 1995 : Sub lethal effect on mercury and lead on monoamine oxidase in different regions of the brain in three fresh water telosts. *Revista Espanola De Fisiologica*, 51 (3) 125 – 128.
46. Sheril,D. ;Janice,L. ;Amichand ,D. and Watkins,S.(2003):Through metal biding,
47. Curcumin protects against lead-cadmium-induced lipid per oxidation in rat brain homogenates and against lead-induced tissue damage in rat brain .*j of Inorganic Biochem* .98:266-575.
48. Silbert5hynjm,rgeld, E. K., 1992 : Management of lead neurotoxicity. *FASEB*, 6 : 3201 -3206.
49. Suszkiw .J.B. , 2004:Presynaptic disruption of transmitters release by lead .*Neurotoxicology* 25:599-604.
50. Swaran, J. S.; Flora, S. G. and Mehta, A, (2007): Reversal of lead – induced neuronal apoptosis by chelation treatment in rats: Role of reactive O₂ species and intracellular Ca⁺². *J. Pharm. Exp. Therap.* 322 : 108 – 116.
51. Szcerbak, G.; Nowak, P.; Kostrzewa, R. M. and Brus. R., 2007: Maternal lead exposure produces long – term enhancement of dopaminergic reactivity in rat offspring. *Neurochem. Res.* 32 (10) : 1791-8.
52. Teresa, M. A.; Laura, L. and Luisa, L.2003:Study of the activity of several brain enzymes like markers of the neurotoxicity induced by Perinatal exposure to lead and / or cadmium. *Toxicology* 143(3):331 – 340.
53. Wright, R. D. and Baccavell, A., 2007 : Metals and neurotoxicology. *J. Nutr.* 173 (12):2809 -2809 – 13.
54. Yu YM.; Wu CH. ; Tseng CE.; Tsai CE. and Chang WC. 2002:Antioxidative and hypolipidemic effects of barley leaf essence in a rabbit model of atherosclerosis. *Jpn J Pharmacol* .89:142-148.
55. Zahir,F. ; Rizwi,J.S. ; Haq,S.A. and khan,S.R.2005: Low dose mercury toxicity and human health. *Environmental Toxicol. and Pharmacol.* 20(2)
56. Zheng W1, Aschner M, Ghersi-Egea JF.2003: Brain barrier systems: a new frontier in metal neurotoxicological research. *oxicol Appl Pharmacol.* (1):1-11.
57. Zhu, Z. W.; Yang, L. R.; Dong, J. G. and Zheo,Y. Z., 2005 : Study on the neurotoxic effects of low – level lead exposure. *J. Zhejiang. Univ. Sci.* 6(7): 686 - 692.