

New Phthalic Acid Diamide Flubendiamide-Its Effects on Non-Target Species-A Review

C P Anitha Devi*, Bismi Antony

Department of Zoology, N.S.S. Hindu College, Changanacherry, Kottayam, Kerala,
India

Review Article

Received: 02-Feb-2022,
Manuscript No. JAAS-22-
57330; **Editor assigned:** 07-
Feb-2022, Pre QC No. JAAS-22-
57330 (PQ); **Reviewed:** 22-Feb-
2022, QC No. JAAS-22-57330;
Revised: 04-Apr-2022,
Manuscript No. JAAS-22-57330
(R); **Published:** 12-Apr-2022,
DOI: 10.4172/E2347-
226X.11.04.005

***For Correspondence:** C P Anitha
Devi, Department of Zoology,
Research and Post Graduate,
University of N.S.S. Hindu College,
Changanacherry, Kottayam, Kerala,
India, **Tel:** 9400725260;
Email: cpanitha@gmail.com

Keywords: Biochemical changes;
Enzyme profile; Hematological
changes; Histopathological
alterations; Lepidopteran
insecticide; Morphological changes

ABSTRACT

Pesticides are used to protect food crops, but can also threaten organisms unrelated to their intended targets. With population growth and pesticide use increasing, food is becoming scarcer and pesticides are used to combat them. Flubendiamide is an insecticide that is toxic to lepidopterans and other non-target organisms. The objective of our review is to identify how the insecticide affects non-target organisms. Several recent studies are systematically reviewed and analyzed. A review of a number of different studies undertaken on different organisms exposed to the chemical showed that the pesticide is capable of causing morphological, biochemical, hematological, enzymatic, histopathological, and behavioral changes.

INTRODUCTION

Crop protection is very important in sustainable agriculture. In order to meet the increasing need for food, growing as well as stored grains should be protected. For this purpose, we use pesticides [1]. Pesticides are a group of chemicals which are used to kill insects, weeds, fungi, bacteria. They are generally called insecticides, herbicides, fungicides, bactericides and rodenticides. Most of these chemicals are designed in such a way that they can act as broad spectrum. But some of them have a specific action [2]. Besides target, pesticides have a serious impact on non-target species also [3]. Most of the pesticides used in the agricultural fields reach the soil. From there they reach the water bodies by surface runoff and leaching. The persistence of these pesticide residues in soil and water affects the soil as well as aquatic life [4]. The present review focuses on the effect of Flubendiamide, a new phthalic acid diamide, a

new generation insecticide on various non-target organisms. It registered in India on 2007 under the trademark Fame [4]. Phthalic acid diamides used as a substitute for organochlorine and organophosphate pesticides due to their high specificity towards lepidoptera and low persistence in the soil.

Flubendiamide (N_2 - [1,1- dimethyl-2-(methyl sulphonyl ethyl)-3-iodo- N_1 -[2-methyl-4-[1,2,2,2-tetrafluoro-1-(trifluoromethyl) ethyl] phenyl]-1,2-benzene dicarboxamide) is registered in 2008 for sale distribution and use under the Federal Insecticide, Fungicide, Rodenticide Act (FIFRA). It is a new stomach poison insecticide which is used to control important lepidopteran pests. Unlike all the other insecticides which act on the nervous system of the target pest, Flubendiamide act on the ryanodine sensitive receptors in the muscles of insects and causes the cessation of feeding thereby killing them. The insecticide resistance action committee has classified the diamides into a new group (Ryanodine receptor modulators) [4]. Its insecticidal activities are very relevant. It is fast acting and long lasting. Even though no mammalian toxicity of Flubendiamide is reported not so far, most of other non-target organisms are affected by the same. Pesticide usage which is desirable for pest control ultimately results in pollution and will affect non-target organisms also [5,6]. The present review focuses on the effect of Flubendiamide on its various non-target organisms.

Overview about Flubendiamide

Flubendiamide is a novel class of insecticide, the phthalic acid diamide. It has a unique structure. The uniqueness is due to the presence of three structural components (Figure 1 and Table 1).

- A heptafluoroisopropyl unit in the anilide moiety
- A sulphonyl alkyl group in the aliphatic amide moiety
- An iodine atom in the 3rd position in the phthalic acid moiety [7].

The chemical structure of Flubendiamide is given in Figure 1.

Figure 1. Chemical structure of Flubendiamide.




Table 1. Physico-chemical properties of Flubendiamide.

1. Common name: Flubendiamide
2. Chemical name: (N_2 - [1,1- dimethyl-2-(methyl sulphonyl ethyl)-3-iodo- N_1 -[2-methyl-4-[1,2,2,2-tetrafluoro-1-(trifluoromethyl) ethyl] phenyl]-1,2-benzene dicarboxamide
3. Molecular formula: $C_{23} H_{22} F_7 I N_2 O_4 S$
4. Appearance: Colorless crystal
5. Molecular weight: 682.39
6. Odour: No characteristic odour

7. Melting point: 217°C
8. Density (at 20.8°C): 1.659 g/cm ³

MODE OF ACTION

Flubendiamide activates ryanodine sensitive intracellular calcium release channels (RyR receptors) in insects. Rather than affecting the sodium-potassium ion balance in the cell, it affects the calcium ion balance which causes the contraction of insect muscles. Flubendiamide keeps the RyR Receptors in an open state causing massive calcium release leading to the rapid contraction of muscles followed by death. It will induce cessation of feeding by the insects. It is effective against a broad range of pests belonging to the order lepidoptera. It is considered to be a new stomach poison, Oral intoxicant, long lasting and fast acting insecticide. It is marketed with different trade names including Belt, Fame™, Tenos™ and Synapse.

LITERATURE REVIEW

The concentration of toxicant when less than the lethal dose, the exposed organisms might show change in morphological, behavioral, physiological and biochemical changes. Few of such studies are listed below.

Earthworm *Eudrilus eugeniae*

The avoidance behavior and neurotoxicity of earthworm *Eudrilus eugeniae* studied by [8]. The neurotoxicity was studied based on the production of marker enzyme acetyl choline esterase. Toxicity signs such as increased mucus secretion and bloody lesions in the body were observed. The earthworms tend to avoid the Flubendiamide contaminated soil. Apparent changes in the body pigmentation were also there in exposed group compared to the control as noted by [8]. Morphological and physiological alteration in earthworms was observed in response to decreased acetyl choline esterase levels. Kinetic study of the enzyme in the presence and absence of the inhibitor was studied and the results showed that it acted as a competitive inhibitor. The effects of Flubendiamide on *Eudrilus eugeniae* is given in Table 2.

Table 2. The effects of Flubendiamide on *Eudrilus eugeniae*.

Dose of Flubendiamide	Parameters studied	Results
54-94 µg/cm ²	Determination of median lethal concentration	LC ₅₀ - 94.4 µg cm ⁻² (48 hr. paper contact test) 332.21 and 238.31 mg kg ⁻¹ respectively, for 7 and 14 d of artificial soil test
54-94 µg/cm ²	Avoidance test	Animals tend to avoid regions where higher concentrations of Flubendiamide were present.
54-94 µg/cm ²	Morphological changes	Sluggish movement, excessive mucous secretion, coiling and swelling of clitellum, extrusion of coelomic fluid, formation of bloody lesions.
54-94 µg/cm ²	Acetyl choline Esterase activity	As the concentration of Flubendiamide increased, the enzyme inhibition also increased drastically. Concluded that the inhibition is competitive in nature

Daphnia magna

The effect of three diamides (Chlorantraniliprole, Cyantraniliprole and Flubendiamide) on life history, embryonic development and oxidative stress biomarkers of *Daphnia magna* was experimented by [9]. They observed increased ROS, decreased activities of antioxidant enzymes (SOD and Glutathione peroxidase) which was consistent with the downregulated transcription of the corresponding genes. An increased catalase activity was noted but there was no change in the gene expression [9]. Most xenobiotic compounds like pesticides can lead to the production of reactive oxygen species in response to oxidative stress via several mechanisms such as inhibition of electron transport chain in the mitochondrial membrane and subsequent accumulation of its reactive intermediates, inactivation of antioxidant enzymes and decrease in the non-enzymatic antioxidants and membrane lipid peroxidation. The adverse effects of exposure to Flubendiamide in *Daphnia magna* are given on Table 3.

Table 3. Effects of Flubendiamide on *Daphnia magna*.

Dose of Flubendiamide	Parameters studied	Results
-----------------------	--------------------	---------

EC ₅₀ -63.5 µ/L (Acute toxicity study)	ROS	Significant increase
	Antioxidant status.	Significant decrease in the activity of enzymes such as superoxide dismutase and glutathione peroxidase, which is consistent with the downregulated transcription of their genes. Increased catalase activity as a result of increased transcription of cat genes.
EC ₅₀ -36.8 µg/L (chronic toxicity)	Mobility	Decreased
Chronic LOEC19.36 µg/L	Survival growth and reproduction	Affected
48 hr. EC ₅₀ - 30.8 µg/L (acute exposure)	Developmental abnormalities	Embryos had underdeveloped second antennae, curved tail spine and abnormal body region

Drosophila melanogaster

The studies on the effect of Flubendiamide on locomotory activities of non-target species *Drosophila melanogaster* for three generations were reported by [10]. Their studies suggest that chronic Flubendiamide exposure might induce alteration in neurotransmission leading to inconsistent behavioral responses (vision and flight) in other beneficial insects and insect dependent organisms [10]. The results of the study are consolidated in Table 4.

Table 4. Effect of Flubendiamide on locomotory activities of *Drosophila melanogaster*.

Dose of Flubendiamide	Parameters studied	Results
0.25-20 µg/ml	Phototactic behavior	Reduction in light seeking behavior (P, F ₁ and F ₂)
0.25-20 µg/ml	Climbing or negative geotaxis behavior	Decline in climbing behavior. P, F ₁ and F ₂ flies failed to reach the top of the glass tube
0.25-20 µg/ml	Adult acetylcholine esterase activity.	Showed dose dependent decrease in the choline esterase activity.
0.25-20 µg/ml	Adult cytochrome P ₄₅₀ activity	Increased resorufin production indicating the increased enzyme activity.

Toxicological evaluation of Flubendiamide in non-target *Drosophila melanogaster* for different developmental stages showed that the acute LC₅₀ values for both larvae and adult are higher than that of the chronic ones. The study revealed that the acute LC₅₀ value for adult was 28 folds higher than the LC₅₀ for the larvae. Similarly, the chronic LC₅₀ for adults was 32 folds higher than for larvae. This showed that the chemical is more efficient against larvae than adult (larvicidal activity) [11]. The results are given in Table 5.

Table 5. Toxicological evaluation of Flubendiamide for different developmental stages of *Drosophila melanogaster*.

Dose of Flubendiamide	Parameters studied	Result
5000, 7500, 10000 and 15000 µg/ml	LC ₅₀ values for larvae	After 24 hr-12011.64 µg/ml After 48 hr-8317.64 µg/ml After 72 hr-4897.79 µg/ml
100000, 200000, 300000 and 500000 µg/ml	Acute LC ₅₀ values for adult	After 24 hr-338844.26 µg/ml After 48 hr-257038.58 µg/ml After 72 hrs-154881.66 µg/ml
10000, 25000, 50000 and 75000 µg/ml	Chronic LC ₅₀ determination	For larvae-562.34 µg/ml For pupae-676.08 µg/ml For adult flies-17782.79 µg/ml
250, 500, 1000 and 1500 µg/ml	Adult emergence	As the concentration of the Flubendiamide increases, there was a significant reduction in percentage of adult emergence.

Trichogramma chilonis

The effect of Flubendiamide against egg parasitoid *Trichogramma chilonis* (Ishii) (Hymenoptera: Trichogrammatidae)

under laboratory conditions was noted by [12]. The highest percentage parasitisation was noted in Flubendiamide W.G at 50 g.a.i./h. and maximum adult emergence (89.70) was shown at Flubendiamide W.G. at 50 g.a.i./h. (g.a.i/h–grams of active ingredient per hectare)

The effect of Flubendiamide on parasitisation of *Trichogramma chilonis* is given on Table 6.

Table 6. The effect of Flubendiamide on parasitisation of *Trichogramma chilonis*.

Dose of Flubendiamide	Parameters studied	Results
a)50 g.a.i. ha ⁻¹ b)60 g.a.i. ha ⁻¹ c)70 g.a.i. ha ⁻¹	Parasitisation	Little effect on the parasitisation. Untreated group showed greatest parasitisation. Among Flubendiamide treated groups, the highest percentage parasitisation was found to be at 50 g.a.i./ha followed by 60 and 70.
	Adult emergence	Maximum adult emergence was found at the 50 g.a.i./ ha followed by 60 and 70 at 72 hours after treatment.

Fishes

Pesticide pollution in water affects the fish and other aquatic organisms, relatively sensitive to the changes in the surrounding environment [43]. The greatest hazard is to the aquatic organisms, especially fish, which are more susceptible to them as they got concentrated into their organs more readily than terrestrial organisms [14]. The toxicants are accumulated in fish through food chain or by absorption through general body surface which affect severely their life support systems at molecular and biochemical levels [15].

The nutritional value of different species of fish and shellfish depend on their biochemical components such as proteins carbohydrates and lipids. It is important to study the effect of specific concentrations of pesticides on various systems of aquatic organisms. When exposed to certain chemicals, organisms show alterations in their biochemical parameters. An animal uses their muscle protein as an energy source during acute respiratory stress [16]. It is reported that many of the toxicants alter the protein metabolism in fishes such as bleached craft pulp mill effluent, textile mill effluent [17,18], DDT, malathion and mercury [19-21].

Changes in the blood parameters of fish are an indication of its pathophysiological status and their determination is essential for the diagnosis of structural and functional status of fish when exposed to toxicants [22]. Genotoxic compounds will induce alterations of hematological parameters of fish and this is always associated with their physiological state. The biochemical parameters in fish are validated for evaluating their physio pathological conditions and they are sensitive for detecting potential adverse effects and relatively early events of pollutant damage [23]. Alterations in the blood parameters of organisms are due to the damage caused to their hemopoietic organs by the water born pollutants [24-28]. Hematological studies in fishes revealed that the alterations in the blood parameters are indicators of a stressed environment and it gives details about the health conditions of fishes [29]. Studies on the impact of pesticides on fishes revealed that they will interfere with their ability to resist diseases and increases mortality rates [30].

The effect of Flubendiamide on protein metabolism of freshwater fish *Labeo rohita* was studied by [31]. They exposed the fishes with sublethal concentrations of Flubendiamide and found that the exposed ones showed decrease in protein content in various organs. They concluded that the protein depletion may be due to the increased pesticide stress when compared with the control group [31]. The reduction in the protein content is due to the increase use of protein to meet the high energy demand caused by the toxins such as detoxification reactions, tissue repair and homeostasis and stress metabolism [20].

The assessment of the health status of wild fish inhabiting a cotton basin, heavily impacted by pesticides, including Flubendiamide in Benin (West Africa) was done by [32]. Their findings revealed that the pesticides disrupt the endocrine regulation of fish living in the basin and also it causes severe damage to the liver and gonads. Many significant changes were observed in the biochemical profile of *Heteropneustes fossilis* such as decrease in glycogen and total protein content and increased cholesterol content when exposed to Flubendiamide as reported by [33]. Biochemical analysis of Flubendiamide toxicity on gill of *Heteropneustes fossilis* showed that it is not only the gill, but other organs such as liver, kidney, intestine and blood also got affected [34,35]. Determined the consequences of Flubendiamide exposure on hematological parameters of the fish *Oreochromis mossambicus*. The results of the experiment showed that all the blood parameters except WBC count showed reduction. The concentration of the leucocytes was high throughout the experiment. The enzyme profile also showed significant increase. This is an indication of the impact of the toxin to the organism. Table 7 gives a summary of the effect of Flubendiamide on various biological parameters of different fishes.

Table 7. Effects of Flubendiamide on various parameters of different fishes.

Name of fishes	Dose of	Parameters studied	Results
----------------	---------	--------------------	---------

	Flubendiamide		
Labeo rohita	11 µg/L	Total Protein content	Depletion of protein content in all types of tissues was observed when compared to the control.
<i>Heteropneustes fossilis</i>	0.1 ml/L	Biochemical parameters of the intestine:	
		1.Total glucose	Reduces as exposure period increases
		2.Total protein	Lower than the control
<i>Heteropneustes fossilis</i>	0.1 ml/L	3.Total cholesterol	Higher than the control.
		Biochemical analysis of gill:	
		1.Total protein	Decreased as the exposure time increases.
<i>Heteropneustes fossilis</i>	0.1 ml/L	2.Glycogen	Showed decline in the glycogen concentration as exposure time increases.
		3.Cholesterol	Declined initially but after 14th day, there was a sudden increase
		Heamatological parameters.	
<i>Oreochromis mossambicus</i>	10 ppm	1.TEC (Total Erythrocyte Count)	Decreased
		2.TLC (Total Leucocyte Count)	Increased
		3.Hb (Hemoglobin)	Decreased
		4.PCV (Packed cell volume)	Decreased
		5.MCV (Mean Corpuscular Volume)	Increased
		6.MCH (Mean Corpuscular Hemoglobin)	Decreased
		7.MCHC (Mean Corpuscular Hemoglobin Concentration)	Decreased

Wistar rats (*Rattus norvegicus*)

When an organism is in toxic stress, the study of histochemistry and pathology is very important. Studies on the in-vitro and in-vivo effects of Flubendiamide and copper in cytogenotoxicity, oxidative stress and spleen histology of rats showed that they induce oxidative stress, cytogenetic effects along with histoarchitectural changes in spleen [36]. In- vitro studies on Wistar rats and the results are given in Table 8 and In-vivo studies are given in Table 8 and 9.

Table 8. In-vitro studies of Wistar rats when exposed to Flubendiamide and the observed effects.

Dose of Flubendiamide	Parameters studied	Results
1-80 µM	Median lethal concentration	40 µM was selected as the median lethal concentration Flubendiamide treated cells showed 71.88 non-viable cells when compared to control.
40 µM	Viability of splenocytes	The combined treatment of Flubendiamide with antioxidants and xenobiotics decreased the number of non- viable cells.
40 µM	Tunnel assay	Showed higher number of tunnel positive cells when compared to control.
40 µM	Micronuclei formation	7.86% cells show micronuclei formation.
40 µM	DNA fragmentation	Flubendiamide treated splenocytes showed more DNA fragmentation.
40 µM	Comet formation	Flubendiamide alone treatment in splenocytes produced 22.90% comet formation when compared to control.

Table 9. *In-vivo* studies in Wistar rats when exposed to Flubendiamide and the observed effects.

Dose of Flubendiamide	Parameters studied (In-vivo)	Results
40 µM	Oxidative stress biomarkers: 1.Lipid peroxidation levels	Higher when compared to control
	2. Reduced glutathione	Lower when compared to the control.
	3.Glutathione peroxidase	Significant decrease when compared to control.
	4.Glutathione-S-transferase	Significantly lower
	5.Total protein content	No significant decrease.
	6.Catalase activity	No change when comparing to control group.
	Histology	Flubendiamide and copper treated group showed separation of splenocytes and small sized splenic parenchyma. Treatment along with antioxidants showed normal histoarchitecture of spleen.

Sprague Dawley rats (*Rattus norvegicus*)

Studied the hematological alterations on sub-acute exposure to Flubendiamide in Sprague Dawley Rats. They observed significant changes in the female rats than male rats. Thus, concluded that many of those changes were dose independent, but sex specific [4]. The various heamatological parameters studied and the obtained results are consolidated on Table 10.

Table 10. Effects of Flubendiamide on various hematological parameters of Sprague Dawley rats.

Dose of Flubendiamide	Parameters studied	Results
Between 1-2 ml/ 100gm body weight	1.RBC (Red Blood Cells) 2.Hb (Hemoglobin) 3.PCV (Packed Cell Volume) 4.CHCM (Cellular Hemoglobin Concentration Mean) 5.CM (Cellular Hemoglobin Concentration) 6.MCV (Mean Corpuscular Volume) 7.MCH (Mean Corpuscular Hemoglobin) 8.MCHC (Mean Corpuscular Hemoglobin Concentration) 9.RDW (Red Cell Distribution Width) 10.HDW (Hemoglobin Distribution Width) 11.WBC (White Blood Cells) 12.PCT (Procalcitonin) 13.MPC (Mean platelet Component) 14.LPT (Lymphocyte Proliferation test) 15.PLT (Platelet count) 16.MPV (Mean Platelet	<ul style="list-style-type: none"> • In females, Significant decrease in RBC count with respect to dose rate. Rest of the parameters had no significant decrease when compared to control. • In males, Significant difference was exhibited by various parameters. WBC, LPT, PCT, PDW, MPV and PCs showed significant decrease whereas parameters such as RBC, PCV, and HDW showed significant increase. For some parameters there was bidirectional response. • Significant differences in some parameters such as RBC, Hb, PCV, RDW, MCHC CHCM, CM and LPTs were observed in males and female rats of same groups. • Sex dependent differences were consistent in Hb and PCV • In males’ dose dependent increase in RBC, Hb and PCV • In females, a dose dependent decrease of RBC, Hb and PCV was observed.

	Volume) 17.PDW (Platelet Distribution Width)	
--	--	--

CONCLUSION

Flubendiamide is a new generation insecticide belonging to a new chemical class, the phthalic acid diamide. They are used against lepidopteran pests. They have a greater larvicidal activity too. It has a novel action- it affects the muscular system rather than the nervous system. It acts on the ryanodine sensitive calcium channels and causes the contraction of insect muscles. It will result in the rapid cessation of feeding thereby crop damage is prevented.

Although it is designed against lepidopteran pests, they have a greater impact on the non-target species also. They affect different organisms differently. No mammalian toxicity is reported so far, there were reports of Flubendiamide poisoning in buffalos in response to consumption of fodder recently sprayed with pesticide. Morphological, biochemical, histopathological, hematological parameters of various organisms towards sub-lethal concentrations of Flubendiamide was studied. The results of the study revealed that Flubendiamide has a significant impact on all these parameters. The decrease in the antioxidant enzymes and production of ROS reveals the oxidative stress caused by the chemical. Inhibition of acetyl choline esterase activity showed the ability of the chemical for competitive inhibition. The studies on *D. melanogaster* showed that the insecticide have larvicidal activity on non-target insects also. Since water bodies are the ultimate destination of the pesticides from agricultural fields, fishes were studied mostly. The effects of this insecticide should be monitored properly for protecting the non-target organisms, which accidentally comes in contact with the chemical.

REFERENCES

1. Das SK, et al. Flubendiamide as New Generation Insecticide in Plant Toxicology; A Policy Paper. Adv Clin Toxicol. 2017;2
2. Ravindran J, et al. Organochlorine pesticides, their toxic effects on living organisms and their fate in the environment. Interdiscip Toxicol. 2017;9: 90-100.
3. Sharma DR, et al. Use of pesticides in Nepal and impacts on human health and Environment. The J of Agric Environ. 2012;13:67-74.
4. Vemu B, et al. Hematological alterations on sub-acute exposure to Flubendiamide in Sprague Dawley Rats. Toxicol Int. 2014;21:288-293.
5. Pimental D, et al. Pesticides: amount applied and amount reaching pests. Bioscience. 1986;36:86-88.
6. Matsumura F, et al (1975). Toxicology of insecticides. 1st Ed. Plenum press, Springer New York, and London.
7. Tonishi M, et al. Flubendiamide, a Novel Insecticide Highly Active against lepidopterous Insect Pests. Pestic Sci. 2005;30:354-360.
8. Jadhav SS, et al. Effect of Flubendiamide on morphology, avoidance behaviour and acetylcholinesterase activity in earthworm *Eudrilus eugeniae*. Int J Pharm Pharm Sci. 2017;9:233-238.
9. Cui F, et al. Effects of three diamides (Chlorantraniliprole, Cyantraniliprole and Flubendiamide) on life history, embryonic development and oxidative stress biomarkers of *Daphnia magna*. Chemosphere. 2017;169:107-116.

10. Sarkar S, et al. Flubendiamide affects visual and locomotory activities of *Drosophila melanogaster* for three successive generations (P, F1 and F2). *Invert Neurosci*. 2018;18.
11. Sarkar S, et al. Toxicological evaluation of a new lepidopteran insecticide, Flubendiamide, in non-target *Drosophila melanogaster* Meigen (Diptera: *Drosophilidae*). *Iran J Toxicol*. 2018;12:45-50.
12. Madhu Sudhanan E, et al. Toxicity of Flubendiamide 20 WG against egg parasitoid, *Trichogramma chilonis* (Hymenoptera; trichogrammatidae) under laboratory conditions. *J Biol Control*. 2014;28:147-150.
13. Ayas Z, et al. Histopathological changes in the livers and kidneys of fish in Sariyar reservoir, Turkey. *Environ Toxicol Pharmacol*. 2007;23:242-249.
14. Kamble GB, et al. Effect of acute exposure of Endosulfan and Chlorpyrifos on the biochemical composition of the freshwater fish *Sarotherodon mossambicus*. *Indian J Environ Sci*. 2000;4:97-102.
15. Rawat DK, Bais VS, Agarwal NC. The correlative study on liver glycogen and Endosulfan toxicity in *Heteropneustes fossilis*. *J Environ Biol*. 2000;23:205-207.
16. Creach Y, et al. Le Jeune et la réalimentation chez la carpe (*Cyprinus Carpio L.*). *J Physiol Paris*. 1974;68:245-260.
17. McLeay DJ, et al. Growth stimulation and biochemical changes in juvenile Coho salmon (*Oncorhynchus kisutch*) exposed to bleached craft pulp mill effluent to 200 days. *J Fish Res board Can*. 1974;31:1043-1049.
18. MurugesanAG, et al. Histopathological and histochemical changes in the oocytes of the air breathing fish *Heteropneustes fossilis* (Bloch) exposed to textile mill effluent. *Bull Environ Contam Toxicol*. 1992;48:929-936.
19. Verma SR, et al. Toxicity of Swascol IP (SLS) to *Channa punctatus* and *Cirrhina mrigala*: Biochemical alterations. *Bull Environ Contam Toxicol*. 1979;21:711-718.
20. Ramalingam K, et al. Effects of sublethal levels of DDT, malathion and mercury on tissue proteins of *Sarotherodon mossambicus* (Peters). *Indian Acad Sci (anim sci)* 1982;91:501-505.
21. Sharma PD, et al. (1989). *Toxicology*, Rastogi publications, Meerut.
22. SampathK, et al. Effects of copper and zinc on blood parameters and prediction of their recovery on *Oreochromis mossambicus*. *Indian J Fish*. 1998;45:129-139.
23. Osman KA, et al. Monitoring of pesticide residues in vegetables marketed in Al-Qassim region, Saudi Arabia. *Ecotoxicol Environ Saf*. 2010 ;73:1433-1439.
24. Reichenbach, et al. The blood components of fish with relation to parasites, infections and water pollution. *Bull Epizoot Dis*. 1966;65:1039-54.

25. Gardner GR, et al. Histological and haematological responses of an estuarine teleost to cadmium. J Fish Res Board Can. 1970;27:2185-2196.
26. Saad MAH, et al. The effect of pollution on the blood characteristics of *Tilapia zillii* G. Wat Air and Soil Poll. 1973;2:171-179.
27. Dawson MA, et al. Effects of long-term mercury exposure on haematology of striped bass *Morone saxatilis*. Fish Bull. 1975;80:389-392.
28. Bhatkar NV, et al. Chromium (III) induced haematological alterations in Indian common carp, *Labeo rohita* (Ham). J Appl Nat Sci. 2011;3:258 -263.
29. Meraj M, et al. Alteration in haematology of *Cyprinus carpio* under the stress of pollution of water bodies of Kashmir valley. Int J Fish Aquat. 2017;5:176-179.
30. Roy B, et al. Some haematological investigations on *Oreochromis niloticus* (Trewavas) following exposure to Thiamethoxam. Acta Zool Litu. 2011;21:301-305.
31. Nirmalakallagda, et al. Flubendiamide a phthalic acid diamide effect on protein metabolism of freshwater fish *Labeo rohita*. Int J Recent Sci Res. 2014;5:1554-1557.
32. Agbohessi PT, et al. Assessment of the health status of wild fish inhabiting a cotton basin heavily impacted by pesticides in Benin (West Africa). Sci Total Environ. 2015;15: 506-507,567-84.
33. Avinashe AM, et al. Effect of biochemical parameters of intestine of freshwater fish, *Heteropneustes fossilis* (Bloch), when exposed to Flubendiamide. Vidyabharati Int interdiscipl Res J. 2015;4:89-94.
34. AvinasheAM, et al. Biochemical analysis of Flubendiamide toxicity of gill of *Heteropneustesfossilis* (Bloch). Int J appl Univ Res. 2017;4:27-30.
35. AmpiliM, et al. Consequences of FAME (Flubendiamide 480cc) Exposure on hematological parameters of the fish *Oreochromis mossambicus*. Int J Sci Res Publ. 2018;8:302-307.
36. Mandil R, et al. *In vitro* and *in vivo* effect of Flubendiamide and copper on cytogenotoxicity, oxidative stress and spleen histology of rats and its modulation by resveratrol, catechin, curcumin and α -tocopherol. BMC Pharmacol Toxicol. 2020;21:1-17.