

Prevalence of Parasitic Infections in Relation to CD4⁺ and Antiretroviral (ART) Usage of HIV Sero-Positive Patients Attending Irrua Specialist Teaching Hospital (ISTH) Irrua, Edo State, Nigeria

Iyevhobu KO* and Obodo BN

Department of Medical Laboratory Science, College of Medical Sciences, Ambrose Alli University, Ekpoma, Edo State Nigeria

Research Article

Received date: 07/05/2020
Accepted date: 18/05/2020
Published date: 25/05/2020

*For Correspondence

Department of Medical Laboratory Science, College of Medical Sciences, Faculty of Basic Medical Sciences, Ambrose Alli University, Ekpoma, Edo State, Nigeria.

E-mail: kennylamai@yahoo.com

Keywords: Parasites; Infections; CD4⁺; Antiretroviral Therapy

ABSTRACT

Intestinal parasitic infections are globally endemic and constitute the greatest single worldwide cause of illness and disease. This study was designed to determine the Prevalence of parasitic infections in relation to CD4⁺ and Antiretroviral Usage of HIV sero-positive patients Attending Irrua Specialist Teaching Hospital (ISTH) Irrua, Edo State, Nigeria. A total of two hundred (200) subjects from Irrua Specialist Teaching Hospital (ISTH), Irrua, were recruited for this study. They were categorised into two groups based on their HIV/AIDS clinical and laboratory test results. Based on this, 170(85.0%) were HIV sero-positive and 30 (15.0%) were HIV sero-negative subjects and were between the age ranges of 18-89 years and 18-59 years respectively. The parasites isolated were hookworms, *Strongyloides stercoralis*, *Trichuris trichuria*, *Ascaris lumbricoides* and *Enterobius vermicularis* with a prevalence of 17(10.0%), 7(4.1%), 4(2.4%), 16(9.4%) and 7(4.1%) respectively for the HIV sero-positive subjects while HIV sero-negative had a prevalence of 1(3.3%), 0(0.0%), 0(0.0%), 10(33.3%), 0(0.0%) for the same parasites respectively and in addition 1(3.3%) of *Schistosoma mansoni* was found respectively. The mean and standard deviation for the age of subjects were 36.4±15.92 and 30.1±10.26 for HIV sero-positive and HIV sero-negative subjects respectively. Parasitic infections were higher within the age range 30-44 for HIV sero-positive and 18-29 age range for HIV sero-negative. The prevalence of parasitic infections among the HIV sero-positive subjects was 44.7%, as compared to 40.0% of HIV sero-negative. A significant difference (p<0.05) was observed in the prevalence of opportunistic parasitic infections, between CD4⁺ count, ART use and age group.

INTRODUCTION

Intestinal parasitic infections caused by protozoans and helminths have been reported to be globally endemic and constitutes the greatest single worldwide causes of illness and disease [1,2]. In fact about one third of the world, more than two billion people are infected with intestinal parasite [1]. These intestinal parasites cause intestinal obstruction, malnutrition, iron deficiency anaemia, diarrhoea, malabsorption and other damage to the hosts [3]. It is estimated that as much as 60% of the World's population is infected with intestinal parasites which may play a significant role in morbidity due to intestinal infections [4]. The rate of infection is also remarkably high in Sub-Saharan Africa, where the majority of Human Immune Deficiency Virus (HIV)/Acquired Immunodeficiency Syndrome (AIDS) cases are concentrated where factors including poverty and malnutrition could promote transmission of both infections in the region [5]. Intestinal parasitic infections have a worldwide distribution with high prevalence found in people with low socio-economic status and poor living conditions as well as people in over-crowded areas with poor environmental sanitation, improper garbage disposal, unsafe water supply and unhygienic personal habits [6,7]. These factors are the causes of a major proportion of the burden of disease and death in developing countries [6]. Intestinal protozoa and faeco-oral transmitted helminths (STH) constitute major health problems, especially in the tropical and sub-tropical regions [8]. They can be

the cause of a wide spectrum of clinical problems ranging from apparently symptomless infections (**Figure 1**) to life-threatening conditions such as intestinal obstruction as seen in *Ascaris* infestation^[8,9].




Figure 1. Mixed infection of Hookworm and *Ascaris lumbricoides* (X40 Objectives).

Furthermore, Intestinal parasites are widely distributed in Nigeria largely due to the low level of environmental and personal hygiene, contamination of food and drinking water that results from improper disposal of human excreta^[10]. Intestinal parasites as a major concern in most developing countries have been pronounced with the co-occurrence of malnutrition and HIV/AIDS. Opportunistic parasitic infections are a common feature in HIV/AIDS infections where almost 80% of AIDS patients die of AIDS-related infections including intestinal parasites rather than of the HIV infection itself which usually occur late in the course of HIV infection when Cluster of Differentiation (CD4)⁺ T-cell count has been severely depleted mostly below 200 cells/mm³^[11,12]. In the absence of Anti-Retroviral Therapy (ART) HIV/AIDS patients in developing countries unfortunately continue to suffer the consequences of opportunistic parasites^[13]. Patients enrolling into ART programmes with very low CD4 cell counts have heightened risk of morbidity and mortality before ART^[14]. There is evidence that the control of these opportunistic parasitic infections in HIV-positive persons under Highly Active Anti-Retroviral Therapy (HAART) is also induced by the inhibition of the aspartyl protease of the parasites and by the reconstitution of the immune system of the patient^[15]. However, patients in resource limited settings typically start ART programmes with advanced symptomatic disease and very low blood CD4 cell counts which predisposes them to high rates of both clinical and subclinical opportunistic infections^[14]. Intestinal obstruction, anaemia, malnutrition, dysentery syndrome, fever, dehydration, vomiting and colitis are the major complications associated with soil transmitted helminthes infections^[16].

The clinical course of infection with larger parasites, such as the helminthes, can vary widely from asymptomatic to severe secondary complications and these asymptomatic hosts may experience several health effects depending on the number, type, and severity of infections present^[17,18]. Therefore, this study is set to assess the prevalence of intestinal parasites in Irrua Specialist Teaching Hospital (ISTH) Irrua. Parasitic infections are infections or opportunistic infections that affect persons who are immune-compromised, where HIV positive subjects are highly implicate^[1,2]. Therefore, it is important to determine the relationship between CD4⁺ count, ART and parasitic infections as this will help be of immense benefit to government and healthcare policies.

MATERIALS AND METHODS

Study area

This study was carried out in Irrua Specialist Teaching Hospital (ISTH), Irrua, the administrative headquarters of Esan Central Local Government Area of Edo State, Nigeria. Majority of the population are civil servants, traders, businessmen/women, transporters, farmers, teachers/lecturers and students. The samples were examined in the Research Diagnostic Laboratory, of the Department of Medical Laboratory Science, College of Medical Sciences, Ambrose Alli University, Ekpoma.

Study population

Two hundred (200) stool samples comprising one hundred and seventy (170) samples from HIV Sero-positive individuals and thirty (30) samples from HIV sero negative individuals were recruited from the study area. HIV sero positive individuals with other underlying health and who did not give consent were excluded from this study. HIV sero negative individuals with any health condition, exhibiting signs and symptoms of any illness, and who did not give consent were excluded from the study. These information were obtained from the medical records of these subjects.

Ethical approval

This was obtained from the Ethics and Research Committee of Irrua Specialist Teaching Hospital (ISTH), Irrua. Also, informed consent was sought from the subjects while explaining the aim and objectives, economic importance and Health benefits of the study to the subjects.

Collection of samples/ Data collection

Specimen containers (Wide Mouthed Screw Capped Plain Plastic Universal Containers), which were clean, dry and leak proof

^[19], and properly labelled with subjects' names, sex, age, and serial number were used for sample collection. The subjects were properly educated on how to collect the sample, the part of the stool sample to be collected and as well as the quantity needed. Early morning stool was advised to be taken. The samples were received the next morning with each subjects name, age, sex and serial number entered into the record book. The samples were then transported immediately to the laboratory for examination. Samples that could not be examined early enough were preserved with 10% formol saline (1 g of stool to 3 ml with 10% formol saline). Data of ART use and CD4⁺ count of the subjects were obtained from their respective registers in the Hospital. Stool samples were collected from those who are positive to HIV sero-test, after informed consent was obtained from them and they equally completed a structured questionnaire.

Examination and analysis of samples

The samples were examined using the Macroscopic and Microscopic methods. Samples were examined macroscopically for colour, consistency and constituents such as mucus, blood, pus, adult worms such as *Enterobius vermicularis* or *Ascaris lumbricoides* and tape worm segments. The consistency was used as a guide as to whether the trophozoites or the cyst stage, egg or worm of the parasite is likely to be present^[19]. The microscopic methods included the use of wet preparations (Saline and Lugol's iodine) and concentration methods. Iodinated formol-ether concentration method was employed as described by WHO,^[19]. Also, modified cold Ziehl-Neelsen's stain (for *Cryptosporidium* species) was employed as described by Current. Lymphocyte subset, CD4⁺ count was analyzed using FACScan flow cytometry (Becton Dickinson Immunocytometry system, and Jose, CA., USA).

STATISTICAL ANALYSIS

The percentage prevalence was calculated in each case and comparative analysis of the results was done using Chi-square. A p-value less than 0.05 (p<0.05) was considered statistically significant.

RESULTS

A total of 200 subjects from Irrua Specialist Teaching Hospital (ISTH), Irrua were recruited for this study. They were categorised into two group based on their HIV/AIDS clinical and laboratory test results. Based on this, 170(85.0%) were HIV sero-positive and 30(15.0%) were HIV sero-negative where the age ranges between 18-89 years for HIV sero positive and 18-59 years for HIV sero-negative. The results of the socio-demographic data revealed that the mean age of the subjects were 36.4±15.92 and 30.1±10.26 for HIV sero-positive and HIV sero-negative respectively. The majority of the subjects 112(56.0%) were female and 84(44.0%) were male (**Table 1**).

Table 1. Socio-demographic characteristic of the study subjects.

	HIV+n(%)	HIV-n(%)	Total n(%)	X ² p value
Sex				0.229(0.6322)
Male	76(44.7)	12(40.0)	88(44.0)	
Female	94(55.3)	18(60.0)	112(56.0)	
Age(Mean ± SD) years	36.4±15.92	30.1±10.26	35.3±15.21	
Residence				7.38(0.0066)
Urban	122(71.8)	14(46.7)	136(68.0)	
Rural	48(28.2)	16(53.3)	64(32.0)	
Marital Status				3.58(0.058)
Married	105(61.8)	13(43.3)	118(59.0)	
Not married	65(38.2)	17(56.7)	82(41.0)	
Occupation				19.067(0.0079)
Bankers	6(3.5)	0(0.0)	6(3.0)	
Traders	32(18.8)	8(26.7)	40(20.0)	
Civil Servants	39(22.9)	1(3.3)	40(20.0)	
House Wives	14(8.2)	2(6.7)	16(20.0)	
Students	47(27.7)	18(60.0)	65(32.5)	
Farmers	13(7.7)	1(3.3)	14(7.0)	
Drivers	13(7.7)	0(0.0)	13(6.5)	
Health Workers	6(3.5)	0(0.0)	6(3.0)	

Also, the results showed the age and sex distribution of HIV sero-positive and sero-negative of the study subjects (**Table 2**).

Table 2. Age and sex distribution of HIV sero-positive and sero-negative subjects.

Age (years)	Sex	Clinical Status of HIV+n (%)	Study Subjects HIV-n(%)
0-14	Male	7	3
	Female	10	5

15-29	Male	21	5
	Female	22	2
30-44	Male	23	6
	Female	34	3
45-59	Male	16	2
	Female	19	4
60-74	Male	6	0
	Female	7	0
75-89	Male	3	0
	Female	2	0
Total		170(85.0)	30(15.0)

The results on the parasites encountered among HIV sero positive and HIV sero negative subjects revealed that Hookworms with 17(10.0%) had a higher prevalence for helminthic infections while cryptosporidium species with a prevalence of 25(14.7%) was the only protozoan recorded for HIV sero-positive subjects. Also, *Ascaris lumbricoides* with a prevalence of 10(33.3%) was the most occurring helminthic infection (**Figure 2**) in HIV sero negative subjects and there was no protozoan isolated.




Figure 2. Multiple infection of *Acaris lumbricoides* (X10 Objectives).

The overall prevalence for parasitic infections for both HIV sero-positive and negative subjects were 88(44.0%). The overall prevalence of parasitic infections for HIV sero-positive was 76(44.7%) while HIV sero-negative was 12(40.0%) (**Table 3**).

Table 3. Parasites isolated among HIV sero-positive and negative subjects.

Parasite identified	HIV+	HIV-	Total	X2 (p value)
	n= 170 (%)	n= 30(%)	n= 200(%)	
Helminths				27.27(0.00011)
Hookworms	17(10.0)	1(3.3)	18(9.0)	
Strongloidesstercolaris	7(4.1)	0(0.0)	7(3.5)	
Trichuristrichura	4(2.4)	0(0.0)	4(2.0)	
Ascarislumbricoides	16(9.4)	10(33.3)	26(13.0)	
EnterobiumVermicularis	7(4.1)	0(0.0)	7(3.5)	
Schistosomamansonii	0(0.0)	1(3.3)	1(0.5)	
Protozoans				
Cryptosporidium species	25(14.7)	0(0.0)	25(12.5)	
Total	76(44.7)	12(40.0)	88(44.0)	

The results on the parasites encountered, as it relate with HIV status, ART status and CD4⁺count revealed that the prevalence of helminthic and protozoan infections for HIV sero positive subjects were 51(81.0%), and 25(100.0%) respectively while HIV sero negative subjects recorded prevalence of 12(19.0%) and 0(0.0%) for helminthic and protozoan infections respectively (**Figure 3**).




Figure 3. Multiple infections with *Cryptosporidium species*(X10 Objectives).

HIV sero positive subjects on ART recorded a prevalence of 22 (43.1%) for helminths and 9(36.0%) for protozoa infection while those not on ART recorded a prevalence of 29(56.9%) for helminthic infections and 16 (64.0%) for protozoa infection. As it relates to CD4⁺ count, the helminthic infection was 32(62.7%),11(21.6%) and 8(15.7%) for <200 cells/μL, 200-499 cells/μL and ≥ 500 cells/μL CD4⁺ count respectively while the protozoan infection was 18(72.0%), 5(20.0%) and 2(8.0%) for<200 cells/μL, 200-499 cells/μL and ≥ 500 cells/μL CD4⁺ counts respectively (**Table 4**).

Table 4. Parasite isolated as it relates with HIV Status, ART Status and CD4⁺ Count.

	Type of parasitic intestinal infection		
Characteristics	Helminthic infection	Protozoanic infection	X2 (p value)
HIV Status			5.51(0.019)
HIV+	51(81.0)	25(100.0)	
HIV_	12(19.0)	0(0.0)	
ART Status			0.35(0.552)
ON ART (31)	22(43.1)	9(36.0)	
NOT ART(45)	29(56.9)	16(64.0)	
CD4⁺ count (170) (Seropositive Only)			0.99(0.607)
<200 cells/μL	32(62.7)	18(72.0)	
200-499 cells/μL	11(21.6)	5(20.0)	
>500 cells/μL	8(15.7)	2(8.0)	

The results on the parasites encountered in relation to ART status showed that the prevalence of parasitic infections were 31(40.8%) for those on ART and 45(59.2%) for those not on ART (**Table 5**).

Table 5. Parasites isolated in relation to ART Status.

Parasite identified	On ART	Not on ART	X2 (p value)
Helminths			1.497(0.91)
Hookworms	7	10	
Strongloidesstercolaris	3	4	
Trichuristrichura	1	3	
Ascarislumbricoides	7	9	
EnterobiumVermicularis	4	3	
Protozoans			
Cryptosporidium species	9	16	
Total	31(40.8)	45(59.2)	

The results on the magnitude of parasitic infection as related to HIV sero positive and HIV sero negative subjects revealed that the HIV sero-positive and HIV sero-negative subjects had prevalence of 63(88.7%) and 8(11.3%) for light infections respectively,8(72.7%) and 3(27.3%) for medium infections respectively, 5(83.3%) and 1(16.7%) for heavy infections respectively, while 7(87.5%) and 1(12.5%) for mixed parasitic infections respectively. From the results it was observed that the individuals who were HIV sero-positive had the highest parasitic infection as compared to those of the HIV sero-negative subjects (**Table 6**).

Table 6. Magnitude of Parasitic infection as related to Sero-Positive and Negative Subjects.

Clinical status of the Study Subject	Magnitude of Infection			
	Light Infection (%)	Medium Infection (%)	Heavy Infection (%)	Mixed Infection (%)
HIV+ n=76	63(88.7)	8(72.7)	5(83.3)	7(87.5)
HIV- n=12	8(11.3)	3(27.3)	1(16.7)	1(12.5)
Total=88	71(74.0)	11(11.4)	6(6.3)	8(8.3)

The results on the prevalence of parasitic infections in relation to ART use and CD4⁺ Counts showed that those subjects with CD4⁺ count <200 had the highest prevalence with 50(65.8%) with prevalence of 16(21.1%) for CD4⁺ count of 200-499 and 10(13.1%) for CD4⁺ count of ≥500 (**Table 7**).

Table 7. Prevalence of Parasitic Infection in Relation to ART use and CD4+ Counts.

CD4+ cell/mm ³	On ART n(%)	No ART n(%)	Total n(%)	X ² (p value)
< 200	31(100.0)	19(42.2)	50(65.8)	
200-499	0(0.0)	16(35.6)	16(21.1)	27.23(0.0000012)
≥500	0(0.0)	10(22.2)	10(13.1)	
Total=170	31(40.8)	45(59.2)	76(44.7)	

DISCUSSION

Human Immune-deficiency Virus (HIV)/ AIDS pandemic has brought about a great change in intestinal fauna [10]. As the spectrum of immune-deficiency processes, HIV infected individuals become susceptible to a variety of opportunistic parasitic infection that occur with greater frequency and severity. Kumie [20], recorded that almost 80% of AIDS patients die from AIDS-related infections. Several intestinal parasites previously considered non-pathogenic in immune-competent individuals are opportunistically becoming aggressive and causing debilitating illness in HIV/AIDS patients [21]. In this present study, the majority of HIV sero-positive subjects were found in the age range 30-44 years, the most productive and reproductive group of the society, 76(44.7%) and 94(55.3%) of the study participants were males and females respectively, showing high HIV exposure rate in both sexes.

The prevalence of parasitic infections among the HIV sero-positive subjects was 44.7% and this in agreement with work of Sow et al., [22,23] and Zeynudin et al.[24], who reported 43.6%, 33.6% and 39.56% respectively. But it is in contrast to the work of Henry et al. [25], Therizol et al. [26], Babatunde et al., [27], and Inabo et al., [28], WHO reported 7.5%, 8.0%, 87.8 and 70.6% respectively. The reason while this study is in contrast to some work reported may be due to the sample size used, the advancement of technique use, distribution of parasitic infections geographically, the use of antiretroviral drugs(ART) and hygienic practices according to the report of WHO and CDC [29].

In terms of identified parasites specific prevalence, Cryptosporidium species which had a prevalence of 25(14.7%) was the highest occurring parasites and the only opportunistic parasites recorded in this study, while the prevalence of hookworms, *Strongyloides stercoralis*, *Trichuris trichiura*, *Ascaris lumbricoides* and *Enterobius vermicularis* were 17(10.0%),7(4.1%), 4(2.4%), 16(9.4%) and 7(4.1%) respectively for HIV sero-positive subjects. Also, HIV sero-negative subjects had prevalence of 1(3.3%), 0(0.0%), 0(0.0%), 10(33.3%), 0(0.0%) for hookworms, *Strongyloides stercoralis*, *Trichuris trichiura*, *Ascaris lumbricoides* and *Enterobius vermicularis* respectively and in addition 1(3.3%) of *Schistosoma mansoni* was isolated while no Cryptosporidium species were found in HIV sero-negative individuals. There was statistically significance (p-value=0.00011, p< 0.05) when the parasites isolated from HIV sero positive and negative subjects were compared. This is in agreement with report of CDC [29] were parasitic infections were found to be more significant with those who their CD4⁺ count is <200, and in this study the mean CD4⁺ count was less than 200 according to CDC classification.

Furthermore, parasitic infections were found among those that are not on ART with prevalence of 29(56.9%) and 16(64.0%) of helminthic and protozoan infection respectively, while those on ART had 22(43.1%) and 9(36.0%) respectively for helminthic and protozoan infections, and this comparison was not statistically significant (p-value=0.552, p>0.05), and this is in contrast with the work of Nigus [23] and Missage et al., [30] who reported that the difference was statistically significant with p-value (0.001) and (0.001), both p-values less than 0.05. On the account of CD4⁺, the helminthic and protozoan infections of those subjects whom their CD4⁺ count was <200, 200-499 and ≥500 were 32(62.7%),11(21.6%),8(15.7%) and 18(72.0%)5(20.0%), 2(8.0%) respectively and their difference in distribution was not significant (p-value= 0.607) and this result was in disagreement to the work of Akinbo et al., [31]and Nigus [23], who showed that there was significant difference in the CD4⁺ count with p-values of 0.0001 and 0.005 in there independent study.

On the distributions of parasites species for HIV sero-positive subjects, on ART and Not on ART, it was found that the distributions was not statistically significantly (p-value=0.9, p>0.05). This may not be unconnected to the fact that the area of study is endemic to the parasites isolated and the fact that most of those that have not started ART and those on ART's mean CD4⁺ count was between <200 to ≤250 cells/μL. There were more mixed parasitic infections i.e polyparasitism with HIV sero-positive than HIV sero-negative with 7(87.5%) and 1(12.5%) respectively.

In terms of magnitude of parasitic infection of HIV sero-positive and HIV sero-negative subjects, there was prevalence of

63(88.7%), 8(72.7%), 5(83.3%) and 8(11.3%), 3(27.3%), 1(16.7%) recorded for light, medium and heavy parasitic infection respectively. This is in agreement with the work of Missage et al.,^[30,31] Therizol et al.^[26], Babatunde et al.,^[27], and Inabo et al.,^[28] who reported same but in contrast with work of Nigus^[23], who reported differently, reason be that most of Nigus's^[23], HIV sero-positive subjects were on Antiretroviral(ART) drugs, which improved their immunity and fight most parasitic infections which most times are opportunistic infections. When the distribution of parasitic infections from subjects on ART and not on ART was correlated with CD4⁺ count, it was found to be statistically significant (p-value=0.0000012, p<0.05) and the correlation was also significant, this finding equally agreed with the works of Missage et al.^[30,31].

CONCLUSION

It is evident from this report that parasitic infections were highly prevalent among the HIV infected patients. This was particularly so in patients with very low CD4⁺ cells count and those that are not on ART. The majority of the infections were prevalent among the age range 30-44. The prevalence of opportunistic infection among those of HIV sero-positive was 25(14.7%). A significant difference (p<0.05) was observed in the prevalence of opportunistic parasitic infection, between CD4⁺ count, ART use and age group.

CONTRIBUTION TO KNOWLEDGE

Results from this study will help to assess the prevalence of intestinal parasite infection among HIV sero-positive in the study area. This will help to create awareness on the prevalence rate for the public to recommend the best management and preventive measures to reduce the rate of infection and transmission via personal hygiene, better public awareness and public policies. It will also help the care giver to manage patients effectively and emphasis on clean environment in the prevention and control of parasitic infections among the study population when the work finally published.

ACKNOWLEDGMENT

Thanks to all who contributed to the success of this research and the presentation of this manuscript. Special appreciation goes to Mr. Iyevhobu Kenneth Oshiohayamhe and Mr. Obodo Basil Nnaemeka, Chief Researchers in St Kenny Consult, Ekpoma, Edo State, Nigeria for taking their time to prepare, arrange and make the required corrections in this work.

REFERENCES

1. Chan MS. The global burden of intestinal nematode infections – Fifty years on. *Parasitol.* 1997; 13: 438-443.
2. Pillai DR, et al. Common Intestinal Parasites. *Curr Treat Opt Infect Dis.* 2003; 5: 207–217.
3. Buchini MLT, et al. Spatial distribution of enteroparasites among school children from Guarapuava, State of Paraná, Brazil. *Revista Brasileira de Epidemiologia.* 2007; 10: 568-578.
4. World Health Organization (WHO) Prevention and control of intestinal parasitic infections, WHO Technical Report, 1987; 749:1–86.
5. UNAIDS/WHO (2002): HIV Epidemic Update. Geneva: UNAIDS.
6. Adamu H, et al. Prevalence of intestinal parasite. *Ethiop J Health Div.* 2006; 20: 39–47.
7. Noor Azian MY, et al. Prevalence of intestinal protozoa in an aborigine community in Pahang, Malaysia. *Trop Biomed.* 2007; 24: 55-62.
8. Savioli L, et al. Intestinal parasitic infections: a soluble public health problem. *Trans R Soc Trop Med Hyg.* 1992; 86:353–354.
9. Glickman LT, et al. Nematodes intestinal parasites in children in rural Guinea, Africa: Prevalence and relationship to Geophagia. *Int J Epidemiol.* 1999; 28: 169–174.
10. Uzairue LI, et al. Gastrodiscoides Hominis Infestation On Vegetables (Cabbages) Sold In Ekpoma Markets, Edo State, Southern Nigeria- A Case Report. *IJB AIR; 2013; 2: 37-39.*
11. Kam KM. Gut parasites in general practice in Hong Kong, *J Trop Med Hyg.* 1994; 97:117–120.
12. Kelly P. Diarrhoea and AIDS: recent developments in the African settings. *African Hlth.* 1998; 1:16–18.
13. Maggi P, et al. Effect of antiretroviral therapy on cryptosporidiosis and microsporidiosis in patients infected with human immunodeficiency virus type 1. *Eur J Clin Microbiol Infect Dis.* 2000; 19:213–217.
14. Lawn SD, et al. Early mortality among adults accessing a community-based antiretroviral service in South Africa: implications for programme design, *AIDS.* 2005; 19:2141–2148.
15. Willemot P, et al. Prevention of HIV-associated opportunistic infections and diseases in the age of highly active antiretroviral therapy, *Expert Rev Anti Infect Ther.* 2004; 2:521–532.

16. Cooper E. Intestinal parasitoses and the modern description of diseases of poverty. *Trans R Soc Trop Med Hyg.* 1991; 85: 168-170.
17. Higgins DA, et al. Human intestinal parasitism in three areas of Indonesia: a survey. *Ann Trop Med Parasitol.* 1984; 78: 637-648.
18. Checkley W, et al. Asymptomatic and symptomatic cryptosporidiosis: their acute effect on weight gain in Peruvian children. *Ame J Epidemiol.* 1997; 145: 156-163.
19. Cheesbrough M. Parasitological Test. In: *District Laboratory practice in Tropical Countries.* 2nd edition, part 1 (Low Price Ed), Cambridge University Press, 2006; 5:183-215.
20. Kumie A, et al. An overview of environmental health status in Ethiopia with particular emphasis to its organization, drinking water and sanitation: a literature survey, *Ethiop J Health Dev.* 2005; 19:89-103.
21. Gomez MMA, et al. Opportunistic and non-opportunistic intestinal parasites in HIV-positive and negative patients with diarrhoea in Tanzania. *Trop Med Parasitol.* 1995; 46: 109-111.
22. Sow PG, et al. About Intestinal Parasitic Infections in a Cohort of HIV-Infected Patients, *Adv Bioesrch.* 2012; 3: 32 - 35.
23. Nigus FT. Intestinal parasitic infection among HIV sero-positive and sero-negative adults patients presented with diarrhea in Gondar, Northwest Ethiopia, a paper presented at the interes workshop on 25-28 May, 2010, in Mapulo Mozambique in Gondar, Northwest.
24. Zeynudin A, et al. Prevalence of opportunistic intestinal parasitic infection among HIV infected patients who are taking antiretroviral treatment at Jimma Health Center, Jimma, Ethiopia. *Euro Review for Medl and Pharmacolo Sci.* 2013; 17: 513-516.
25. Henry MC. Parasitological observation of chronic diarrhoea in suspected AIDS adult patients in Kinshasa (Zaire), *Trans Rcoc Trop Med Hyg.* 1986; 80: 309-310.
26. Therizol FM. Chronic diarrhoea and parasitosis in adult patients suspected to have AIDS in Ivory Coast., *bull de la soc de path exo et de sel92(Suppl 2).* 1989; 690-693.
27. Babatunde SK, et al. Prevalence of intestinal parasitic infestation in HIV seropositive and seronegative patients in Ilorin, Nigeria. *Ann Afr Med.* 2010; 9: 123-128.
28. Inabo H, et al. Profile of Intestinal Parasitic Infections Associated with Diarrhoea in HIV/AIDS Patients in a Tertiary Hospital in Zaria, Nigeria. *World J Life Sci Med Res.* 2012; 2: 43-47
29. CDC. Control measures for Outbreaks – Intensified Cryptosporidiosis (Crypto). *Parasitol Res.* 5: 89-90.
30. Missage A, et al. Prevalence of intestinal parasites and associated risk factors among HIV/AIDS patients with pre-ART and on-ART attending dessie hospitalART clinic, Northeast Ethiopia, *AIDS Res Thpy.* 2013; 10: 7
31. Akinbo FO, et al. Prevalence of intestinal parasites in relation to CD4 counts and anaemia among HIV infected patients in Benin City, Edo State, Nigeria, Tanzania. *J Hlth Reserch.* 2011; 13: 12-32.