

Technical Sheet of Process of Attieke Production in Cote Divoire

Kouame AK*, Jean-Paul KMB, Boli ZA, Djeni N and Marcellin KD

Department of Food Sciences and Technology, Laboratory of Biotechnology and food Microbiology,
University Nangui Abrogoua, Abidjan, Cote Divoire, West Africa

Research Article

Received Date: 26/09/2017

Accepted Date: 30/09/2017

Published Date: 05/10/2017

*For Correspondence

Alfred K Kouame, Department of Food Sciences and Technology, Laboratory of Biotechnology and food Microbiology, University Nangui Abrogoua, Abidjan, Cote Divoire, Tel: +22509994291.

E-mail: kohialfred2@yahoo.fr

Keywords: Attieke, Process, Cote divoire

ABSTRACT

Attieke is the major fermented plant food in Côte d'Ivoire. It is a steamed granular cassava (*Manihot esculenta* Crantz) meal, couscous-like product, with slightly sour taste and whitish colour. Our study is a technical sheet of process of attieke production in cote d'Ivoire. process of attieke production in côte d'Ivoire was : Cassava roots, peeling, pulp washing, grinding, fermentation, wrinding, sieving, drying, winniwing and fibre removal, cooking, packaging and attieke. Attieke not contained coliforms, moulds and anaerobic sulfite reducing bacteria, Salmonella and Echerichia coli. However, the values obtained fall within the standard specifications set for attieke by CODINORM (Côte d'Ivoire standards Board)

INTRODUCTION

Cassava, the enlarged root of *Manihot esculenta* Crantz, is an important staple food for about 80% of Cote d'Ivoire's estimated population, especially those living in the southern regions^[1]. Cassava may be processed by boiling, roasting, drying, or by fermentation, depending on the variety^[2] before eaten. However, the most popular processing method, especially for the varieties which are high in the cyanogenic glucosides, is by fermentation. In Africa, over 90% of the processed cassava are consumed as fermented products^[3], and one of the most popular fermented foods derived from cassava in Cote d'Ivoire is *attieke*. *Attieke*, a steamed cassava fermented semolina, is one such fermented cassava product and is of significant importance for an increasing number of people in Cote d'Ivoire^[4] and other countries in the world. Recent data on attieke consumption do not exist, but Aboua et al.^[1] estimated the consumption between 28,000 and 34,000 tons per year, the equivalent of 40,000 and 50,000 tons of fresh cassava. The popularity of *attieke* to urban dwellers is associated with its cheapness, lower bulk (as compared to other cassava product) and its characteristic of ready to eat food. The processing of cassava into *attieke* needs several and hard steps. Our study is a technical sheet of process of *attieke* production in Cote d'Ivoire.

MATERIALS AND METHODS

Material Vegetable

The amino acid that has been shown to affect brain neurotransmission most readily is tryptophan. It is an essential amino acid and precursor of neurotransmitter serotonin and hormone melatonin. Central serotonergic system is also involved in the regulation of other mood status, including depression. However, a single tryptophan deficient meal has been clearly shown to produce transient depression in normal human. It produces hypnotic like effect, accordingly, it could be considered as a drug not a food.



Figure 1. Cassava roots.

The used plant material consisted of 12 months-old freshly harvested cassava roots of the bitter variety, IAC cultivar (Improved African Cassava) (Figure 1).

Attieke Preparation

Roots are peeled, cut into pieces and then washed three times with fresh water. The milling takes place in a cooperative mill located in the village. Before milling, 5–10% (w/w) of inoculum, 10% (v/w) water and about 01% (v/w) of palm oil are added and the pieces are ground to a fine paste, which is placed in large bowls. The mash is left to ferment for about 12–15 hours at ambient temperature (30–37 °C). After fermentation, the mash is placed in a jute sack and pressed continuously in a hand press for an hour. The press cake is then passed through two sieves to obtain a fine powder. The grains are formed by shaking and rotating the powder in a large bowl. The grains are sun-dried on black plastic canvas or flat bowls for a time period ranging from a few minutes up to half an hour. After drying, fibres and dirt are removed by sprinkling the grains. The grains are poured onto the sieve up to a height of 15–20 cm for steaming for about 20–25 hours on a cauldron filled with boiling water. Attieke obtained is the filled into plastic bags, sealed airtight and sold on local markets or transported in cars at ambient temperature (30–37 °C) in other localities (Figures 2-8).




Figure 2. Process of attieke production in Cote d'Ivoire.



Figure 3. Crushed cassava dough.




Figure 4. Pressed cassava paste.



Figure 5. Cassava semoula.




Figure 6. Cooking of semoulas.



Figure 7. Attieke.



Figure 8. Attieke packed.

Determination of pH and Total Titrable Acidity (TTA)

Thirty grams of cassava traditional inocula samples were blended with 70 ml of sterile distilled water and filtered through a Whatman filter paper. The pH of 30 ml of the filtered solution was determined using a pH-meter (pH-meter P107, Consort, BioblockScientific, Illkirch, France). TTA was determined using the standard method described by Amoa-Awua et al. [5]. Ten millilitres of filtered solution were titrated with NaOH 0.1 N, using 1% phenolphthalein as indicator. The volume of aliquot used was recorded to determine the amount of acid in the sample. The titrable acidity was calculated as percentage of lactic acid. The determinations were carried out in triplicates and the mean value recorded.

Enumeration and Identification of Spoilage Microorganisms

Preparation of stock solutions, inoculation of agar plates, cultivation and quantification of micro-organisms were carried out according to [6]. For all determinations, 10 g of the traditional cassava inocula samples was homogenized in a stomacher with 90 ml of sterile buffered peptone water (AES Laboratoire, Combourg, France). Tenfold serial dilutions of stomacher fluid were prepared and spread plated for determination of micro-organism counts. Enumeration of coliforms was carried out using plates of Violet Red Bile Lactose agar (VRBL, Merck 10660, Merck, Darmstadt, Germany). The cultures were incubated for 48 h at 30 °C for total coliforms and 44 °C for faecal coliforms. Yeasts and moulds were enumerated on plates of Sabouraud–chloramphenicol agar (Fluka, Bochemica 89579, Sigma-Aldrich Chemie GmbH, Bangalore, India), incubated at 30 °C for 4 days. Aerobic mesophiles were enumerated on plates of plate count agar (PCA Oxoid Ltd, Basingstoke, UK) and incubated at 30 °C for 2 days.

Isolation and Identification of Food-borne Pathogens

Staphylococcus aureus

Staphylococcus aureus was isolated and enumerated according to the method described by Capita et al. [7]. A volume of 0.1 ml of each dilution was surface plated on Baird-Parker agar (BPA) containing egg yolk tellurite emulsion (Oxoid) and incubated at 37 °C for 24 and 48 h. The total number of colonies, colonies with different morphology to those of *Staphylococcus aureus* was counted. Five colonies from each sample were randomly selected, purified and tested for cell morphology, arrangement of the cells, Gram reaction, catalase activity, oxidase test, ability to produce acid anaerobically in a glucose-containing growth medium, coagulase activity, thermo-stable nuclease activity, acid production from mannitol and acetoin production. Only, the gram positive cocci were identified using the identification schemes proposed by Schleifer [8]. After the identification, the percentages of *Staphylococcus aureus* and the other strains were calculated. These percentages were later used to correct the results of the counts obtained from each BPA plate.

Bacillus

The quantitative estimation of spores of *B. cereus* was performed by a standard plate-counting method. Isolations were achieved from heat-treated dilutions by plating on mannitol egg yolk polymyxin B agar [9]. Presumptive colonies of *B. cereus* were randomly selected based on characteristic colony feature, purified on the same medium and identified by morphological, cultural and biochemical characteristics according to the documented procedures [10].

Salmonella

The research of *Salmonella* in cassava traditional inocula, palm oil and water samples were achieved according to the procedure described in the global *Salmonella* surveillance and laboratory support project of the World Health Organization [11]. From each sample, 25 g was aseptically weighed and macerated in 225 ml of buffered peptone water (Oxoid) and incubated at 37 °C for 24 h. A selective enrichment in Tetrathionate broth (Muller-Kauffmann) and Rappaport Vassiliadis soy peptone broth using 1 ml of previously incubated buffered peptone water was achieved at 37 °C for 24 h, followed by a subcultivation on *Salmonella* Shigella agar incubation at 35 °C for 24–48 hours [12]. Colourless, transparent and with a black centre colonies were further identified using biochemical tests.

Statistical Analysis

Descriptive statistics for microbiological data were calculated with Excel (Microsoft, Redmond, WA, USA). All statistical

analyses were implemented in STATISTICA for Windows ver. 10 (Statsoft Iberica, Lisbon, Portugal). Parametric tests (one-way variance analysis with Duncan's test) at 5% significance level were performed to determine whether there were significant differences between markets regarding microbiological data collected.

RESULTS AND DISCUSSION

In Côte d'Ivoire; *attieke* plays an important role in the population diet. It is part of the diet of many peoples. It is a typically Ivorian food, whose annual local consumption is estimated at over 450000 tons^[6]. *Attieke* production in Côte d'Ivoire is usually prepared by method described above (**Figure 1**). *Attieke* just after cooking and *attieke* packed had acidic pH. The production of *attieke* depends on a fermentation step which gives an intermediate product (fermented paste) of acid pH. Other unit operations that result in the finished product do not cause significant pH changes^[13]. pH values were 4.37 and 4.36, respectively. The titratable acidity values were respectively 2.44% and 1.78%. However, the values obtained fall within the standard specifications set for *attieke* by CODINORM (Côte d'Ivoire standards Board) (4–5 for pH) and are also similar to the results of Coulin et al.^[14]. Just after steaming and packaging in plastic bags, *attiéké* did not contain coliforms, moulds and anaerobic sulfite reducing bacteria, *Salmonella* and *Escherichia coli* (**Table 1**).

Table 1. pH, total titratable acidity and microbial population in cassava traditional inocula used in *attieke* process.

Parameters	Values
pH	4.94 ± 0.8
Titratable acidity (%)	1.67 ± 0.2
Aerobic mesophiles (CFU.g ⁻¹)	Ab
Moulds (CFU.g ⁻¹)	Ab
<i>Staphylococci</i> (CFU.g ⁻¹)	Ab
<i>Bacilli</i> (CFU.g ⁻¹)	(1.41 ± 3.2)10 ³
Total coliforms (CFU.g ⁻¹)	Ab
Faecal coliforms (CFU.g ⁻¹)	Ab
<i>Escherichia coli</i>	Ab
<i>Salmonella</i> (CFU.g ⁻¹)	Ab

The absence of *Salmonella*, *E. coli* and anaerobic sulfite reducing bacteria in *attieke* samples could be due to the low pH. In fact, the combined effect of organic acids produced during the fermentation period may possibly exert bacteriostatic effect on spoilage organisms and pathogens that might be present^[15,16]. They all contained *Bacillus* spore at mean loads of (1.41 ± 3.2)10³. Due to the low level of microbial detection, these *attieke* samples were of satisfying quality in regard to the standards recommended by CODINORM.

CONCLUSION

Attieke is a typically Ivorian food. Its production necessarily involves stages of production that pass from the roots of cassava to the cooking of the semolina. The *attieke* obtained at an acid pH and is of satisfactory microbiological quality.

ACKNOWLEDGEMENT

This work was supported by the International Foundation for Science (IFS) under Grant E/4955-1. The authors gratefully acknowledge all the women *attieke* producers. There is no conflict of interest concerning this manuscript.

REFERENCES

1. Aboua F, et al. Analysis of some constituents of cassava during the preparation of the *attieke*. In The Post-Harvest in Africa: Seminar International Abidjan. Montmagny QC Marquis Publishers, Ivory Coast. 1990;217-221.
2. Kouadio NA, et al. Etude comparative des méthodes traditionnelles de la preparation de l'*attieke* dans le Sud de la Côte d'Ivoire. Cahiers de la recherche scientifique et technique. 1991;108:703-706.
3. Westby A. Cassava utilization, storage and small-scale processing. In Cassava: Biology, Production and Utilization. CAB International, Wallingford, UK. 2002;281-300.
4. Assanvo B, et al. (2006) Microflora of traditional starter made from cassava for "attieke" production in Dabou (Cote d'Ivoire). Food Control. 2006;17:37-41.
5. Amoa-Awua WK, et al. Lactic acid fermentation of cassava dough into agbelima. Int J Food Microbiol. 1996;31:87-98.
6. Djéni NT, et al. Quality of *attieke* (a fermented cassava product) from the three main processing zones in Cote dlvoire. Food Res Int. 2011;44:410-416.

7. Capita R, et al. Assessment of Baird-Parker agar as screening test for determination of *Staphylococcus aureus* in poultry meat. *J Microbiol.* 2001;390:321-325.
8. Schleifer KH and Kloos WE. Isolation and characterization of *staphylococci* from human skin. *Int J Syst Bacteriol.* 1975;25:50-61.
9. Kouame AK, et al. Postprocessing microflora of commercial *attieke* (a fermented cassava product) produced in the south of Cote d'Ivoire. *Letters in Applied Microbiology.* 2012;56:44-50.
10. Cappucino JG and Sherman N. *Microbiology: A Laboratory Manual.* 6th ed. Person Education, Singapore; 2004.
11. Hendriksen RS. *Laboratory protocols level 1: Training course isolation of salmonella.* A Global *Salmonella* Surveillance and Laboratory Support Project of the World Health Organization, 4th ed. WHO, Geneva; 2003.
12. Feng P, et al. Enumeration of *Escherichia Coli* and the coliform bacteria. *Bacteriological Analytical Manual,* 8th ed. FDA/ Center for Food Safety and Applied Nutrition, USA; 2007.
13. Kouame AK. Identification des dangers et des des points critiques de contrôle pour la mise en place d'un système HACCP pour la production de l'*attieke* en Côte d'Ivoire. Thèse de doctorat unique Université Nangui Abrogoua, UFR des Sciences et technologies des Aliments, 2013;210.
14. Coulin P, et al. Characterisation of the microflora of *attieke*, a fermented cassava product during traditional small-scale production. *International Journal of Food Microbiology.* 2006;106:131-136.
15. Sengun IY and Karapinar M. Microbiological quality of Tarhana, Turkish cereal based fermented food. *Qual Assur Safety Crop Food.* 2012;4:17-25.
16. Tomkins A, et al. L'emploi de produits fermentés pour améliorer l'alimentation des enfants d'Afrique australe et orientale. In: *Pour Améliorer l'Alimentation des Jeunes Enfants en Afrique Orientale et Australe: une Technologie à la Portée des Ménages* eds Alnwick D. 1987;156-192.