

# Research & Reviews: Journal of Medical and Health Sciences

## Hypertension: Causes, Prevention and Treatment

Kavya SR\*

Department of Biotechnology, Sapthagiri College of Engineering, Visvesvaraya Technological University, India

### Review Article

Received: 25/01/2015  
Revised: 09/02/2015  
Accepted: 25/02/2015

#### \*For Correspondence

#14/5, Sapthagiri College of Engineering Chikkasandra, Hesaraghatta Main Road  
Bangalore – 560057, India.

Hypertension, likewise alluded to as hypertension, is a condition in which the corridors have perseveringly lifted pulse. Each time the human heart pulsates, it pumps blood to the entire body through the conduits.

Circulatory strain is the power of blood pushing up [1] against the vein dividers. The higher the weight the harder the heart needs to pump [2].

Hypertension can prompt harmed organs, and a few diseases, for example, renal disappointment [3] (kidney disappointment), aneurysm, heart disappointment, stroke, or heart assault [4]. Specialists from UC Davis reported in the Journal of the American Academy of Neurology that hypertension amid middle age may raise the danger of intellectual decrease sometime down the road [5].

The typical level for circulatory strain is underneath 120/80[6], where 120 speaks to the systolic estimation (crest weight in the courses) [7] and 80 speaks to the diastolic estimation (least weight in the veins) [8]. Circulatory strain between 120/80 and 139/89 is called prehypertension (to indicate expanded danger of hypertension) [9], and a pulse of 140/90 or above is considered hypertension [10].

Hypertension may be named key or optional. Vital hypertension is the term for hypertension with obscure reason. It represents around 95% of cases [11]. Auxiliary hypertension is the term for hypertension with a known direct cause, for example, kidney illness [12], tumors, or contraception pills [13].

Practically 70 million grown-ups in the United States are influenced by hypertension [14]. The condition likewise influences around two million teenagers and youngsters. As indicated by a report issued by the Centers for Disease Control and Prevention (CDC) in September 2012 [15], over a large portion of all Americans with hypertension don't have their hypertension under control.

#### Causes

Despite the fact that the definite reasons for hypertension are generally obscure [16], there are a few variables that have been very connected with the condition. These include:

- Smoking
  - Obesity or being overweight
  - Being stout/overweight as a kid
  - Diabetes
  - Sedentary way of life
  - Lack of physical action
  - High levels of salt admission (sodium affectability).
  - Insufficient calcium, potassium, and magnesium utilization
  - Vitamin D lack

- High levels of liquor utilization
- Stress
- Aging
- Medicines, for example, conception prevention pills
- Genetics and a family history of hypertension (a few qualities in the kidneys may add to hypertension)
- Chronic kidney sickness
- Adrenal and thyroid issues or tumors.

Measurements in the USA demonstrate that African Americans have a higher occurrence of hypertension than different ethnicities [17-21].

### **Side effects of Hypertension**

There is no ensure that a man with hypertension will introduce any indications of the condition. Around 33% of individuals really don't have the foggiest idea about that they have hypertension [22], and this obliviousness can keep going for a considerable length of time. Consequently, it is prudent to experience occasional pulse screenings notwithstanding when no indications are available [23].

Amazingly hypertension may prompt a few indications [24,25] are as follows:

- Severe migraines
- Fatigue or perplexity
- Dizziness
- Nausea
- Problems with vision
- Chest torments
- Breathing issues
- Irregular pulse
- Blood in the pee.

### **Hypertension Diagnosis**

Hypertension may be analyzed by a wellbeing proficient who measures circulatory strain with a gadget called a sphygmomanometer - the gadget with the arm sleeve, dial, pump, and valve [26]. The systolic and diastolic numbers will be recorded and contrasted with a graph of qualities. In the event that the weight is more noteworthy than 140/90, you will be considered to have hypertension.

A hypertension estimation, on the other hand, may be spurious or the consequence of anxiety at the season of the exam [27]. With a specific end goal to perform a more exhaustive analysis, doctors for the most part direct a physical exam and request the restorative history of you and your gang. Specialists will need to know whether you have any of the danger components for hypertension, for example, smoking, elevated cholesterol, or diabetes [28].

In the event that hypertension appears to be sensible, tests, for example, electrocardiograms [29] and echocardiograms will be utilized as a part of request to quantify electrical action of the heart and to evaluate the physical structure of the heart. Extra blood tests will likewise be obliged to recognize conceivable reasons for auxiliary hypertension and to quantify renal capacity, electrolyte levels, sugar levels, and cholesterol levels [30].

### **Hypertension consequences for different sicknesses**

#### ***Extreme immune system hyperthyroidism***

Immune system thyroid infection has been depicted in patients with pneumonic blood vessel hypertension (PAH) [31]. In any case, the season of onset and sort of thyroid ailment connected with PAH is variable. Case reports recommend a higher pervasiveness of hypothyroidism than hyperthyroidism in patients with PAH [32,33]. Epoprostenol, an endothelium-determined prostacyclin is a systemic and pneumonic vasodilator that hinders platelet total through expanded intracellular cyclic adenosine

monophosphate (cAMP) movement [34]. Prostacyclins have antiproliferatives, antithrombotics and immunomodulatory movement and are much of the time utilized as a part of the treatment of PAH. Start of epoprostenol has been accounted for with thyroid infection in a few studies [35]. We portray two instances of intense hyperthyroid immune system malady in already euthyroid patients who were begun on epoprostenol for PAH. The seriousness of hyperthyroidism brought about exacerbating aspiratory/cardiovascular status and fast development of a compressive goiter in one patient proposing that thyroid capacity ought to be observed nearly in PAH patients.

### ***Intense pneumonic embolism***

Endless thromboembolic pneumonic hypertension (CTEPH) is the consequence of clump constancy and fibrosis in the aspiratory vasculature after intense aspiratory embolism. This condition prompts lifted pneumonic weights and in the end right heart disappointment. Patients are frequently extremely constrained by exertional dyspnea and can encounter huge horribleness and mortality when left untreated [36-38]. The condition is likewise connected with disability of left ventricular diastolic capacity and filling. Pneumonic thromboendarterectomy (PTE) is as of now the main remedial treatment for CTEPH and significantly changes the clinical course of the infection.

Depicted in subtle element by Moser, et al. in 1987, PTE has significantly enhanced CTEPH-related mortality [39-40]. The quick hemodynamic and echocardiographic changes taking after PTE have given a perfect model to evaluating the association between the left and right heart. While early studies concentrated on the privilege ventricular brokenness seen in CTEPH, later studies have tended to the related left ventricular (LV) brokenness. This article highlights two of the most recent echocardiographic productions in CTEPH, one which addresses right ventricular capacity utilizing strain imaging, and the other which investigates the part of left atrial volume in the related LV diastolic disability.

### ***Renal capacity of females with gestational diabetes***

Renal disappointment is one of the main sources of mortality. Diabetes mellitus is the most widely recognized reason for end-stage renal infection and it has been evaluated that 366 million individuals will diabetes mellitus by 2030 [41]. Diabetes frequently coincides with a bunch of other strong cardiovascular danger variables including hypertension [42]. The joined vicinity of hypertension and diabetes associatively quickens the reduction in renal function [43]. Gestational diabetes mellitus (GDM) is characterized as any level of glucose narrow mindedness withonset or first acknowledgment amid pregnancy [44]. Approximately 7% of all pregnancies are entangled by GDM, bringing about more than 200,000 cases annually [45]. Women with a background marked by gestational diabetes are at expanded danger of cardiovascular and kidney illness, paying little respect to whether they have gone ahead to create plain diabetes, developing proof proposes [46]. Hypertensive issue of pregnancy confuse [5-10] precents of all pregnancies and can bring about a mixture of maternal and fetal complexities, including renal disappointment [47]. In a considerable lot of the females, hypertension may coincide with gestational diabetes or it might be available independently. In spite of the fact that the hypertension and diabetes mellitus are the known danger variables for kidney disappointment [48], which of these clinical conditions disturb renal capacity all the more extremely in females is not clear. Consequently this study was embraced to assess the impact of hypertension, gestational diabetes with and without hypertension on renal capacity in pregnant ladies.

### ***Cardiovascular danger***

Blood vessel hypertension is viewed as a critical danger component for the advancement of cardio and cerebral-vascular pathologies [49], and is one of the issue with the most noteworthy pervasiveness in cutting edge age [50]. In Italy, 31% of the overall public experiences hypertension and 17% is marginal [51].

Couple of studies have inspected the relationship in the middle of work and wellbeing state, particularly considering delayed working life and the vicinity of constantly more seasoned subjects at work [52]. A few studies considering these components have demonstrated a high recurrence of blood

vessel hypertension in particular working gatherings [53]. However, autonomously from age and other additional expert danger conditions, some work conditions are related to a high danger of creating blood vessel hypertension [54]. In a few studies, blood vessel hypertension has been related all the more regularly with "blue collars" than with "white collars" [55], a positive affiliation is likewise present in transport and cabbies [56], seriously higher contrasted with mechanical support experts [57]. Laborers with composition to clamor better than 85 dB (A) for more than 30 years have demonstrated a danger of hypertension 1.5 times higher [58-60].

### ***Expanded body mass***

Hypertension is a noteworthy cardiovascular ailment and is in charge of lethal cardiovascular results, for example, stroke, myocardial dead tissue, heart disappointment and renal disappointment [61,62]. Hypertension is exceptionally common among blacks for reasons not yet clear [63-65]. In Nigeria, the predominance of hypertension and its related intricacies are expanding at a disturbing rate [66]. The predominance relies on upon the area of the study and the cut-off point for meaning of hypertension. Lower predominance was found in studies that utilized BP 160/95 mmHg contrasted and BP 140/90mmHg as meaning of hypertension [67,68] In Meta-investigation by Ekwunife et al. [69], the pervasiveness ran somewhere around 12.4% and 34.8% while learn at the group in the Eastern piece of Nigeria by Ejim et al. [70] demonstrated that 46.4% of the study populace had hypertension. The commonness is higher in guys contrasted with females.

The etiology of hypertension is complex and it is getting to be clearer that the ascent in pervasiveness of hypertension in Nigeria can't be totally clarified by racial or hereditary programming. It is of note that while mulls over on hereditary determinants of hypertension are at different phases of conclusion overall including Nigeria [71-75], the part of natural elements, for example, physical idleness, heftiness, and smoking can't be disregarded.

Accompanying with the expanding predominance of hypertension is the bounce in pervasiveness of heftiness [76,77]. In reality relationship between expanded body mass and hypertension and diabetes is entrenched in the writing [78]. Corpulence is a standout amongst the most essential danger components for hypertension and cardiovascular ailment [79]. The worldwide heftiness pandemic has been ascribed to expanding inactive way of life and horrible dietary pattern, particularly in the western created nations [76]. A few physiologic systems including actuation of the reninangiotensin- aldosterone and thoughtful sensory system, endothelial brokenness and renal capacity anomalies have been cited for the relationship in the middle of stoutness and hypertension [80].

### ***Hypertension Treatment***

The primary objective of treatment for hypertension is to lower circulatory strain to under 140/90 - or even lower in a few gatherings, for example, individuals with diabetes, and individuals with ceaseless kidney maladies [81]. Treating hypertension is vital for diminishing the danger of stroke, heart assault, and heart disappointment.

Hypertension may be dealt with medicinally, by changing way of life components, or a blend of the two. Imperative way of life changes incorporate getting more fit, stopping smoking, eating an energizing eating routine, decreasing sodium admission, practicing consistently [82], and constraining liquor utilization.

Medicinal choices to treat hypertension incorporate a few classes of medications. Pro inhibitors, ARB drugs, beta-blockers, diuretics, calcium channel blockers, alpha-blockers, and fringe vasodilators are the essential medications utilized as a part of treatment. These prescriptions may be utilized alone or as a part of blend, and some are just utilized as a part of mix [83]. Likewise, some of these medications are wanted to others contingent upon the attributes of the patient (diabetic, pregnant, and so on.).

Calcium-channel blockers and disease hazard - postmenopausal females [84] who took calcium-channel blockers for a long time were discovered to be 2.5 times more inclined to create bosom malignancy contrasted with ladies who never took them or those on other hypertension solutions.

In the event that circulatory strain is effectively brought down, it is savvy to have continuous checkups and to take preventive measures to maintain a strategic distance from a backslide of hypertension [85].

Beetroot juice - an examination group from Queen Mary, University of London, wrote in the diary Hypertension that some beetroot squeeze every day can lessen pulse in hypertensive patients [86].

The analysts began off analyzing what the effect of expending nitrates may be on research facility rats, and after that affirmed their discoveries with 15 volunteer people, all with hypertension [87].

#### **The accompanying sustenances are high in nitrates:**

- Beetroot
- Fennel
- Cabbage
- Lettuce
- Radishes
- Carrots. [88-91]

Yoga - Dr. Debbie Cohen and associates from the University of Pennsylvania reported at the "28th Annual Scientific Meeting" that yoga is powerful in lessening pulse [92].

Telemonitoring enhances uncontrolled hypertension - analysts reported critical changes in the wellbeing of hypertensive patients who utilized telemonitoring, which can be utilized at home. Patients utilize a convenient framework permitting them to record and send their pulse readings straight to the specialist's office progressively.

"Exchanging off" hypertension in the body - researchers from University of California San Diego have planned particles that could in the long run be utilized as a part of solutions that "switch off" [93] hypertension in the human body.

#### **Anticipating hypertension**

Hypertension can best be anticipated by changing your way of life so that legitimate eating regimen and activity are key segments. It is essential to keep up a solid weight, lessen salt admission, diminish liquor admission, and decrease stress [94].

#### ***Diet***

Eliminate the measure of salt in your sustenance and eat a lot of products of the soil. The eatwell plate highlights the diverse sorts of sustenance that make up our eating routine, and demonstrates the extents we ought to eat them into have a very much adjusted and sound eating regimen [95].

Salt raises your pulse. The more salt you eat, the higher your pulse. Intend to eat under 6g (0.2oz) of salt a day, which is around a teaspoonful. Figure out all the more about how to eliminate salt.

Eating a low-fat eating routine that incorporates bunches of fiber (for instance, wholegrain rice, bread and pasta) and a lot of products of the soil helps lower circulatory strain [96,97]. Leafy foods are brimming with vitamins, minerals and fiber that keep your body in great condition.

#### ***Liquor***

Frequently drinking liquor above levels will raise your pulse after some time. Staying inside of the prescribed levels is the most ideal approach to lessen your danger of growing hypertension [98].

- men ought not frequently drink more than three to four units a day
- women ought not frequently drink more than a few units a day

Liquor is likewise high in calories, which will make you put on weight. This will likewise expand your circulatory strain [99,100].

### **Weight**

Being overweight powers your heart to work harder to pump blood around your body, which can raise your circulatory strain [101]. See whether you have to get thinner with the BMI solid weight number cruncher.

On the off chance that you do need to shed some weight, it merits recalling that simply losing a couple pounds will have a major effect to your pulse and general wellbeing [102-104].

### **Exercise**

Being dynamic and taking customary activity brings down pulse by keeping your heart and veins in great condition. Normal activity can likewise help you get thinner, which will likewise help bring down your pulse [105].

Grown-ups ought to do no less than 150 minutes (2 hours and 30 minutes) of moderate-force oxygen consuming movement, (for example, cycling or quick strolling) consistently. For it to check, the action ought to make you understand warm and somewhat of breath [106,107]. Somebody who is overweight may just need to stroll up a slant to get this inclination.

Physical action can incorporate anything from game to strolling and cultivating [108].

### **Smoking**

Smoking doesn't specifically cause hypertension however it puts you at much higher danger of a heart assault and stroke [109]. Smoking, similar to hypertension, will bring about your supply routes to thin. On the off chance that you smoke and have hypertension, your veins will limit a great deal all the more rapidly and your danger of heart or lung ailment later on is significantly expanded [110].

### **Caffeine**

Drinking more than some espresso a day may expand your pulse. In the event that you are a major enthusiast of espresso, tea or other caffeine-rich beverages, (for example, cola [111] and some caffeinated beverages [112]), consider chopping down.

It is fine to drink tea and espresso as a feature of an adjusted eating routine yet it is critical that these beverages are not your just wellspring of liquid [113].

### **Decrease of anxiety**

Anxiety or uneasiness can briefly build circulatory strain. Take sooner or later to consider what makes you feel pushed, for example, work, family, funds or sickness [114]. When you comprehend what's bringing about your anxiety [115].

Different supplements might likewise help avert hypertension [116]. those are potassium, calcium, magnesium, fish oils, garlic [117-119].

With a specific end goal to anticipate harm to discriminating organs and conditions, for example, stroke, heart assault, and kidney disappointment that may be brought about by hypertension, it is vital to screen, analyze, treat, and control hyper pressure in its most punctual stages [120]. This can also be accomplished by increasing public awareness and increasing the frequency of screenings for the condition [121].

### **References**

1. Srimatkandada P, et al. Severe Autoimmune Hyperthyroidism in Two Patients with Pulmonary Arterial Hypertension after Treatment with Epoprostenol. *Thyroid Disorders Ther.* 2015;4:176.
2. Divekar A, et al. Non-Restrictive Ductal Patency in Management of Cardiac Failure in Congenital Diaphragmatic Hernia – Non-Invasive Biventricular Assist. *J Neonatal Biol.* 2015;4:171.

3. Wasim M. Role of Leptin in Obesity. *J Obes Weight Loss Ther.* 2015;5:258.
4. Marston N, et al. Update in Echocardiography:Chronic Thromboembolic Pulmonary Hypertension. *J Pulm Respir Med.* 2015;5:252.
5. Bhutani K, et al. Predictive and Prognostic Value of Thyroid Profile and Lipid Profile in Pregnancy Induced Hypertension. *J Preg Child Health.* 2015;2:144.
6. Kincaid BB, et al. Manifestations of Anxiety? Explaining Tachycardia and Hypertension in a Patient with POTS . *Fam Med Med Sci Res.* 2015;4:153.
7. Yamamoto T, et al. Relationship between Digit Ratio and Idiopathic Pulmonary Arterial Hypertension in Japanese Women. *J Vasc Med Surg.* 2015;3:175.
8. Aziz A, et al. Proteomic Profiling of Early Chronic Pulmonary Hypertension: Evidence for Both Adaptive and Maladaptive Pathology. *J Pulm Respir Med.* 2015;5:241.
9. Reyes LR, et al. Clinical Characteristics and Survival of Patients with Pulmonary Hypertension: A 40-Month Mean Follow-Up . *J Pulm Respir Med.* 2015;5:235.
10. Speziale M and Barbini N. Cardiovascular Risk for Workers in Big, Infrastructure Projects. *Occup Med Health Aff.* 2015;3:191.
11. Maatoug J, et al. Three-Year Intervention Program to Prevent Hypertension in Workplaces in Tunisia: A Pre-Post Quasi-Experimental Design with Control Group. *J Clin Trials.* 2015;5:202.
12. Nusee Z, et al. The Effects of Maternal Serum Lipid on Maternal Blood Pressure and Fetal Birth Weight: A Prospective Cohort Study. *Gynecol Obstet (Sunnyvale).* 2015;5:276.
13. Azhar R and Gauhar A Haemodynamic Response to Tracheal Extubation: Verapamil versus Metoprolol. *General Med.* 2015;3:170.
14. Helvaci RM and Ozcan A White Coat Hypertension is a Pioneer Sign of Metabolic Syndrome. *J Metabolic Syndr.* 2015;4:172.
15. Aleyeidi N and Aseri K. The Efficacy of Wet Cupping on Blood Pressure among Hypertension Patients in Jeddah, Saudi Arabia: A Randomized Controlled Trial Pilot Study. *Altern Integ Med* 2015;4:183.
16. Kaul N, et al. Status of T2D In Punjabi Indian Females Based on Risk Variable Factor Analysis. *J Biomol Res Ther.* 2015;4: 122.
17. Kyle D Maxwell, et al. The Plasmalemmal Na/K-ATPase: An Amplifier for Reactive Oxygen Species?. *J Hypertens.* 2015;4:197.
18. Adamu Umar G, et al. Prevalence of Metabolic Syndrome According to Three Defining Criteria in Hypertensive Population in a Rural Hospital Setting. *J Hypertens.* 2015;4:196.
19. Deveci OS, et al. A Novel BioTarget in Treatment of Heart Failure:Changes in Serum Galectin-3 Levels after Spironolactone Therapy. *J Hypertens.* 2015;4:195.
20. Okeahialam BN. Twelve Month Follow-Up Audit of Nigerian Hypertensives on Back Titration. *J Hypertens.* 2015;4:193.
21. Abbas Al Ubaidi B. Putting Evidence Based JNC 8 Guideline into Primary Care Practice. *J Hypertens.* 2015;4:193.
22. Ukimura A, et al. Azilsartan is More Effective as Compared to Olmesartan in Hemodialysis Patients with Uncontrolled Hypertension. *J Nephrol Ther.* 2015;5:193.
23. Asarkua H, et al. Serious Influence of Yersinia Enterocolitis on Pregnancy in a Woman Complicated With Chronic Hypertension and Gestational Diabetes Mellitus: A Case Report. *J Preg Child Health.* 2015;2:135.
24. Kalyani HB, et al. The Effect of *Murraya koenigii* Extract on Therapeutic Efficacy of Amlodipine in Rats: Possible Drug-Herb Interaction. *General Med.* 2015;3:153.
25. Gunes ARIK and Burcu Balam YAVUZ. Hypertension in Older Adults-Geriatrician Point of View. *J Gerontol Geriat Res.* 2014;3:182.
26. Zhu Q, et al. Single Nucleotide Polymorphism of the CYP2J2 Gene is Associated with Essential Hypertension in Uygur Population in China. *Biochem Anal Biochem.* 2015;4:159.

27. Mesiano P, et al. Vascular Injury by Arterial Hypertension in Anti-Phospholipid Antibody Syndrome . J Vasc Med Surg. 2015;3:170.
28. Zafar F, et al. Drug Utilization Pattern in Cardiovascular Diseases: A Descriptive Study in Tertiary Care Settings in Pakistan. J Bioequiv Availab. 2015;7:059-062.
29. Adeyeye OO, et al. Correlates of Abnormal Pulmonary Function Tests in Persons with Type 2 Diabetes Mellitus. J Pulm Respir Med. 2015;5:231.
30. Aliyu SU, et al. Prevalence of Overweight/obesity and Undiagnosed Hypertension among Military Personnel in Maiduguri, Nigeria. J Nov Physiother. 2014;4:237.
31. Giquel J, et al. Cardiovascular Manifestations of Hyperglycemia: A Review Article. J Gen Practice. 2014;2:191.
32. Elnaggar NK . Diabetes and Hypertension. J Hypertens. 2014;3:192.
33. Vaclavik J and Sliva J. ACE Inhibitors or Sartans in the Treatment of Hypertension: A Needless Discussion?. J Hypertens. 2014;3:191.
34. García-Familiar A, et al. Transcatheter Embolisation with Onyx of a Persistent Type 1a Endoleak. J Hypertens. 2014;3:190.
35. Aronow WS. What Should the Systolic Blood Pressure Goal be in Adults with Hypertension?. J Hypertens. 2014;3:189.
36. AtienoJalang'o G, et al. Do Healthcare Workers Adhere to Diabetes Clinical Care Guidelines? A Study at a National Hospital, Kenya. J Hypertens. 2014;3:188.
37. Wang Q, et al. Renin-Dependent Hypertension in Mice Requires the NLRP3-Inflammasome. J Hypertens. 2014;3:187.
38. Adeoye AM, et al. Prevalence of Hypertension and Association with Increased Body Mass in a Semi-Urban Settlement in Nigeria. J Hypertens. 2014;3:186.
39. Montrezol FT, et al. Resistance Training Promotes Reduction in Blood Pressure and Increase Plasma Adiponectin of Hypertensive Elderly Patients. J Hypertens. 2014;3:185.
40. Dallatu MK, et al. The Role of Hypoxia-Inducible Factor/Prolyl Hydroxylation Pathway in Deoxycorticosterone Acetate/Salt Hypertension in the Rat. J Hypertens. 2014;3:184.
41. Miyata S and Noda A. Effects of a Custom-Made Pillow on Sleep and Blood Pressure Circadian Rhythm. J Hypertens. 2014;3:183.
42. Lv Y, et al. Grading of Peripheral Cytopenias due to Splenomegaly and Hepatitis B Cirrhotic Portal Hypertension. J Hypertens. 2014;3:182.
43. Zapata-Sudo G, et al. Targeting of the Adenosine Receptors as A Novel Strategy for the Treatment of Arterial Hypertension . J Neurol Neurophysiol. 2014;5:243.
44. Delacroix S, et al. Hypertension: Pathophysiology and Treatment. J Neurol Neurophysiol. 2015;5:250.
45. Patel JN, et al. Congenital Hepatic Fibrosis Associated with Polycystic Kidney Disease. J Liver. 2015;4:171.
46. Babisch W, et al. Road Traffic, Location of Rooms and Hypertension. J Civil Environ Eng. 2014; 4:162.
47. Dan X, et al. Abdominal Tuberculosis Leading to Prehepatic Portal Hypertension: A Case Report. J Cytol Histol. 2014;5:278.
48. Ikeo S, et al. A Surgical Lung Biopsy Case of Pulmonary Venous-occlusive Disease Associated with Limited Cutaneous Systemic Sclerosis. Rheumatology (Sunnyvale). 2014;S16:005.
49. Batsh MMEI and Batch MMEI. Telmisartan is More Effective than Taurine in Protecting High Fat Diet Induced Obesity in Rats from Hypertension, Some Metabolic, Oxidative Stress and Vascular Complications. Cardiol Pharmacol. 2014;3:119.
50. Bitner A, et al. The Role of Multidrug Interactions in the Safety of Pharmacotherapy for Concomitant Parkinson's Disease and Arterial Hypertension in Poland. J Pharmacovigilance 2014;2: 151.
51. Aliyu SU, et al. Prevalence of Overweight/Obesity and Undiagnosed Hypertension among Military Personnel in Maiduguri, Nigeria. J Nov Physiother. 2014;4:223.

52. Wang M, et al. Case Detection Testing for Primary Aldosteronism in Male Patients with Hypertension and Snoring. *J Hypertens* 2014;3:180.
53. Laneelle D, et al. Home Blood Pressure Self-Monitoring in Patients Treated With Anti-Angiogenic Drugs for the Detection of Arterial Hypertension. *J Hypertens*. 2014;3:179.
54. Lagi A and Cencetti S. Hypertensive Emergencies: a New Clinical Approach. *J Hypertens*. 2014;3:178.
55. Bcheraoui TCEI, et al. Reported Stroke Symptoms and their Associated Risk Factors in the Kingdom of Saudi Arabia, 2013. *J Hypertens*. 2014;3:177.
56. DeJongste MJL, et al. Can Too Few and Too Many Climato-Economic Resources Elevate Blood Pressure? A 120-Nation Study. *J Hypertens*. 2014;3:176.
57. Girma F, et al. Compliance with Anti-Hypertensive Treatment and Associated Factors among Hypertensive Patients on Follow-Up in Jimma University Specialized Hospital, Jimma, South West Ethiopia: A Quantitative Cross- sectional Study. *J Hypertens*. 2014;3:174.
58. Zuo HJ, et al. Relationship between Four Blood Pressure Indexes and Ischemic Stroke in Patients with Uncontrolled Hypertension. *J Hypertens*. 2014;3:173.
59. Lv YF, et al. Prognostic Risk Factors for Surgery in Patients with Cirrhotic Portal Hypertension. *J Hypertens*. 2014;3:172.
60. Dhawan V, et al. Implication of Endothelin-2 and Oxidative Stress Biomarkers in Essential Hypertension. *J Hypertens*. 2014;3:170.
61. Kuhn-Thiel AM and Wehling M. Antihypertensive Drugs in the Elderly: Risks and Benefits as Evaluated by the FORTA Method. *J Gerontol Geriatr Res*. 2014;3:1000171.
62. Seimetz M, et al. Pulmonary Hypertension Precedes Emphysema: Paradigm Shift or Artifact of Rodent Studies?. *Angiol Open Access*. 2014;2:e108.
63. Hussein M, et al. Assessment of Drug Related Problems Among Hypertensive Patients on Follow up in Adama Hospital Medical College, East Ethiopia. *Clin Pharmacol Biopharm*. 2014;3:122.
64. Moayedi S. Acute Management of Cardiac Complications in Pregnancy. *Gynecol Obstet (Sunnyvale)*. 2014;4:245.
65. Blicharska et al. Examination of the Elemental Composition of Hair in Cholelithiasis, Kidney Stone, Hypertension and Diabetes by Scanning Electron Microscopy and Energy Dispersive Spectrometry SEM/EDS. *J Anal Bioanal Tech*. 2014;5:207.
66. Shinde S. Treprostinil: Safety Signal Detection Based on Adverse Event Reporting System Database. *J Pharmacovigilance*. 2014;2:140.
67. Speziale M and Barbini N. Cardiovascular Risk for Workers in Big, Infrastructure Projects. *Occup Med Health Aff*. 2014;2:175.
68. Houston MC. The Role of Mercury in Cardiovascular Disease. *J Cardiovasc Dis Diagn*. 2014;2: 170.
69. Rao GHR, et al. Clinical Complications of Type- 2 Diabetes Mellitus in South Asian and Chinese Populations: An Overview. *J Diabetes Metab*. 2014;5:420.
70. Dharwadkar MN, et al. Study of Methyl Dopa Versus Labetalol in Management of Preeclampsia and Gestational Hypertension. *Gynecol Obstet (Sunnyvale)*. 2014;4:242.
71. Ahmad S, et al. Malignant Pheochromocytoma: A Case Report. *Emerg Med (Los Angel)*. 2014;4:209.
72. García AR, et al. Recognition and Initial Treatment of Suspected Intracranial Hypertension by Primary Care Pediatricians: An Advanced Simulation Observational Study. *Emerg Med*. 2014;4:205.
73. Ishikawa Y. Obstructive Sleep Apnea and Hypertension; A New Way to Go. *J Sleep Disorders Ther*. 2014;3:e128.
74. Sang LY, et al. A Self-assembling Nanomaterial Reduces Acute Brain Injury and Enhances Functional Recovery in a Rat Model of Hypertensive Intracerebral Hemorrhage. *J Nanomed Nanotechnol*. 2014;5:224.
75. Jennings CG, et al. Review of Hyperuricaemia and Hypertension: A Target for Treatment. *J Hypertens*. 2014;3:164.

76. Karter Y. Oxidative Stress and Endothelial Dysfunction in White Coat Hypertension: Data of the Group from Cerrahpasa Medical Faculty. *J Hypertens.* 2014;3:163.
77. Liu Z, et al. The Association Study of PRKG1 Gene Polymorphism and Salt-Sensitive Hypertension among the Essential Hypertension in Beijing. *J Hypertens.* 2014;3:162.
78. Ding X, et al. Mitochondrial Long Chain Fatty Acid Oxidation Related Enzyme Changes in Different Preeclampsia- Like Mouse Models. *J Hypertens.* 2014;3:161.
79. Cao T, et al. Brain-Targeted (Pro) Renin Receptor Knockdown Modulates Body Fluid Homeostasis. *J Hypertens.* 2014;3:160.
80. Wei G, et al. Assay Profiles of Prostanoids and Female Sex Hormones for Studying Potential Biomarkers of Hypertension. *J Hypertens.* 2014;3:159.
81. Parlak A, et al. Investigation of Pentraxin 3 Levels in Hypertensive Patients with Stroke, Retinopathy and Nephropathy. *J Hypertens.* 2014;3:158.
82. Skultetyova D, et al. The Impact of Blood Pressure on Carotid Artery Stiffness and Wave Intensity in Patients with Resistant Hypertension after Renal Sympathetic Denervation. *J Hypertens.* 2014;3:157.
83. Shukla AN, et al. The Prevalence of Hypertension: Role of Hereditary in Young and Obesity in all. *J Hypertens.* 2014;3:156.
84. Tibaldi M, et al. Prevalence and Clinical Implications of Orthostatic Hypotension in Elderly Inpatients. *J Hypertens.* 2014;3:155.
85. Hoshida S, et al. Analysis of the Difference in the Pressure-Natriuresis Relationship Based on the Class of Medication Using Spot Urine Tests in Hypertensive Patients. *J Hypertens.* 2014;3:154.
86. Colvin KL and Yeager ME. Animal Models of Pulmonary Hypertension: Matching Disease Mechanisms to Etiology of the Human Disease. *J Pulm Respir Med.* 2014;4:198.
87. Jasper US. Magnitude of Obesity, Abdominal Adiposity and their Association with Hypertension and Diabetes- A Cross Sectional Study. *J Metabolic Synd.* 2014;3:146.
88. Chakore and Eklare. Carcinoma Lung Presenting as Intracranial Hypertension and Lower Cranial Nerve Palsies. *J Neurol Disord.* 2014 2:163.
89. Colvin L, et al. The Use of Angiotensin Converting Enzyme Inhibitors during the First Trimester of Pregnancy. *J Pharmacovigilance.* 2014;2:129.
90. Gemalasar Liman NG, et al. Education-Oriented Control Book for Patients with Hypertension in Primary Health Care. *J Community Med Health Educ.* 2014;4:296.
91. Genel S, et al. All Things are Lawful, but not all Things are Helpful. Reflections on the Risk Factors in Cardiovascular Disease . *J Diabetes Metab.* 2014;5:e114.
92. Chik Z, et al. A Bioequivalence Comparison of Two Captopril Formulations (25 mg Tablets): An Open-Label, Randomized, Two-Treatment, Two-Way Crossover Study in Healthy Volunteers. *J Bioequiv Availab.* 2014;6:080-085.
93. Ihle F, et al. An Integrated Outpatient Training Program for Patients with Pulmonary Hypertension - the Munich Pilot Project. *Int J Phys Med Rehabil.* 2014;2:204.
94. Emelianova YP, et al. Oscillator Suppression in the Blood Flow Regulation of Interacting, Non-Identical Nephrons. *J Hypertens.* 2014;3:153.
95. Denault AY, et al. Pilot Randomized Controlled Trial of Inhaled Milrinone in High-Risk Cardiac Surgical Patients. *Surgery Curr Res.* 2014;4: 192.
96. Brinth L, et al. Baroreflex Sensitivity in Relation to Clinical Characteristics in Subject Aged 40 to 80 Years. *J Hypertens.* 2014;3:152.
97. Zhu H, et al. Relationship between Nocturnal Blood Pressure and Left Ventricular Hypertrophy in Hypertensive Patients. *J Hypertens.* 2014;3:151.
98. Ortiz RM, et al. Increased Physical Activity Reduces the Odds of Elevated Systolic Blood Pressure Independent of Body Mass or Ethnicity in Rural Adolescents. *J Hypertens.* 2014;3:150.

99. Ando T, et al. Sudden Onset of Pyrexia and Severe Blood Pressure Elevation Due to IL-6 Production by Pheochromocytoma. *J Hypertens.* 2014;3:149.
100. Juncos LI and Joaquin C. Dual Blocking of the Renin Angiotensin System: A Settled Issue? *J Hypertens.* 2014;3:148.
101. Williams EA, et al. Silent Crisis: Epidemic Hypertension in Rural West Africa. *J Hypertens.* 2014;3:147.
102. Araújo AV, et al. Function and Protein Expression of Potassium Channels in Mesenteric Resistance Arteries Isolated from 2K-1C Hypertensive Rats. *J Hypertens.* 2014;3:146.
103. Jurado SR. Effects of the Maternal Hypertension in Renal Development in Offspring of Rats. *J Clin Exp Cardiol.* 2014;5: 306.
104. Andreas Fette. Sildenafil in Use for Treatment of Pulmonary Hypertension in Congenital Diaphragmatic Hernia: Short Literature Review. *Pediat Therapeut.* 2014;4:197.
105. Ciccone MM, et al. Endothelial Function in Pre-diabetes, Diabetes and Diabetic Cardiomyopathy: A Review. *J Diabetes Metab.* 2014;5:364.
106. Hao XQ, et al. Telmisartan Protect Broiler Chickens from Pulmonary Arterial Remodeling Induced by Low Ambient Temperature. *J Clin Exp Cardiol.* 2014;5: 300.
107. Soejima H, et al. The Changes of Biomarkers by Telmisartan and their Significance in Cardiovascular Outcomes: Design of a Trial of Telmisartan Prevention of Cardiovascular Diseases (ATTEMPT-CVD). *J Clin Trials.* 2014;4:162
108. Joison AN, et al. Study the Small and Dense Lipoprotein Levels (LDLsd) in Coronary Artery Disease. *J Cardiovasc Dis Diagn.* 2014;2:151.
109. Lemaire A, et al. (2014) Refractory Hypertension Leads to both Type A and B Aortic Dissections in a Single Patient. *J Cardiovasc Dis Diagn.* 2014;2:155.
110. Aziz R and Mahboob T. Serum Calcium, Magnesium and Parathyroid Hormone in Normal Pregnant and Pre-eclamptic Women in Karachi, Pakistan. *J Hypertens.* 2014;3:143.
111. Hänninen MR. Is it Practical to Search for Masked Hypertension? *J Hypertens.* 2014;3:144.
112. Yang Q and Sun M. Role of MicroRNAs in Hypoxia-Induced Pulmonary Hypertension. *Cardiol Pharmacol.* 2014;3:e124.
113. Zeng R, et al. The Use of an Amlodipine“atorvastatin Single Pill Combination in the Management of Hypertension and Hypercholesterolemia: Is Time an Important Problem? *J Clin Exp Cardiol.* 2014;5:297.
114. Lisa G, et al. Clinical Impact and Prognosis of Pulmonary Hypertension after Transcatheter Aortic Valve Implantation. *J Cardiovasc Dis Diagn.* 2014;2:143.
115. Liu PP, et al. Association between ADD1 Gly460Trp Polymorphism and Essential Hypertension in Han Chinese. *J Hypertens.* 2014;3:145.
116. Berezin AE and Lisovaya OA. Predictive Value of Circulating Vascular Endothelial Growth Factor-1 Level Measured Repeatedly During Long-Term Follow-Up in Patients with Arterial Hypertension after Acute Ischemic Stroke. *Angiol.* 2014;2:119.
117. Ando D, et al. Hypoalbuminemia and Hyperalbuminuria as Major Factors Regulating Circadian Blood Pressure Rhythm in Normal Subjects and Non-diabetic Patients with Early Stages of Chronic Kidney Disease. *J Hypertens.* 2014;3:142.
118. Imbalzano E, et al. Renal Denervation, “Last Resort” or Alternative Therapy? *J Hypertens.* 2014;3:141.
119. Dézsi CA, et al. New Procedure in a New Situation: Short-term Effect of Renal Denervation for Resistant Hypertension in Case of a Patient Living with Pacemaker. *J Hypertens.* 2014;3:140.
120. Lovica D and Lovica B. Insights into Pathophysiology of Carotid Baroreceptor Stimulation as a Method for Treatment of Resistant Hypertension. *J Hypertens.* 2014;3:139.

121. Fukuda N, et al. Olmesartan Improves the Formation of Impaired EPCs and Renal Degeneration through Activation of the ACE2/Ang-(1<sup>7</sup>)/Mas Receptor Axis in SHR. *J Hypertens*. 2014;3:138.