

Representation of a Full Transformation Semi-group Over a Finite Field

Munir Ahmed^{1*}, Muhammad Naseer Khan², Muhammad Afzal Rana^{2*}

¹Department of Mathematics, Islamabad Model College For Boys, Islamabad, Pakistan

²Department Of Mathematics and Statistics, Riphah International University, Islamabad, Pakistan

Research Article

Received date: 09/01/2018
Accepted date: 10/05/2018
Published date: 11/06/2018

*For Correspondence

Department Of Mathematics, Islamabad Model College For Boys, F-10/4, Islamabad, Pakistan

E-mail: irmunir@yahoo.com

Keywords: Semigroup of transformations;
Representation of semigroup; Finite field

ABSTRACT

In this paper we discuss the representations of a full transformation semigroup over a finite field. Furthermore, we observe some properties of irreducibility representation of a full transformation semigroup and discuss the linear representation of a zero-adjointed full transformation semigroup. Moreover, we characterize the linear representation of a full transformation semigroup over a finite field F_q (where q is a prime power) in terms of Maschke's Theorem. Finally, we observe that there exists an isomorphism between the full matrix algebra $(F_q)_m$ and the space of all linear transformation $L(F_q^m)$ on an m -dimensional vector space F_q^m .

INTRODUCTION

Serre has given a comprehensive theory of linear representation of finite groups in [1]. It has been obtained in the group theory that the number of simple FG- modules is equal to the number of conjugacy classes of the group G such that the characteristic of the field F does not divide the order of G . A lot of work is done for the classification of groups in terms of its representation and characterization.

By Clifford, each element of a semigroup is uniquely determined by a matrix over a field and a complete classification of the representations of a particular class of a semigroups is given in [2-4]. Moreover, irreducible representations of a semigroup over a field is obtained as the basic extensions to the semigroup of the extendible irreducible representations of a group, and the representations of completely simple semigroup is also constructed in [2-4].

Stoll has given a characterization of a transitive representation, and obtained a transitive representation of a finite simple semigroup, see [5]. The construction of all representations of a type of finite semigroup which is sum of a set of isomorphic groups is also obtained. Munn obtained a complete set of inequivalent representations of a semigroup S which are irreducible in terms of those of its basic groups of its principal factors. He also introduced the principal representations of a semigroup in [6]. A representation of semigroup whose algebra is semisimple is characterized in [7,8]. The representation of a finite semigroup for which the corresponding semigroup algebra is semisimple is also obtained. An explicit determination of all the irreducible representations of T^n is due to Hewit and Zuckerman in [9].

There is a one-to-one correspondence between the representations of a group G and the nonsingular representations of the semigroup S , which preserves equivalence, reduction and decomposition [10].

In the case of an irreducible representation of a finite semigroup, the factorization can be avoided and an explicit expression of such representation is given in [11]. We consider a full transformation semigroup T^n to obtain its combinatorial property with regard to its irreducible representations. There exists a non-zero linear transformation satisfying some specific conditions in Theorem 7.3.

It is observed that for the basis \mathcal{B} of a vector space F_q^m , there is a natural one-to-one correspondence (between the rep-

representations of a full transformation semigroup \mathcal{T}_q over a finite field F_q and those of the algebra $F_q[\mathcal{T}_q]$ which preserves, equivalence, reduction and decomposition into irreducible constituents.

Consequently, we reinterpret the Maschke Theorem [12] regarding the algebra $F_q[\mathcal{T}_q]$, i.e., the algebra $F_q[\mathcal{T}_q]$ is semi-simple if and only if the characteristic of F_q does not divide the order mm of the full transformation semigroup \mathcal{T}_q .

The representation of full transformation semigroup over a finite field is discussed in Section-8, specially the Maschke's theorem is restated for the semisimplicity of the semigroup algebra $F_q[\mathcal{T}_q]$, see Theorem 8.1 Finally, a linear algebraic result regarding the isomorphism between the full matrix algebra $(F_q)_m$ and the space of all the linear transformations on F_q^m is given in Theorem 8.2.

PRELIMINARIES

Definition

A transformation semigroup is a collection of maps of a set into itself which is closed under the operation of composition of functions. If it includes identity mapping, then it is a monoid. It is called a transformation monoid.

If (X,S) is a transformation semigroup then X can be made into semigroup action of S by evaluation, $x.s=xs=y$ for $s \in S$, and $x,y \in X$. This is the monoid action of S on X , if S is a transformation monoid.

Hewitt and Zuckerman gives a treatment of the irreducible representation of the transformation semigroup on a set of finite cardinality [8]. The result for the case of a finite semigroup S with $F[S]$ semisimple was given by Munn in [13].

The full reducibility and the proper extensions of irreducible representations of a group to those of a semigroup are the basic extensions.

THEOREM 2.2

Full reducibility holds for the representations of a semigroup S over the field F if and only if

Full reducibility holds for the extendible representations of G over F , and

The only proper extension of a proper representation of G to S is the basic extension [14].

A representation M of S is homomorphism of S into the multiplicative semigroup of all (α, α) matrices (where α is an arbitrary positive integer) such that $M(x) \neq 0$ for some $x \in S$. If the set $\{M(x) : x \in S\}$ is irreducible i.e., if every (α, α) matrix is a linear combination of matrices $M(x)$, then M is said to be an irreducible representation of S . The identity representation is the mapping that carries every $x \in S$ into the identity matrix.

Full transformation semigroup

The idea of studying T_n was suggested by Miller (in oral communication). The problem of obtaining representations of semigroup as distinct from groups have been first studied by Suskevic. Clifford has given a construction of all representations of a class of semigroups closely connected with T_n . Popizovski has pointed out some simple properties of T_n . In the present discussion, we relate the irreducible representations of T_n to that of its semigroup algebra $L(T_n)$. The set of all transformations of set X into itself is called the full transformation semigroup under the binary operation of multiplication as the composition of transformation analogue of the symmetric group G_x . Let $X_n = \{1,2,3,\dots,n\}$ be a finite set and denote the semigroup TX_n of all the self-maps of X_n into X_n . If cardinality of X_n is n , denote T_n for TX_n then the cardinality of T_n is n^n [15].

Example

The set $S=\{e,a,x,y\}$ is a semigroup under the multiplication. The Cayley's multiplication table of S is given as follows [16].

.	e	a	x	y
e	e	a	x	y
a	a	e	x	y
x	x	y	x	y
y	y	x	x	y

If the mapping $\phi : S \rightarrow \mathcal{S}_X = \{1,2\}$ is given by $x\phi = \beta$, $a\phi = \beta$, $e\phi = \beta$, and $y\phi = \gamma$, then ϕ embeds S in $\mathcal{T}_{\{1,2\}}$. It can also be seen that the map $\psi : S \rightarrow \mathcal{T}_{\{a,e,x,y\}}$ is defined by

$$\psi(e) = \begin{pmatrix} e & a & x & y \\ e & a & x & y \end{pmatrix},$$

$$\psi(x) = \begin{pmatrix} e & a & x & y \\ x & x & x & x \end{pmatrix},$$

$$\psi(y) = \begin{pmatrix} e & a & x & y \\ x & x & x & x \end{pmatrix},$$

and

$$\psi(y) = \begin{pmatrix} e & a & x & y \\ y & y & y & y \end{pmatrix}.$$

embeds S into $\mathcal{T}_{\{a,e,x,y\}}$.

Notice that ψ is a right regular representation of S, where $\psi : S \rightarrow \mathcal{T}_S$ as defined above (where $\psi(e), \psi(a), \psi(x), \psi(y) \in \mathcal{T}_S$) is such that for any $s \in S$, we have

$$(\psi e)(s) = se$$

$$(\psi a)(s) = sa$$

$$(\psi x)(s) = sx$$

$$(\psi y)(s) = sy$$

So ψ is a right regular representation of S.

Regular representation of a transformation semigroup

Let K denote the set of right zero elements of a semigroup S. Then, $s \in K$ if and only if

(i) for all x in K, and all a,b in S, xa=xb implies a=b;

(ii) if α is any transformation of K, then there exists a in S such that $x\alpha = xa$ for all $x \in K$.

An element α of \mathcal{T}_X is idempotent if and only if it is the identity mapping when restricted to $X\alpha$. Suppose that X is a set of cardinality n. Then, the full transformation semigroup \mathcal{T}_X contains the symmetric group G_X of degree n. If $\alpha \in \mathcal{T}_X$, then the rank r of α is defined by $r = |X\alpha|$, and the defect of the element α is given by n-r. If β is an element of \mathcal{T}_X of rank $r < n$, then there exists elements γ and δ of \mathcal{T}_X such that β has the rank $r+1$, δ has the rank $n-1$, and $\beta = \gamma\delta$ (we can choose δ as an idempotent, and γ different from β at only one part of X). By induction, every element of \mathcal{T}_X of defect $k (1 \leq k \leq n-1)$ can be expressed as the product of an element of G_X and k number of (idempotent) elements of defect 1, see also [17].

If $\alpha \in \mathcal{T}_X$ is of defect 1, then every other element of \mathcal{T}_X of defect 1 can be expressed in the form $\lambda\alpha\mu$ with λ and μ are in G_X . If α is an element of \mathcal{T}_S of defect 1, then $\langle G_X\alpha \rangle = \mathcal{T}_S$.

Let $X=S$ be a semigroup, an element $\rho \in \mathcal{T}_S$ is said to be a right translation of S if $x(\rho y) = (xy)\rho$ for all $x,y \in S$ and $\lambda \in \mathcal{T}_X$ is said to be a left translation of S if $(x\lambda)y = (xy)\lambda$ for any $x,y \in S$. The left and a right translations λ and ρ , respectively, are called linked if $x(\lambda y) = (x\lambda)y$ for all $x,y \in S$.

Note that $\lambda_a\lambda = \lambda_{a\lambda}$ and $\rho_a\rho = \rho_{a\rho}$, if λ and ρ are linked, then

$$\lambda\lambda_a = \lambda_{a\rho}, \quad \rho\rho_a = \rho_{a\lambda}$$

Let $S = \{e,f,g,\alpha\}$ be a semigroup with the operation “.” given by the Cayley's table

.	e	a	x	y
e	e	a	x	y
a	a	e	x	y
x	x	y	x	y
y	y	x	x	y

Cayley's table

The transformation

$$\lambda = \begin{pmatrix} e & f & g & a \\ g & g & e & g \end{pmatrix}$$

is a left translation which is not linked with any right translations of S. We recall the following proposition regarding the semisimple algebra.

PROPOSITION

An algebra A is a semisimple if and only if A-module of A is semisimple.

Definition

Let S be a semisimple with zero element z. The contracted algebra $F_0[S]$ of S over F is an algebra over F containing a basis \mathfrak{S} such that $\mathfrak{S} \cup \{0\}$ is a subsemigroup of $F_0[S]$ isomorphic with S. A semisimple algebra can also be regarded as a contracted semigroup algebra.

We recall the following facts regarding the representations of a semisimple algebra.

Lemma

(a) Let \mathfrak{R} be an algebra having finite order over the field F, and let \mathfrak{R} be a radical of \mathfrak{A} . Then, every non-null irreducible representation of \mathfrak{A} maps \mathfrak{R} into 0, and so it is effectively a representation of the semisimple algebra $\mathfrak{A} / \mathfrak{R}$.

(b) Let ϕ be any faithful representation of a semisimple algebra \mathfrak{A} and let P be an $n \times n$ matrix over \mathcal{T} . Then, P is non-singular if and only if $\phi^{(n)}(P)$ is non-singular [18].

THEOREM 4.4

(6, Th. 5.7). An irreducible algebra of linear transformations is simple.

If $A \in (F)_n$, then the transformation $x \rightarrow Ax$ of a vector space V is linear transformation τ of V to V, and the mapping $A \rightarrow A$ is an isomorphism of $(F)_n$ upon the algebra $\mathcal{L}[\mathcal{T}_V]$ of all linear transformations of V. A homomorphism ϕ of \mathfrak{A} into $(F)_n$ is called a representation of \mathfrak{A} of degree n over F. In other words, to each element x of \mathfrak{A} there corresponds an $n \times n$ matrix $\phi(x)$ such that

$$\phi(x+y) = \phi(x) + \phi(y);$$

$$\phi(xy) = \phi(x)\phi(y);$$

$$\phi(\alpha x) = \alpha\phi(x);$$

for all x,y in \mathcal{T}_V and α in F.

The irreducible representations of semigroups

Let f be an element of \mathcal{T}_V . Then, f splits the set $\{1,2,\dots,n\}$ into a number p of nonvoid disjoint subsets, each of the form $\{x: f(x)=a\}$ for some $a \in \text{rang}(f)$. Obviously, f is determined by these sets and the corresponding a's. For nonvoid subset s of $\{1,2,\dots,n\}$, let s^* be the least element of s. Write the sets $\{x: f(x)=a\}$ in the order s_1, s_2, \dots, s_p where $s_1^* < s_2^* < \dots < s_p^*$, and represent f by the symbol

$$\begin{pmatrix} s_1 s_2 \dots s_p \\ a_1 a_2 \dots a_p \end{pmatrix},$$

where $1 \leq s_i \leq n$, the class of sets s_1, \dots, s_p is a decomposition of $\{1,2,\dots,n\}$ of the kind described above, and a_1, a_2, \dots, a_p are any distinct integers lying between 1 and n. The expression s_1, \dots, s_p will always mean a decomposition of $\{1,2,\dots,n\}$ into nonvoid, disjoint subsets with $s_1^* < s_2^* < \dots < s_p^*$. The letters t and w will be used similarly. Also a_1, a_2, \dots, a_p will always mean any ordered sequence of distinct integers from 1 to n; the letters c and d will be used similarly.

For $p = 1, 2, \dots, n$, let \mathfrak{S}_p be the set of all elements of \mathfrak{S}_n whose range contains just p elements, that is,

$$\begin{pmatrix} s_1 s_2 \dots s_p \\ a_1 a_2 \dots a_p \end{pmatrix},$$

for a fixed p. Strictly speaking, \mathfrak{S}_p depends upon n as well as p. However, only one value of n will be treated at one time. The set \mathfrak{S}_n is obviously the symmetric group S_n . The set $\mathfrak{S}_{p,1}$ is a semigroup with the trivial multiplication $fg=f$. No other \mathfrak{S}_p is a subgroup of \mathfrak{S}_n . It will be convenient to have the semigroup $\mathfrak{S}_p \cup \{z\}$, with multiplication defined by

$$z \circ z = f \circ z = z \circ f = z, \text{ for all } f \in \mathfrak{S}_p$$

$$f \circ g = \begin{cases} fg & \text{if } fg \in \mathfrak{S}_p, \\ z & \text{if } fg \notin \mathfrak{S}_p. \end{cases}$$

Using a linear algebraic result, we have the following formula regarding the rank of a linear representation of T_n .

THEOREM 5.1

Let M be an irreducible linear representation of T_n , and let $S = \{f \in T_n \text{ and } M(f) = 0\}$, then

$$\text{rank}[M(T_n)]$$

$$= \begin{cases} n^n, & \text{if } S \text{ is void} \\ n^n - \sum_{j=1}^p j!, & \text{if } S \text{ is nonvoid, i.e., if } S = \cup_{j=1}^p B_j \end{cases} \quad \gg \ll$$

Proof

Suppose the irreducible linear representation $M : T_n \rightarrow L(T_n)$ is as given above. Since M is irreducible representation of T_n . Thus, using a result in, the set S is void or $S = \cup_{j=1}^p B_j$.

Since,

$$\dim F[T_n] = \dim F[S] + \dim F[M(T_n)],$$

where F is a field of characteristic 0.

Since,

$$\dim F[T_n] = n^n,$$

and,

$$|S| = \begin{cases} 0 & \text{if } S \text{ is void,} \\ \sum_{j=1}^p j! & \text{if } S \text{ is nonvoid.} \end{cases}$$

We have

$$\text{rank} F[M(T_n)] = \dim F[M(T_n)].$$

Thus,

$$\text{rank} F[M(T_n)] = \dim F[T_n] - \dim(F[S]) = \begin{cases} n^n - 0 & \text{if } S \text{ is void,} \\ n^n - \sum_{j=1}^p j! & \text{if } S \text{ is nonvoid.} \end{cases}$$

$$\text{rank} F[M(T_n)] = \begin{cases} n^n - 0 & \text{if } S \text{ is void,} \\ n^n - \sum_{j=1}^p j! & \text{if } S \text{ is nonvoid.} \end{cases}$$

Therefore,

This completes the proof.

Let $X = \{x_1, x_2, \dots, x_n\}$ be a set of cardinality n and let S_n denote the set of all single-valued maps of X to itself. We have the following characterization of a map from S_n into the set of all $n \times n$ matrices D_n over the field F , see also.

THEOREM 5.2

Let $M : S_n \rightarrow D_n$ be a map defined by $M(f) = A_f \in D_n$, for $f \in S_n$. Then, M forms a homomorphism of S_n into D_n . If, in particular, S_n is a semigroup S , then M becomes a representation of S into D_n (where z is a zero element).

Proof

For any two single valued maps f and g in S_n , the product fg is also a single valued map, therefore $fg \in S_n$.

Moreover, since $M(f) = A_f \in D_n$ and $M(g) = A_g \in D_n$, therefore $M(fg) = A_{fg} = A_f A_g = M(f) \cdot M(g) \in D_n$. In particular, if i is the identity map

on X , then $M(i)=A_i=I_n \in D_n$, then we have;

$$M(ig)=M(g)=A_g=I_n:A_g=A_i:A_g=M(i)M(g), \text{ and}$$

$$M(f i)=M(f)=A_f=A_f:I_n=A_f:A_i=M(f):M(i).$$

Therefore, M defines a homomorphism of S_n into D_n .

If, in particular, if $S_n=S=\mathcal{T}_N$ the semigroup of all maps from X into itself, then we can define an induced structure on the adjoined zero semigroup $'N$, where z is a zero element, i.e., for any $f \in \mathcal{T}_N$, we have

$$z.z = f.z = z.f = z \quad \forall f \in \mathcal{T}_N.$$

The induced structure on $\mathcal{T}_N \cup \{z\}$ is defined as follows:

$$f \circ g = \begin{cases} fg \in \mathcal{T}_N & \text{if } f \text{ and } g \text{ in } \mathcal{T}_N, \\ z & \text{if one of } f \text{ and } g \text{ is not in } \mathcal{T}_N. \end{cases}$$

Then, the homomorphism M can be extended into a map \overline{M} of the semigroup $\overline{S} = \mathcal{T}_N \cup \{z\}$ into D_n , i.e., $\overline{M} : \overline{S} \rightarrow D_n$ is defined by

$$\overline{M}(z) = M_0 = 0_{n \times n} \in D_n,$$

$$\overline{M}(z) = M(f) \quad \forall f \in S.$$

Therefore,

$$\overline{M}(af) = M_{af} = M(af) = M(af) = aM(f) = a\overline{M}(f) \in D_n,$$

And

$$\overline{M}(f + g) = M(f + g) = A_{(f+g)} = A_f + A_g = M(f) + M(g) = \overline{M}(f) + \overline{M}(g),$$

$$\overline{M}(fg) = M(fg) = A_{fg} = A_f \cdot A_g = M(f)M(g) = \overline{M}(f) \cdot \overline{M}(g).$$

Thus, \overline{M} becomes a representation on S .

Representation of a semigroup of linear transformations in green's

Relations

Two things that can be associated with an element $\alpha \in \Pi_{\alpha-\alpha\alpha}$ are as follows:

1. the range $X\alpha$ of α , and
2. the partition $\Pi_\alpha = \alpha \circ \alpha^{-1}$ of X by $x \Pi_\alpha y (x, y \in X)$ if $x\alpha=y\alpha$ which defines an equivalence relation on X .

Let Π_α^h be the natural mapping of X upon the set X / Π_α of equivalence classes of $X \text{ mod } \Pi_\alpha$. Then, $x \Pi_\alpha^h \rightarrow x\alpha$ becomes a one-to-one mapping of X / Π_α upon $X\alpha$. It follows that $|X \Pi_\alpha| = |X\alpha|$, and this cardinal number is called the rank of α .

Remark

The Ex.2.2.6 in [4] can be rewritten as follows,

Let F be a field and V be a vector space over F . By the dimension $\dim V$ of we mean the cardinal number of a basis of V over F . Let $\mathcal{L}(V)$ be the multiplicative semigroup (i.e., under the operation of composition of maps) of all linear transformations of V with each element t of $L(V)$ we associate two subspaces of V that are given as follows:

1. the range V^τ of τ , consisting of all $(x)^\tau$ with $x \in V$ and,
2. the null space N^τ of τ , consisting of all y in V such that $(y)^\tau = 0$.

(a) Let $\tau \in \mathcal{L}(V)$, and W be a subspace of V , complementary to the null space N^τ , so that $V = \dots$.

Then, τ induces a non-singular matrix A .

Hence, $\dim(V=N^\tau)=\dim(W)=\dim(V)$; is called rank of t . The difference or quotient space of V modulo N^τ is denoted by V/N^τ or by V/N_{τ} . If $\dim V$ is finite, this notation of rank is the usual one as for the matrix A , since VA is the row-space of A . Also N_A is the orthogonal complement of the column-space of A .

- (b) Two elements of the space $\mathcal{L}(\mathcal{T}_V)$ are $\mathcal{L}(\mathcal{R})$ -equivalent if and only if they have the same range (null-space).
- (c) If N and W are subspaces of V such that $\dim(V/N^\tau) = \dim W$, then there exists at least one element ρ of \mathcal{T}_V such that $N = N\rho$ and $W = V\rho$.
- (d) Two elements τ_1 and $\tau_2 \in \mathcal{L}(V)$ are \mathcal{R} -equivalent if and only if $\text{rank}(\tau_1) = \text{rank}(\tau_2)$.
- (e) The Th. 2.9 holds for $\mathcal{L}(V)$ instead of \mathcal{T}_X if we replace “subset Y of X ” by “the subspace W of V ”, \mathcal{T}_V by $\dim W$, “partition \mathcal{T}_V of X ” by “subspace N of V ”, and $|X/\Pi|$ by $\dim(V/N)$.

Linear representation of a full transformation semigroup over a finite field

Definition

Let V be a vector space over the field $F(=C)$ the complex numbers and let the finite subset $\{e_i\}_{i=1}^n$ of V be a basis for V , i.e., $\dim V = n$, let \mathcal{T}_V denote the full transformation semigroup over V . The space $\mathcal{L}(\mathcal{T}_V)$ denotes the space of all linear transformations on V . If a is in $\mathcal{L}(\mathcal{T}_V)$, a linear transformation, then, each $a:V \rightarrow V$ is represented by a square matrix (a_{ij}) of order n . The coefficients a_{ij} are complex numbers for all i and $j=1, \dots, n$ and are obtained by

$$a(e_j) = \sum_{i=1}^n a_{ij} e_i$$

where a can be identified as a morphism which is equivalent to saying that $\det(a) = \det(a_{ij}) \neq 0$. The linear space $\mathcal{L}(\mathcal{T}_S)$ of full transformation semigroup can be identified with the semigroup of all transformations of degree n .

A representation $\phi : S \rightarrow \mathcal{L}(\mathcal{T}_S)$ is faithful if and only if ϕ is one-to-one homomorphism. A representation ϕ of a semigroup S , of degree n over the field F , we mean a homomorphism of S into the semigroup $\mathcal{L}(\mathcal{T}_{F^n})$ of all linear transformation over F^n , where $F^n \cong F[S]$, the vector space is generated by S over the field F . Thus, to each element s of S there corresponds a linear transformation $\phi(s) \in \mathcal{L}(\mathcal{T}_{F^n})$ such that $\phi(st) = \phi(s)\phi(t)$ for all $s, t \in S$.

We denote the algebra of all linear transformations over the n -dimensional vector space F^n over the field F by $F(\mathcal{T}_{F^n})$. Obviously, $F(\mathcal{T}_{F^n})$ appears as a subspace of $\mathcal{L}(\mathcal{T}_{F^n})$.

If ϕ is an isomorphism of S upon a subsemigroup of $F(\mathcal{T}_{F^n})$; then ϕ is said to be faithful. We shall determine all the representations of various classes of finite semigroups over a finite field F_q . If S is a finite semigroup, then there is a one-to-one correspondence between a representation of S and that of algebra $F_q[S]$ over the finite field F_q . Of course, this correspondence preserves the reduction, decomposition and hence the full reducibility hold for such representations of S if and only if $F_q[S]$ is semisimple that holds if q does not divide the $\dim F_q^n = n$, (the dimension of the vector space F_q^n over a finite field F_q). There is a necessary and sufficient condition on a finite semigroup S in order that $F_q[S]$ is semisimple. An explicit representation of such group is obtained in. They constructed all the irreducible representations of S from those of its principal factors of the full transformation semigroup on a finite set.

If F is algebraically closed, then there are no division algebras over F other than F itself, and in this case Wedderburn’s second theorem tells us that every simple algebra Λ over F is isomorphic with the full transformation semigroup algebra Λ of degree n for some positive integer n .

Any isomorphism of Λ upon semigroup Λ is a representation of Λ , and gives the irreducible representation of Λ . Let Λ be an algebra of order n over F , and let ϕ be a representation of Λ of degree r over F , and let m be a positive integer. For each element $\phi^{(m)}$ of $\mathcal{L}(\Lambda^m)$, construct a transformation $\phi^{(m)} \in \mathcal{L}(\Lambda^m(F^r))$.

such that

$$\Phi^{(m)} = \sum_{i=1}^r a_{mi} \Phi_i^{(m)}$$

$$\Phi_i^{(m)}, \Phi_j^{(m)} \in \mathcal{L}(\Lambda^m(F^r)),$$

if

then

$$\Phi^{(m)} = \sum_{\substack{i,j=1 \\ i+j=k}}^r a_{mi} b_{mj} \Phi_i^{(m)} \Phi_j^{(m)}$$

The map $\phi^{(m)}$ is called the representation of $L(L^m)$ associated with the representation ϕ of Λ . The following lemma is due to Van der Waerden’s modern algebra.

Lemma

Let D be division algebra, and let m be a positive integer. The right regular representation ρ of D is an irreducible, and the only irreducible representation of the simple algebra $\mathcal{L}(D^m)$ is just the representation $\rho^{(m)}$ of $\mathcal{L}(D^m)$ associated with ρ .

THEOREM 7.3

Let \wedge^σ ($\sigma=1, \dots, c$) be the simple components of a semisimple algebra \wedge . By Wedderburn's second theorem, each \wedge^σ may be regarded as a full transformation $\mathcal{L}(D^\sigma)^{m_\sigma}$ of some degree m_σ over the division algebra $\mathcal{L}(\wedge^\sigma)$. Let ρ_σ be the regular representation of D^σ and $\rho\sigma^{(m_\sigma)}$ be the representation of $\mathcal{L}(\wedge^\sigma)$ associated with $\rho\sigma$ then $\rho\sigma^{(m_\sigma)}$ is the only irreducible representation of $\rho\sigma$. Extending $(\rho\sigma)^{(m_\sigma)}$ to \wedge by defining $\phi\sigma(a) = (\rho\sigma)^{(m_\sigma)}(a)$ if $a = \sum_{r=1}^c a_r$ is the unique expression of the element a of \wedge as a sum of elements a_r of the \wedge^r . Then $\{\phi_1, \dots, \phi_c\}$ is the complete set of inequivalent irreducible representations of $D\sigma$. If $d\sigma$ is the order of D^σ , then the degree of ϕ_σ is $d_\sigma m_\sigma$. If F is algebraically closed, each D^σ reduces to F and we may regard L as a direct sum of full transformation semigroup algebra \wedge over F . The irreducible representation of \wedge are then just the projections of τ upon its various components (see Th.7.3 in [4]).

THEOREM 7.4

Let τ be a linear operator on \wedge with an algebra \wedge of finite order over a field F .

If $n > m$, then there exists a non-zero linear transformation $\sigma: \wedge^n \rightarrow \wedge^m$ such that $\tau\sigma = 0$. There exists a non-null transformation $\gamma: \wedge^n \rightarrow \wedge^m$ (over τ) such that $\gamma\tau = 0$, for every $m > n$.

Proof

Let $n > m$ and $\tau = \tau_1 \oplus \tau_2$ with τ_2 an operator on τ_2 and τ_2 a linear transformation from \wedge^{n-m} into \wedge^{n-m} (over τ_1). Suppose that τ_1 is left divisor of zero in $\mathcal{L}(\wedge^m)$, then there exists $\sigma_1 \neq 0$ in $\mathcal{L}(\wedge^m)$ such that $\tau_1\sigma_1 = 0$. We may take $\sigma = (\sigma_1, 0)$. Hence we may assume that τ_1 is not left divisor of zero in $\mathcal{L}(\wedge^m)$. By Lemma 5.8, that can be applied to the algebra $\mathcal{L}(\wedge^m)$, we have that the algebra $\mathcal{L}(\wedge^m)$ contains a left identity element i with respect to which τ_1 has a two-sided inverse ρ_1 in $\mathcal{L}(\wedge^m)$, i.e. $\rho_1\tau_1 = \tau_1\rho_1 = i$. We may take $\sigma = (\rho_1 \wedge \sigma_2, \sigma_2)$, where σ_2 is any non-singular linear transformation from \wedge^m into \wedge^m over the algebra \wedge .

Then,

$$\text{since } \tau_2 \sigma_2 \in \mathcal{L}(\wedge^m) \text{ and } i \text{ is the identity element in } \mathcal{L}(\wedge^m).$$

One can similarly prove that, if $m > n$, then there exists a non-null transformation $\gamma: \wedge^n \rightarrow \wedge^m$ such that $\gamma\tau = 0$

Representation of a full transformation semigroup over a finite field

Let θ be a root of some irreducible polynomial of degree m over a finite field F_q (or the Galois field $GF(q)$), then the set $\{1, \theta, \theta^2, \dots, \theta^{m-1}\}$ becomes a basis for the vector space F_q^m over F_q and is called a polynomial basis for F_q^m . The dimension of the vector space F_q^m over F_q is m . Let $\theta \in F_q^m$ such that the set

$$\mathcal{B} = \{\theta^{q^i} \mid 0 \leq i < m\} = \{\theta, \theta^q, \theta^{q^2}, \dots, \theta^{q^{m-1}}\}$$

form a basis for F_q^m . Let $a = \alpha = a_0\theta + a_1\theta^q + a_2\theta^{q^2} + \dots + a_{m-1}\theta^{q^{m-1}}$ so that a be represented by the vector $(a_0, a_1, a_2, \dots, a_{m-1})$ and let α^q be represented by the shifted vector $(a_{m-1}, a_0, a_1, \dots, a_{m-2})$. The normal basis exists for any extension field of F_q .

Consider the vector space $V = F_q^m$ over F_q (where q is a prime), and let $\mathcal{B} = \{\theta, \theta^q, \theta^{q^2}, \dots, \theta^{q^{m-1}}\}$ be a basis for V . Let $T_{\mathcal{B}}$ be the full transformation semigroup upon the basis \mathcal{B} . Then $|T_{\mathcal{B}}| = m^m$.

Since $\alpha = a_0\theta + a_1\theta^q + a_2\theta^{q^2} + \dots + a_{m-1}\theta^{q^{m-1}}$ is an element of $V = F_q^m$ as described above. Then the element $\sigma \in T_{\mathcal{B}}$ can be defined by $\sigma(\alpha) = \theta^q, \sigma^2(\alpha) = \theta^{q^2}, \dots, \sigma^{m-1}(\alpha) = \theta^{q^{m-1}}$. If $(a_0, a_1, a_2, \dots, a_{m-1}) \in V$, then $\sigma(a) \in T_{\mathcal{B}}$, where

$$\begin{aligned} \sigma(\alpha) &= \sigma(a_0, a_1, a_2, \dots, a_{m-1}) \\ &= (a_{m-1}, a_0, a_1, a_2, \dots, a_{m-2}), \end{aligned}$$

i.e.,
$$\sigma(\alpha) = \sigma \begin{pmatrix} a_0 & a_1 & a_2 & \dots & a_{m-1} \\ a_{m-1} & a_0 & a_1 & \dots & a_{m-2} \end{pmatrix} \in T_{\mathcal{B}}$$

It is obvious to say that $F_q^m = F_q^m$. S is a full transformation semigroup over V^* with a dual basis $\overline{\mathcal{B}} = \{\sigma_0 = 1, \sigma, \sigma^2, \dots, \sigma^{m-1}\}$ of V^* then there exists a mapping $\phi_a: T_{\mathcal{B}} \rightarrow S$ which becomes an isomorphism.

Since $T_{\mathcal{B}}$ is a finite full transformation semigroup on the basis \mathcal{B} of V over the finite field F_q . Therefore $F_q[T_{\mathcal{B}}]$ becomes an algebra of $T_{\mathcal{B}}$ over F_q . Then, there is a natural one-to-one correspondence between the representation of $T_{\mathcal{B}}$ over F_q and those of $F_q[T_{\mathcal{B}}]$, which preserves equivalence, reduction and decomposition into irreducible constituents.

Thus the representations of \mathcal{S} over F_q is transferred to the algebra $F_q[\mathcal{S}]$. If $F_q[\mathcal{S}]$ is semisimple, then by the main representation theorem[4] holds for semisimple algebra $F_q[\mathcal{S}]$. Every representation of $F_q[\mathcal{S}]$ and hence every representation of TB is full reducible into irreducible one.

Let F_q be a finite field, and B be a basis for F_q^m , where $(m,q) = 1$. (i.e., m,q are relatively prime).

Then, we have the following interpretation of the Maschke's theorem regarding the algebra $F_q[\mathcal{S}]$ over the finite field F_q .

THEOREM 8.1

Let $S = \mathcal{S}$ be a finite full transformation semigroup over basis \mathcal{B} of F_q^m of order mm.

Then, the semigroup algebra $F_q[\mathcal{S}]$ over F_q is semisimple if and only if the characteristic q of F_q does not divides the order mm of the full transformation semigroup.

Let \mathcal{A} be an algebra of order r over the vector space $V = F_q^m$, and let n be another positive integer different from m. Denote by $M_n(\mathcal{A})$ the full matrix algebra of all nn matrices over \mathcal{A} , with the additions and multiplication of matrices, and of the multiplication of matrix by a scalar in F_q . Then, the algebra $M_n(\mathcal{A})$ is of order rn^2 over F_q . In particular, $(F_q^m)_n$ will denote the full matrix algebra of degree n over F_q .

An algebra L over a field F is called division algebra if $L \setminus \{0\}$ is a group under multiplication. A result regarding the existence of an isomorphism between a full matrix algebra and the space of all the linear transformations over the vector space F_q^m , is as follows.

THEOREM 8.2

Let F_q^m be a vector space over a finite field F_q . Then, there is an isomorphism from the space of full matrix algebra $(F_q)_m$ to the space $\mathcal{L}(F_q^m)$ of all the linear transformations on F_q^m .

Proof

The set of all m-dimensional vector space (1m matrices) over F_q is an m-dimensional vector space F_q^m over F_q . The natural basis of F_q^m consists of the m vectors $v_1 = \theta, v_2 = \theta^q, v_3 = \theta^{q^2}, \dots, v_m = \theta^{q^{m-1}}$, where v_i has the identity element 1 of F_q for its ith component, and has 0 for the remaining components.

If $A \in (F_q)_m$, then the transformation $t : F_q^m \rightarrow F_q^m$ given by $t(v_i) = Av_i$ is a linear transformation t of F_q^m into itself and the mapping $\phi : (F_q)_m \rightarrow \mathcal{L}(F_q^m)$ is an isomorphism of $(F_q)_m$ upon the algebra $\mathcal{L}(F_q^m)$ of all linear transformations of F_q^m into itself. The ith row of A is the vector $t(v_i)$.

Conversely, if F_q^m is any m-dimensional vector space, and we choose a basis $\{v_1, v_2, \dots, v_m\}$ of F_q^m , then each linear transformation t of F_q^m determines a matrix $A = (\alpha_{ij})$ from the expression

for the m vectors $t(v_i); (1 \leq i \leq m)$ as linear combination of the basis vectors. Then, the mapping $\psi : \mathcal{L}(F_q^m) \rightarrow (F_q)_m$ becomes an isomorphism of $\mathcal{L}(F_q^m)$ upon $(F_q)_m$.

CONCLUSION

A combinatorial result about the rank of a representation of the full transformation semigroup is obtained. It seems that for any homomorphism between the set of single-valued maps and the set of all nn matrices over a field F becomes a representation when the set of single valued maps is replaced by a full transformation semigroup adjoined with a zero element z. There is a one-one correspondence between the set of all representations of some finite semigroup S and those of the algebra of a full transformation semigroup over a finite dimensional vector space over a finite field. Consequently, we observed an isomorphism between the full matrix algebra $(F_q)^m$ and the set of all linear transformations on F_q^m is obtained.

REFERENCES

1. Serre JP. Linear representations of finite groups, translated from the second French edition by Leonard L. Scott. Graduate Texts in Mathematics. 1977;42:172.
2. Clifford AH. Matrix representations of completely simple semigroups. American J Math. 1942;64:327-342.
3. Clifford AH. Basic representations of completely simple semigroups. American J Math. 1960;82:430-434.
4. A. H. Clifford and G. B. Preston, The algebraic theory of semigroups, Mathematical surveys of the Amer. Math. Soc. 1961.
5. Stoll RR. Representations of finite simple semigroups. Duke Mathe J. 1944;11:251-265.
6. W. D. Munn, Irreducible matrix representations of semigroups, Quarterly J. Math. Oxford. 1960;11:295-309.

7. W. D. Munn, On semigroup algebras, Proc Cambridge Phil. Soc. 1955b;51:1-15.
8. W. D. Munn, Semigroups and their algebras, Dissertation, Cambridge University. 1955a.
9. Hewitt E, et al. The irreducible representations of a semigroup related to the symmetric group. Illinois J Math. 1957;1:188-213.
10. G. L. Mullen, et al. Finite Fields and Applications, Amer. Math. Soc. Student Mathematical library. 2007;41.
11. Lallement G, Petrich M. Irreducible matrix representations of finite semigroups. Transactions of the American Math Soc. 1969;139:393-412.
12. O. Ganyushkin, et al. Classical Finite Transformation Semigroups: An Introduction, Springer Science and Business Media. 2008.
13. Munn WD. Matrix representations of semigroups. In Mathematical Proceedings of the Cambridge Philosophical Society. Cambridge University Press. 1957; 53:5-12.
14. Nagy A. On faithful representations of finite semigroups S of degree $|S|$ over the fields. Int J Algebra. 2013;7:115-29.
15. Nagy A, et al. Finite Semigroups whose semigroup algebra over a field has a trivial right annihilator. Int J Contemporary Mathe Sci. 2014;9:25-36.
16. R. J. Warne, Matrix Representation of d -simple semigroups, Trans. Amer. Math. Soc. 1963;106:427-435.
17. N. L. Biggs, Finite group of Automorphisms, Cambridge University press, London. 1971;13.