

A Potential Application of Algae as a Complementary/Integrative Therapy in Veterinary Medicine

Susumu Ishiguro, Damilola Gbore, Grace Calo, and Masaaki Tamura*

Department of Anatomy & Physiology, Kansas State University College of Veterinary Medicine, Manhattan, KS 66506, United States

Review Article

Received: 1/09/2021

Accepted: 15/09/2021

Published: 22/09/2021

***For Correspondence:**

Masaaki Tamura,
Department of Anatomy &
Physiology, Kansas State University
College of Veterinary Medicine,
Manhattan, KS 66506, United
States

E-mail: mtamura@vet.ksu.edu

Keywords: Algae; Therapeutic potential; Potential Application; Integrative Health; Complementary therapies

ABSTRACT

Although the nutritional and medicinal properties of algae have been studied for the last several decades, its value in veterinary medicine is widely unknown. Algae are found in fresh water, saltwater and soil, and they are rich in diversity. It has been shown that algae contain various nutrients that are beneficial for human and animal health as well as food/feed industries. Alga's whole bodies, their extract, and bioactive molecules purified from algae have shown medicinal properties that are potentially beneficial for animal health, however, the usage of algae is still in confusions. In this minireview, potential applications of algae in veterinary medicine and its nutritional value for domestic animals are discussed. In addition, the possible existence of novel immune modulative materials in cell wall membrane of *Chlorella sorokiniana* will be discussed based on our recent discovery.

INTRODUCTION

Over the years, there has been vast research into the use of natural products for medicinal purposes [1]. Algae are some of such naturally occurring resources that show huge promise. "Algae" is a general term for

photosynthetic eukaryotic organisms that can be found throughout the world. The organisms in this group consist of a wide variety of species; from micrometer-sized unicellular microalgae such as *Chlorella*, to meter-sized multicellular macroalgae such as seaweed. Algae have been reported to contain large amounts of various beneficial components, such as proteins, peptides, amino acids, beta-1,3-glucan, vitamins, polyunsaturated fatty acids, polysaccharides, etc. [2]. Therefore, the whole cell powder or crushed cell body powder of algae is taken as a nutritional and functional dietary supplement and contributes to human health [2]. It has been shown that a dried powder of microalgae or its extracts in water or polar organic solvents has potential medicinal properties, such as anticancer [3], antimicrobial [4], antiviral [5] antioxidant [6], anti-inflammatory [7], anti-obesity [8], etc. These medicinal properties of microalgae can potentially develop into novel therapeutics in the field of both human and veterinary medicine. Microalgae and related products are also usable as a potential resources of feed and nutrition for livestock and poultry [9] as well as pets [10-12].

MICROALGAE IN VETERINARY MEDICINE

There are potentially wide applications that can be derived from microalgae, particularly *Chlorella* in the treatment of diseases such as inflammation, cancer, and infectious disease. The use of microalgae as a nutrient source for pets was evaluated and deemed nutritionally beneficial. Consequently, pet foods containing microalgae are commercially available as healthy functional food for dogs and cats. Souza *et al.* (2019) used green microalgae *Schizochytrium* sp. as a source of docosahexaenoic acid (DHA) and evaluated the aptitude as a dietary component and the effect on immunity in dogs [13]. DHA is one of the polyunsaturated fatty acids (PUFA) which is known to be an essential nutrient for dog's growth and survival. In addition, DHA is known to improve immune functions in dogs and other species [14, 15]. However, since the conversion of alpha-linolenic acid to DHA is inefficient in dogs [16] the effect of dietary supplementation of DHA was evaluated in this study. Supplementation of 0.4 % of *Schizochytrium* sp. in the diet as a source of DHA increases the digestibility of nutrients and metabolizable energy. It also increases the number of phagocytic cells and the intensity of phagocytosis in monocytes in the blood. Notably, it does not alter dog's palatability, fecal characteristics, or biochemical blood parameters. As a source of DHA, *Schizochytrium* sp. was also evaluated in cats [17]. In this study, the influence of DHA-rich microalgae diet on the serum level of inflammatory markers including prostaglandin E2 (PGE2), acute-phase proteins (ceruloplasmin, haptoglobin, α -1-acid glycoprotein, albumin and transferrin) and 12-hydroxyeicosatetraenoic acid (12-HETE), before and after neutering was evaluated. The DHA-rich microalgae diet did not affect serum level of 12-HETE, while serum level of PGE2 and albumin were decreased significantly. The number of platelets in blood was observed as an indicator of local inflammation. Feeding of the DHA-rich microalgae diet increased basal level of platelet count. Although platelet count was significantly decreased after neutering in both normal and DHA-rich diet groups, platelet count in the DHA-rich diet group was significantly higher as compared to the normal diet group. In this study, dried *Schizochytrium* sp. containing DHA (*n*-3 fatty acid) was supplemented by replacing poultry fat (*n*-6 fatty acid). It is known that *n*-3 fatty acid has an anti-aggregatory activity, whereas *n*-6 fatty acid has a pro-aggregatory activity. The result suggests that supplementation of DHA (*n*-3 fatty acid)-rich *Schizochytrium* sp. decreased the ratio of *n*-6 fatty acid in the diets, therefore anti-aggregatory activity become predominant in the blood of cats served with a DHA-rich microalgae diet. These results also suggest

that a DHA-rich microalga diet regulates the inflammatory response that occurs after neutering. Another study showed that beta-1,3-glucan stimulates both adaptive and innate immunity in a species of fish [18]. Beta-1,3-glucans are one of the most abundant polysaccharides in algae [19]. This may suggest that feeding or administering *Chlorella* or its extract, being a rich source of beta-1,3-glucan [20] can enhance disease resistance in fish. In another study, the immunomodulatory activity of whole blue-green alga in humans and different animals, such as cats and chickens, was described. Because of the high nutritional value of microalgae, application to livestock feed is noted. Jeon *et al.* (2016) evaluated the dietary effect of lutein-fortified *Chlorella* on the milk production and components of Holstein cows [21]. In this study, three weeks feeding of conventional (15 g/day) or lutein-fortified *Chlorella* (30 g/day) [22] increased concentrations of milk protein and solids non-fat without changing milk yield by either feeding of conventional or lutein-fortified *Chlorella* compared to control. The feeding of lutein-fortified *Chlorella* increased lutein content in milk compared to conventional *Chlorella* and control. Lutein is known to have antioxidant properties [23]. Therefore, these results suggest that *Chlorella* is capable of increasing nutrient components in the milk as well as functional materials, such as antioxidants in dairy products. Similar results were reported by Lamine *et al.* (2019), however researchers pointed out the necessity of improving microalgae palatability [24]. The application of microalgae for livestock feed is also attempted in pigs [25], broiler chickens [26], goats [27], etc.

FUCOIDAN FROM MACROALGAE IN VETERINARY MEDICINE

Fucoidan, a long chain sulfated polysaccharide, has been found in various species of brown macroalgae, such as *Hizikia fusiformis* (Hijiki), *Cladosiphon okamuranus* (Mozuku), *Laminaria japonica* (Kombu), *Undaria pinnatifida* (Wakame), etc. [28]. These brown macroalgae are common food in East Asia and have been cultivated in that area. Multiple medicinal properties of fucoidan have been reported in both human and veterinary medicine [29].

Laura *et al.* (2014) demonstrated that fucoidan extracted from *Cladosiphon okamuranus* inhibits canine distemper virus (CDV) infection *in vitro* by interference in the early stage of CDV infection and by inhibiting CDV-mediated cell fusion [30]. The cytotoxicity of fucoidan (CC50 = 2,089 ± 6 µg/ml) was much lower than that by antiviral drug, ribavirin (CC50 = 88.9 ± 6 µg/ml). Fucoidan treatment (0.1–10 µg/ml) inhibited CDV proliferation in CDV-infected cells. Both pretreatment and simultaneous treatment with various concentrations of fucoidan (0.001–10 µg/ml) inhibited CDV infection into cells. Fucoidan treatment (0.001–10 µg/ml) inhibited cell-cell fusion of CDV-infected cells, which is an important mechanisms for cell to cell spread of progeny viruses.

The property of fucoidan as a prebiotic was also investigated in domestic animals [31]. Many researchers evaluated the effect of fucoidan and laminarin on the gut microbiota in piglets. Laminarin is storage form of glucan found in brown algae and known with fucoidan as a mucopolysaccharide. A notable effect of feeding fucoidan and laminarin in combination to sows is the reduction of the gut *Escherichia coli* (*E. coli*) population in their piglets [32, 33]. Leonard *et al.* (2011) reported that the effect of maternal dietary supplementation with a seaweed extract (10 g/day) composed of 1 g laminarin, 0.8 g fucoidan, and 8.2 g ash on the *E. coli* population in intestinal microbiota of post-weaning piglets. The population of *E. coli* in the caecum and the colon was reduced 16 % compared to piglets from the basal diet fed sow at 9 days post weaning. This effect of fucoidan + laminarin on *E. coli* population in piglets was also observed in separate

studies [34, 35]. Since it is well known that *E. coli* have various pathogenic strains and often causes diarrhea in piglets [36] these results strongly suggest that supplementations of fucoidan and/or laminarin are beneficial to piglet health and the industry.

BLUE-GREEN ALGAE (CYANOBACTERIA) IN VETERINARY MEDICINE

Cyanobacteria are Gram-negative bacteria. Algae belong to eukaryotes, while cyanobacteria belong to prokaryotes. Therefore, these are different organisms. However, cyanobacteria share similar features with green algae, such as photosynthetic properties, their habitat, and their color; blue (phycocyanin) and green (chloroplast). Therefore, cyanobacteria are also called as blue-green algae.

Spirulina refers to the biomass of *Arthrospira platensis* which is one of the species of cyanobacteria. Because of a high nutritional value due to proteins, carbohydrates, fatty acids, vitamins, minerals, etc., whole spirulina powder or its tablet form is commonly consumed as food supplement for human and animals [37]. In addition, therapeutic properties such as antioxidant [38], antiviral [39], anti-inflammatory and immunomodulating [40], anti-obesity [41], anticancer [42], etc. have been reported. In the veterinary field, the effect of spirulina on the gut microbiota was evaluated in dogs [43] and broiler chickens [44].

On the other hand, hazardous properties of cyanobacteria are reported. Cyanobacterial blooms cause severe damage to organisms inhabiting the environment, and economic activities benefitting the environment [45]. Some of cyanobacteria produce a toxin called cyanotoxin. Microcystin produced by *Microcystis aeruginosa* is known to cause serious damage to the liver [46]. Saxitoxin produced by *Alexandrium catenella* is also known as shellfish poisoning, which is caused by shellfish that ate cyanobacteria with their toxin. This is also harmful for livestock, poultry, and pets [47]. In the case of dogs, symptoms appear after exposure to freshwater with cyanobacterial blooms [48, 49].

POTENTIAL BENEFITS OF *CHLORELLA* IN HUMAN AND VETERINARY MEDICINE

One microalgae that has recently become popular among the microalgae populations is *Chlorella*, a single-celled green algae renowned for its highly nutritious and commercial benefits [50]. *Chlorella* is one genus of microalga living in fresh water. The genus of *Chlorella* is composed of more than 20 species and the species are further divided into three varieties: *C. vulgaris*, *C. lobophora*, and *C. sorokiniana* [51]. *Chlorella* has potential protective functions such as immunostimulatory, anticancer, anti-inflammatory, antiviral, antimicrobial, anti-fungal, anti-diabetic, anti-hypercholesterolemic, and anti-atherosclerotic effects, among others, as have been described [51]. Being a natural and less expensive source, *Chlorella* and its extracts could potentially provide a good supplement or alternative to more expensive conventional therapies which may also have high incidence of side effects in treatment of animals. There is a definite need for more research to further evaluate the potential benefits that could be derived from *Chlorella* and other valuable microalga in companion, production, and other categories of animals as a step towards improving animal health, production, and welfare. Studies should also be conducted to determine safety and tolerance levels in different species as there have been reports of side effects in humans, such as allergies, nausea, vomiting, and other gastrointestinal problems [52] as well as acute tubulointerstitial nephritis [53].

LPS-LIKE MOLECULE IN *CHLORELLA*

Bacterial lipopolysaccharide (LPS) is an endotoxin found in the cell wall of gram-negative bacteria. It shows various bioactivities in human and animals *via* cell surface receptor, Toll-like receptor 4 (TLR4). LPS stimulates antigen presenting cell proliferation and induces the secretion of pro-inflammatory cytokines from them. In the *Chlorella* species, the presence of a LPS-like molecule in the cell wall of *Chlorella* has been demonstrated using immunohistochemical techniques [54]. In a recent study, our research team demonstrated that a bioactive factor partially purified from the isolated cell wall membrane fraction of *Chlorella sorokiniana* (*Chlorella* membrane factor, abbreviated as CMF) attenuated the growth of colon carcinoma *via* an alteration of host anticancer immunity [55]. The CMF showed opposite effects compared to bacterial LPS on the CT26 murine colon carcinoma cell growth, in which CMF attenuates cell growth, whereas the LPS stimulates the growth of the cancer cells. It is, therefore, suggested that the action of CMF is functionally different from bacterial LPS in stimulation of cancer cell growth. On the other hand, CMF showed an immune stimulating effect in cell culture and mouse. In the study with the three-dimensional cancer spheroid co-culturing with T lymphocytes, CMF stimulated functional differentiation of T lymphocytes, thereby inhibiting growth of colon carcinoma cells. In a mouse study, the intraperitoneal administration of CMF (10 or 30 mg/kg) was carried out using CT26 cell intraperitoneal dissemination mouse model. The analysis of immune cell population in the mouse ascites, CD4+ and CD8+ T cell, and CD19+ B cell populations were particularly increased, while CD68+ macrophage population was decreased in ascites in the mice treated with 30 mg/kg CMF. The LY6G+ granulocyte (neutrophil) population in ascites was also decreased in mouse groups treated with 10 or 30 mg/kg CMF. These two cell populations, macrophages and neutrophils, were known to participate in tumor growth [56, 57]. Therefore, these results suggest that CMF treatment cause modulation of anticancer immunity by increasing anti-tumorigenic effector T cells and decreasing pro-tumorigenic immune cell populations. It is also suggested that *Chlorella* cell wall membrane may be a potential source for novel therapeutics with immunomodulatory properties.

DISCUSSION AND CONCLUSION

In conclusion, early research supports that algae may be a very useful complementary/alternative treatment, applicable to a wide array of medical issues due to its immunostimulatory, anticancer, anti-inflammatory, antiviral, antimicrobial, anti-fungal, anti-diabetic, anti-hypercholesterolemic, and anti-atherosclerotic properties. However, most research regarding the above-described bioactivities has not been rigorously evaluated by traditional pharmacokinetics and pharmacodynamics studies and patient-based clinical trials. The issue of the biosafety is another concern; since many studies have evaluated the bioactivities of algae using whole algae without purification of bioactive components, potential toxic components in microalgae would be higher when the intake of algae increases. Therefore, it is critically important to conduct additional research to evaluate the topic properly.

ACKNOWLEDGEMENTS

We thank Ms. Nicole Robben (Department of Anatomy and Physiology, Kansas State University College of Veterinary Medicine) for her critical reading and review of this manuscript.

DISCLOSURE OF ANY POTENTIAL CONFLICTS OF INTEREST

No potential conflicts of interest were disclosed by all authors.

FUNDING

This work was supported in part by Kansas State University (KSU) College of Veterinary Medicine Dean's funds [2018 CVM-SMILE] (MT), Euglena Research Fund (MT), KSU Johnson Cancer Research Center [2020 JCRC-CRA] (MT), K-INBRE Bridging Award [P20 GM103418] (MT).

REFERENCES

1. [Newman DJ, et al. Natural products as sources of new Drugs over the nearly four decades from 01/1981 to 09/2019. J Nat Prod. 2020; 83:770-803.](#)
2. [Caporgno MP, et al. Trends in microalgae incorporation into Innovative food products with potential health benefits. Front Nutr. 2018; 5:58.](#)
3. [Galasso C, et al. Microalgal derivatives as potential nutraceutical and food supplements for human health: A focus on cancer prevention and interception. Nutrients. 2019; 11:1226.](#)
4. [Falaise C, et al. Antimicrobial compounds from eukaryotic microalgae against human pathogens and diseases in aquaculture. Mar Drugs. 2016; 14:159.](#)
5. [Carbone DA, et al. Evaluation of microalgae antiviral activity and their bioactive compounds. Antibiotics \(Basel\). 2021; 10:746.](#)
6. [Sansone C, et al. Promises and challenges of microalgal antioxidant production. Antioxidants \(Basel\). 2019; 8:199.](#)
7. [Tabarзад M, et al. Anti-inflammatory activity of bioactive compounds from microalgae and cyanobacteria by focusing on the mechanisms of action. Mol Biol Rep. 2020; 47:6193-6205.](#)
8. [Gomez-Zorita S, et al. Anti-Obesity effects of macroalgae. Nutrients. 2020; 12:2378.](#)
9. [Saadaoui I, et al. Microalgal-based feed: promising alternative feedstocks for livestock and poultry production. J Anim Sci Biotechnol. 2021; 12:76.](#)
10. [Vuorinen A, et al. Safety of algal oil containing EPA and DHA in cats during gestation, lactation and growth. J Anim Physiol Anim Nutr \(Berl\). 2020; 104:1509-1523.](#)
11. [Dahms I, et al. Safety of a novel feed ingredient, algal oil containing EPA and DHA, in a gestation-lactation-growth feeding study in beagle dogs. PLoS One. 2019; 14:e0217794.](#)
12. [McCusker S, et al. Amino acid content of selected plant, algae and insect species: a search for alternative protein sources for use in pet foods. J Nutr Sci. 2014; 3:e39.](#)

13. [Souza CMM, et al. Microalgae schizochytrium sp. as a source of docosahexaenoic acid \(DHA\): Effects on diet digestibility, oxidation and palatability and on immunity and inflammatory indices in dogs. Anim Sci J. 2019; 90:1567-1574.](#)
14. [Kelley DS, et al. DHA supplementation decreases serum C-reactive protein and other markers of inflammation in hypertriglyceridemic Men. J Nutr. 2009; 139:495-501.](#)
15. [Heinemann KM, et al. Long-chain \(n-3\) polyunsaturated fatty acid are more efficient than alpha-Linolenic acid in Improving electroretinogram responses of puppies exposed during gestation, lactation, and weaning. J Nutr. 2005; 135:1960-1966.](#)
16. [Bauer JE, et al. Responses of dogs to dietary omega-3 fatty acids. J Am Vet Med Assoc. 2007; 231:1657-1661.](#)
17. [Scheibel S, et al. DHA from microalgae schizochytrium spp. \(Thraustochytriaceae\) modifies the inflammatory response and gonadal lipid profile in domestic cats. Br J Nutr. 2021; 126:172-182.](#)
18. [Sahoo PK, et al. Effect of dietary beta-1,3 glucan on immune responses and disease resistance of healthy and aflatoxin B1-induced immunocompromised rohu \(Labeo rohita Hamilton\). Fish Shellfish Immunol. 2001; 11:683-695.](#)
19. [Wang Q, et al. beta-Glucans: Relationships between modification, conformation and functional activities. Molecules. 2017; 22:257.](#)
20. [Yaakob Z, et al. An overview: biomolecules from microalgae for animal feed and aquaculture. J Biol Res \(Thessalon\). 2014; 21:6.](#)
21. [Jeon JY, et al. Dietary effects of lutein-fortified chlorella on milk components of holstein cows. Springerplus. 2016; 5:908.](#)
22. [Jeon JY, et al. Optimization of culture media for large-scale lutein production by heterotrophic Chlorella vulgaris. Biotechnol Prog. 2014; 30:736-743.](#)
23. [Obana A, et al. Effect of an antioxidant supplement containing high dose lutein and zeaxanthin on macular pigment and skin carotenoid levels. Sci Rep. 2020; 10:10262.](#)
24. [Lamminen M, et al. Different microalgae species as a substitutive protein feed for soya bean meal in grass silage based dairy cow diets. Animal Feed Science and Technology. 2019; 247:112-126.](#)

25. [Coelho D, et al. A High dietary Incorporation level of Chlorella vulgaris improves the nutritional value of pork fat without impairing the performance of finishing pigs. Animals \(Basel\). 2020; 10: 2384.](#)
26. [Alfaia CM, et al. Influence of dietary Chlorella vulgaris and carbohydrate-active enzymes on growth performance, meat quality and lipid composition of broiler chickens. Poult Sci. 2021; 100:926-937.](#)
27. [Kholif A, et al. Dietary chlorella vulgaris microalgae improves feed utilization, milk production and concentrations of conjugated linoleic acids in the milk of damascus goats. The Journal of Agricultural Science. 2017; 155:508-518.](#)
28. [Fitton HJ, et al. Therapies from Fucoidan: New Developments. Mar Drugs. 2019; 17: 571.](#)
29. [Kwak JY, et al. Fucoidan as a marine anticancer agent in preclinical development. Mar Drugs. 2014; 12:851-870.](#)
30. [Trejo-Avila LM, et al. In vitro anti-canine distemper virus activity of fucoidan extracted from the brown alga Cladosiphon okamuranus. Virusdisease. 2014; 25:474-480.](#)
31. [O'Sullivan L, et al. Prebiotics from marine macroalgae for human and animal health applications. Mar Drugs. 2010; 8:2038-2064.](#)
32. [Leonard SG, et al. Effect of maternal seaweed extract supplementation on suckling piglet growth, humoral immunity, selected microflora, and immune response after an ex vivo lipopolysaccharide challenge. J Anim Sci. 2012; 90:505-514.](#)
33. [Leonard SG, et al. Effect of dietary seaweed extracts and fish oil supplementation in sows on performance, intestinal microflora, intestinal morphology, volatile fatty acid concentrations and immune status of weaned pigs. Br J Nutr. 2011; 105:549-560.](#)
34. [Dillon S, et al. The effects of lactose inclusion and seaweed extract on performance, nutrient digestibility and microbial populations in newly weaned piglets. Livestock Science. 2010; 134:205-207.](#)
35. [McDonnell P, et al. The effect of dietary laminarin and fucoidan in the diet of the weanling piglet on performance, selected faecal microbial populations and volatile fatty acid concentrations. Animal. 2010; 4:579-585.](#)
36. [Melin L, et al. Weaning of piglets. Effects of an exposure to a pathogenic strain of Escherichia coli. J Vet Med B Infect Dis Vet Public Health. 2000; 47:663-675.](#)
37. [Grosshagauer S, et al. The true value of spirulina. J Agric Food Chem. 2020; 68:4109-4115.](#)

38. [Seyidoglu N, et al. The Preventive role of spirulina platensis \(Arthrospira Platensis\) in immune and oxidative insults in a stress-induced rat model. J Vet Res. 2021; 65:193-200.](#)
39. [Petit L, et al. Docking and in silico toxicity assessment of arthrospira compounds as potential antiviral agents against SARS-CoV-2. J Appl Phycol. 2021; 1-24.](#)
40. [Keller M, et al. Antioxidant and inflammatory gene expression profiles of bovine peripheral blood mononuclear cells in response to arthrospira platensis before and after LPS challenge. Antioxidants \(Basel\). 2021; 10:814](#)
41. [Diniz AFA, et al. Spirulina platensis consumption prevents obesity and improves the deleterious effects on Intestinal Reactivity in Rats Fed a hypercaloric diet. Oxid Med Cell Longev. 2021; 2021:3260789.](#)
42. [Subramaiam H, et al. Evaluating anticancer and immunomodulatory effects of spirulina \(Arthrospira\) platensis and gamma-tocotrienol supplementation in a syngeneic mouse model of breast cancer. Nutrients. 2021; 13:2320.](#)
43. [Satyaraj E, et al. Supplementation of diets with spirulina influences immune and gut function in dogs. Front Nutr. 2021; 8:667072.](#)
44. [Chang CT, et al. Effects of mesobiliverdin IXalpha-Enriched microalgae feed on gut health and microbiota of Broilers. Front Vet Sci. 2021; 7:586813.](#)
45. [Piel T, et al. Resilience of microbial communities after hydrogen peroxide treatment of a eutrophic lake to suppress harmful cyanobacterial blooms. Microorganisms. 2021; 9:1495](#)
46. [Zheng S, et al. Effects of environmental contaminants in water resources on nonalcoholic fatty liver disease. Environ Int. 2021; 154:106555.](#)
47. [Wood R, et al. Acute animal and human poisonings from cyanotoxin exposure - A review of the literature. Environ Int. 2016; 91:276-282.](#)
48. [Fastner J, et al. Fatal neurotoxicosis in dogs associated with tychoplanktic, anatoxin-a producing tychonema sp. in mesotrophic lake tegel, berlin. Toxins \(Basel\). 2018; 10:60.](#)
49. [Puschner B, et al. Debromoaplysiatoxin as the Causative Agent of Dermatitis in a Dog after Exposure to Freshwater in California. Front Vet Sci. 2017; 4:50.](#)
50. Rani K, et al. A comprehensive review on chlorella-its composition, health benefits, market and regulatory scenario. The Pharma Innovation Journal. 2018; 7:584-589.
51. [Bito T, et al. Potential of chlorella as a dietary supplement to promote human health. Nutrients. 2020; 12: 2524.](#)

52. [Tiberg E, et al. Allergy to green algae \(Chlorella\) among children. J Allergy Clin Immunol. 1995; 96:257-259.](#)
53. [Yim HE, et al. Acute tubulointerstitial nephritis following ingestion of Chlorella tablets. Pediatr Nephrol. 2007; 22:887-888.](#)
54. [Armstrong MT, et al. Histochemical evidence for lipid A \(endotoxin\) in eukaryote chloroplasts. FASEB J. 2006; 20:2145-2146.](#)
55. [Ishiguro S, et al. Cell wall membrane fraction of chlorella sorokiniana enhances host antitumor immunity and inhibits colon carcinoma growth in mice. Integr Cancer Ther. 2020; 19:1534735419900555.](#)
56. [Fridlender ZG, et al. Polarization of tumor-associated neutrophil phenotype by TGF-beta: "N1" versus "N2" TAN. Cancer Cell. 2009; 16:183-194.](#)
57. [Allavena P, et al. The Yin-Yang of tumor-associated macrophages in neoplastic progression and immune surveillance. Immunol Rev. 2008; 222:155-161.](#)