

## Double seronegative myasthenia gravis: a case report- Linda Carolina Jaramillo- Industrial University of Santander

Linda Carolina Jaramillo, Jenny Paola Garzon and Gustavo Pradilla

Industrial University of Santander, Colombia

A 54-year-old man with chronic exposure to carbamates and organophosphates was referred from the otorhinolaryngology outpatient section in 2014, because of 3 years of dyspnea and dysphonia of spasmodic features, phonasthenia without respiratory distress or dysphagia; associated to progressive weakness in lower limbs. Brain magnetic resonance, angioresonance and chest x-ray had no alterations; laryngeal electromyography revealed a bilateral neuropathic pattern without denervatory activity, with some signs of reinnervation and decreased recruitment. Lambert test displayed motor and sensory distal latencies of median nerves prolonged. Repetitive stimulation test evidenced a decrease greater than 10% in the trapezium, concluding an abnormal repetitive stimulation test, concluding a conduction disorder in the neuromuscular junction. With a diagnostic impression of myasthenia gravis, the patient started a therapeutic test with pyridostigmine worsening the symptoms, so he was

switched to azathioprine and corticosteroids. His anti-MuSK antibodies  $<0.05$  mmol/L and antibodies against acetylcholine receptors  $<15\%$  were interpreted as negative. Due to worsening of the bulbar symptoms, plasmapheresis was performed; after the first session he showed improvement in lower limbs weakness, so five sessions were carried out with continuous improvement. Myasthenia gravis and its subcategories are major diseases that affect the neuromuscular junction. The diagnosis is confirmed by the combination of relevant symptoms and signs, neurophysiological studies and a positive test for specific autoantibodies. About 10% of patients with generalized myasthenia gravis do not have detectable antibodies to acetylcholine receptors or muscle specific kinase (MuSK) (double seronegative myasthenia), so neurophysiological tests and a positive response to therapy secure the diagnosis.