

Research & Reviews: Journal of Microbiology and Biotechnology

Potentially beneficial rhizobacteria associated with banana plants in Juja Kenya

Onchwari Ruth Gechemba¹, Mugweru Julius¹, Huxley Mae Makonde², Nancy L M Bubambula^{3*}, Vivienne N Matiru^{1*}

¹Jomo Kenyatta University of Agriculture and Technology, School of Biological Sciences, P.O. Box 62000-00200 Nairobi, Kenya

²Pure and Applied Sciences, Technical University of Mombasa, Kenya

³Embu University College, Department of Biological Sciences, P.O. Box 6-60100 Embu, Kenya

Research Article

Received date: 06/05/2015

Accepted date: 02/16/2016

Published date: 02/26/2016

*For Correspondence

Jomo Kenyatta University of Agriculture and Technology, School of Biological Sciences, P.O. Box 62000-00200 Nairobi, Kenya, Tel: +254 722 524 051

E-mail: vivienne.matiru@yahoo.com

Keywords: Rhizobacteria; Rhizosphere; Biofertilizers; *Musa* spp.

ABSTRACT

Banana being the world's third most important starchy food and widely grown in developing countries requires large amounts of nitrogen fertilizers for commercial cultivation which are costly and hazardous to the environment when used in excess. Plant growth promoting rhizobacteria are bacteria found in the rhizosphere in association with roots and can enhance the growth of the plant. We isolated and identified the rhizobacteria associated with bananas (*Musa* spp.) from rhizosphere soil samples collected from seven banana farms in Juja, Kenya and evaluated their potential as biofertilizers. We obtained 20 bacterial isolates and 19 had the potential to fix nitrogen as observed from acetylene reduction assay while 18 had phosphate solubilisation activities. Phylogenetic analysis clustered the isolates into three phyla represented by six different genera: *Pseudomonas*, *Bacillus*, *Staphylobacterium*, *Chryseobacterium*, *Streptomyces* and *Paenibacillus* spp. This work has shown that banana rhizospheric bacterial isolates harbor plant growth promoting traits that can be utilized as biofertilizers in agriculture to promote the growth as well as production of bananas.

INTRODUCTION

Rhizosphere is the thin layer of soil that immediately surrounds the plant roots ^[1] and has abundant nutritious elements because of the continuous rot activities of the plants. This zone is mixed with solid particles and an active community of free living beneficial bacteria ^[2], usually known as plant-growth-promoting rhizobacteria (PGPR).

The banana plant (*Musa* spp.) has adventitious and horizontal roots proliferating the topsoil and cannot get water and nutrients from the deeper soil profile unlike other fruit crops. The undeveloped root system inhibits the utilization of essential mineral nutrients thus limiting the large scale production of bananas under adverse tropical soil conditions. In the recent past there has been an increasing interest in soil microorganisms due to their importance in maintaining soil fertility since it has been shown that plant growth may be stimulated by vitamin-related and pathogen suppressing phytohormones produced by rhizospheric bacteria ^[3]. The mutual association where the bacteria obtain the nutrients released from the roots of the plants for growth as they in turn secrete metabolites to the rhizosphere ^[4], is perceived to augment plant growth by the rhizobacteria directly acting as biofertilizers to provide nutrients to the host plant or indirectly by their positive influence on the growth and morphology of roots ^[2].

Previously, Ngamau et al. and Reena et al. have demonstrated diverse banana endophytic bacteria with potential in utilization of the limiting nitrogen and phosphorous nutrients in banana plant growth ^[5,6]. We postulate that the tropical banana rhizosphere harbor's a wide diversity of PGPR that may not only aid in beneficial symbiotic relationships but may stimulate the plant growth

by suppressing pathogenic organisms. Since biofertilizers are widely accepted as a source of fertilizers with significant increase in crop yields ^[7], we set herein to characterize the PGPR associated with banana roots and determine their potential use as a biofertilizer in banana growth.

MATERIALS AND METHODS

Sample collection, isolation and profiling

Soil samples from the rhizosphere region were collected during the month of June 2012 from the cooking and ripening varieties of *Musa* spp. from seven farms in Juja, Kenya. They were collected by digging at 20-40 cm surrounding the banana plant roots and scrubbing the soil attached to the roots. The rhizosphere soil was scrubbed off the roots, serially diluted fivefold and inoculated on nitrogen-free media, yeast extract mannitol agar and nutrient agar (HIMEDIA) for 24 hours at 30°C. Individual colonies were aseptically picked and streaked on sterile fresh media to purify them and generate pure cultures for morphological, biochemical and molecular characterization. The pure cultures were Gram classified and tested for: hydrogen sulphide production, catalase utilization, urease production, nitrate reduction, IMViC tests, starch, casein and gelatin hydrolysis using standard methods as described by Cappuccino and Sherman (2002) ^[8].

Characterization of rhizobacteria for plant growth promoting traits

The isolates were assessed for nitrogenase activity based on their ability to reduce acetylene (C₂H₂) to ethylene (C₂H₄) by acetylene reduction assay by growing an isolate for 72 hours at 30°C in 5 ml nitrogen free semisolid agar in 10 ml vials. Acetylene was added at 12% v/v concentration and the ethylene produced was determined on a Shimadzu Gas Chromatograph (GC-9A, Japan) after 12 hours ^[9]. Using a syringe, 1 ml of ethylene produced was injected into the GC machine and the retention time for each isolate measured.

The isolates were screened for phosphate solubilisation using the National Botanical Research Institute's phosphate (NBRIP) growth medium ^[10]. Formation of visible halo zones on agar plates, which is as a result of organism's production of organic acids into the surrounding medium, was used as a measure of relative efficiency of the isolates. The halo and colony diameters were measured at 13 and 21 days post inoculation. Halo size was described by using solubilization index (ratio of total diameter (colony + halo zone) to the colony diameter). Indole acetic acid production was tested using the calorimetric Salkowski reagent according to Glickmann and Dessaux (1995) ^[11].

Genomic DNA extraction and amplification of the 16S rRNA

The 16S rRNA gene of the bacterial isolates was amplified by polymerase chain reaction using universal primers 27F 5'-GAGTTTGMTCTGGCTCA-3' and 1492R, 5'-TACGGYTACCTACGACT-3' (Bioneer, USA) as described by Embley and Stackebrandt (1994) ^[12] using an Eppendorf AG, model 22331 thermal cycler (Hamburg, Germany).

The PCR mixture comprised of 0.2 Units of Taq polymerase, 20 pmol of 27 F forward primer, 20 pmol of 1492R reverse primer, 1.25 mM dNTPs mix (QIAGEN), 10x PCR buffer (QIAGEN), 1 ml of template DNA and 29.8 ml of PCR water. The thermal program was initial denaturation at 94°C for 5 min, denaturation at 94°C for 30 s, primer annealing at 43°C for 2 min, chain extension at 72°C for 1.5 min for 35 cycles, followed by final extension at 72°C for 5 min. The amplicons were purified with the GeneJET Purification kit (Thermo Fisher Scientific Inc.) and subjected to DNA sequencing according to the manufacturer's protocol (Macrogen, South Korea) using 27 F (5'-GAGTTTGMTCTGGCTCA-3') and 1492R (5'-TACGGYTACCTACGACT-3') primers.

The sequences were processed using the Chromas Pro software (www.technelysium.com.au) and aligned using CLUSTAL Omega program (<http://www.clustal.org/used>). They were used in a BLAST search against sequences hosted at the National Center for Biotechnology Information (NCBI) database to determine their closest neighbours using (BLASTn algorithm, <http://www.ncbi.nlm.nih.gov>) as described by Altschul et al. ^[13]. A phylogenetic tree was constructed using neighbor-joining method ^[14] by using MEGA V5.10 package (www.megasoftware.net) while evolutionary distances were computed using the Maximum Composite Likelihood method ^[15]. To obtain statistical support values for the branches, bootstrapping was conducted with 1000 replicates and all sites, including gaps in the sequence alignment, were excluded pairwise in the phylogenetic analysis ^[16]. The partial sequences for the 16S rRNA genes were deposited in the GenBank sequence database and assigned accession numbers KP797914 to KP797931. The taxonomic assignment was confirmed using the naïve Bayesian rRNA classifier on the RDP website at a 95% confidence level ^[17].

Assessment of growth enhancement effect on tissue culture banana plantlets

Based on plant growth promoting traits screened previously, five strains namely *Chryseobacterium* spp., *Bacillus megaterium*, *Bacillus flexus*, *Pseudomonas montellii* and *Streptomyces* spp. were assessed for plant growth enhancement under controlled greenhouse conditions. Bacterial cultures prepared as previously described by Mia et al. were infected on tissue culture banana plantlets by dipping the 3 month old plantlets into the specific bacterial isolate treatment for 48 hours at room temperature while distilled water was used for the control experiment. The plantlets were grown in a sterile nutrient inert Cocopeat media (Amiran, Kenya) supplemented with nitrogen, phosphorus and potassium (NPK) slow release fertilizer at 5 levels (0 gms, 2.5 gms, 5 gms, 7.5 gms and 10 gms) in a random block design in three replicates ^[18].

Agronomical features of the plantlets: height, diameter and number of green and dry leaves were assessed at 14 days intervals for 3 months [19] after which the plants were destructively sampled by gently uprooting, washing off the media under running tap water and measuring the fresh weight. Dry weight was taken after oven drying the plantlets at 80°C for 14 days then essential nutrient levels of nitrogen, phosphorus, potassium, calcium, iron and magnesium was assessed according to Plank [20].

Statistical analyses

The agronomical data, mean ± SD from three experimental replicates were collected for analysis. Differences in agronomical parameters among the bacterial treatments were tested by analysis of variance procedure (ANOVA) using SPSS software. The student t-test was used to assess the variation between each of the treatments taken against the control experiment (Walpole 1986). A probability of 5% was considered significant ($P \leq 0.05$).

RESULTS AND DISCUSSION

Isolation and profiling of banana rhizosphere isolates

After isolation and culturing, 20 pure isolates were obtained from 67 soil samples based on colony morphological and biochemical characterization revealing relative bacterial diversity. Nitrogenase activity was seen in 19 isolates, 18 (90%) solubilized phosphates while none of the bacterial strains produced the auxin indole acetic acid hormone. Catalase activity was observed in all the isolates (Table 1).

Table 1. Morphology, physiological and biochemical characterization of rhizobacteria isolated from banana farms in Juja, Kenya.

Serial no	Isolate code	Gram test	Morphology	Catalase test	Urease test	Nitrate reduction	Citrate utilization	Hydrogen sulfide production	Phosphate solubilisation	Methyl Red	Voges Proausker	Acetylene Reduction Assay	Indole Acetic Acid	Gelatin hydrolysis	Casein hydrolysis	Starch hydrolysis	Probable identity
1	F1RWL	+	Cocci	+	+	-	-	-	+	+	+	+	-	+	+	+	<i>Staphylococcus</i> spp.
2	F1RWL	-	Rods	+	-	-	-	-	-	-	-	+	-	+	+	+	<i>Chryseobacterium</i> spp.
3	F2RNY	+	Cocci	+	+	-	-	-	-	+	-	+	-	-	+	-	<i>Staphylococcus</i> spp.
4	F1RWL	+	Cocci	+	+	-	-	-	+	+	+	+	-	+	+	+	<i>Staphylococcus</i> spp.
5	F5RNY	+	Rods	+	-	-	-	-	+	+	-	+	-	+	+	+	<i>Bacillus</i> spp.
6	F6RIS	+	Rods	+	-	+	-	-	+	+	-	+	-	+	+	+	<i>Bacillus</i> spp.
7	F5RMS	+	Rods	+	-	-	+	-	+	+	-	+	-	+	+	-	<i>Bacillus</i> spp.
8	F4RKB	+	Rods	+	-	+	+	-	+	+	+	+	-	+	+	+	<i>Bacillus</i> spp.
9	F5RMS	-	Rods	+	+	-	-	-	+	-	-	-	-	+	+	+	<i>Chryseobacterium</i> spp.
10	F1RGC	+	Rods	+	-	-	+	-	+	+	+	+	-	+	+	-	<i>Bacillus</i> spp.
11	F1RWL	+	Rods	+	+	-	+	+	+	-	-	+	-	+	+	+	<i>Streptomyces</i> spp.
12	F1RGN	+	Rods	+	-	+	+	-	+	+	+	+	-	+	-	-	<i>Paenibacillus</i> spp.
13	F1RGC	+	Rods	+	-	+	+	-	+	+	+	+	-	+	+	+	<i>Bacillus</i> spp.
14	F4RNY	+	Rods	+	-	+	+	-	+	+	+	+	-	+	+	+	<i>Bacillus</i> spp.
15	F6DGG	+	Rods	+	-	-	+	+	+	+	+	+	-	+	+	+	<i>Bacillus</i> spp.
16	F6RKB	+	Rods	+	-	-	+	-	+	+	+	+	-	+	+	-	<i>Bacillus</i> spp.
17	F1RVL	+	Rods	+	-	+	+	-	+	+	+	+	-	+	+	+	<i>Bacillus</i> spp.
18	F1RWL	+	Rods	+	-	+	+	-	+	+	+	+	-	+	+	+	<i>Bacillus</i> spp.
19	F6RMB	-	Rods	+	-	-	+	-	+	-	-	+	-	-	+	-	<i>Pseudomonas</i> spp.
20	F5RNY	+	Rods	+	+	-	+	+	+	-	-	+	-	+	+	+	<i>Streptomyces</i> spp.

The *Bacillus* spp. and *Pseudomonas* spp. isolates showed strong phosphorus solubilization activity than the rest of the isolates. *Pseudomonas* spp. had phosphate solubilization activity of 1.6 being the highest, followed by *Bacillus* spp. having 1.5 and 1.45 on the phosphate solubilization index on agar assay (Table 2). Phosphate solubilization ability of the plant growth promoting rhizobacteria is considered to be one of the most important traits associated with plant phosphorous nutrition. Soil or seed inoculations with phosphate solubilizing microbes (PSM) have been used for the improvement of crop growth and production by the solubilization of fixed and applied phosphates [21].

Table 2. The extent of phosphate solubilisation on agar and concentration of ethylene produced by the bacterial isolates.

Serial No.	Isolate code	Phosphate solubilization index on agar assay	Concentration of Ethylene (µl/ml)
1	F1RWL	1.33	0.0309

2	F1RWL	-	0.0145
3	F2RNY	-	0.0113
4	F1RWL	1.25	0.0474
5	F5RNY	1.33	0.0252
6	F6RIS	1.2	0.0635
7	F5RMS	1.25	0.0706
8	F4RKB	1.25	0.0510
9	F5RMS	1.33	-
10	F1RGC	1.33	0.0067
11	F1RWL	1.11	0.1459
12	F1RGN	1.2	0.0611
13	F1RGC	1.33	0.0339
14	F4RNY	1.2	0.0669
15	F6DGG	1.25	0.1515
16	F6RKB	1.5	0.7547
17	F1RVL	1.11	0.0466
18	F1RWL	1.42	0.0631
19	F6RMB	1.6	0.0508
20	F5RNY	1.2	0.0508

All *Bacillus* isolates were found to have nitrogenase activity and one of the *Bacillus* spp. showing the highest concentration of 0.7547 µl/ml followed by another with 0.1515 µl/ml indicating their potential in fixing nitrogen (**Table 2**). From **table 2**, *Chryseobacterium* spp. showed a nitrogenase activity of 0.069 µl/ml and 0.0145 µl/ml which could be a potential nitrogen fixer since members of these genus exhibit plant-growth promoting activities as they are universal symbionts of some higher plants ^[22,23].

Two *Staphylococcus* spp. with nitrogenase and phosphate solubilisation abilities were identified which to our knowledge has not been isolated from banana rhizosphere. Cases of this genus promoting plant growth have rarely been documented. Nitrogen is a crucial element for bananas as it may be a limiting factor in the production and growth of bananas. Ahemad and Khan demonstrated that nitrogen is among the nutrients commonly required by the banana plant for their optimal growth and productivity ^[1,24].

The spore forming *Paenibacillus xylanilyticus* identified showed promising plant growth promoting nitrogenase and phosphate solubilisation activities. Some species of *Paenibacillus* have consistently shown ability to fix atmospheric nitrogen in vitro ^[25].

Though reported to be one of the dominant species in banana rhizosphere, the present study isolated only one *Pseudomonas monteilii* with nitrogenase and phosphate solubilisation ability. This bacterium has been suggested to belong to the sub group of *P. putida* which are ecologically important in colonizing the banana plant roots rhizosphere ^[26]. Only two *Streptomyces* spp. were identified despite being shown as important microorganisms present in the banana root rhizosphere that have antagonism to phytopathogenic fungi ^[27]. *Streptomyces* strains promote banana plantlets' growth by supplying iron to them.

Bacterial identification using the amplification of the 16S rRNA

Phylogenetic relationship of the bacterial isolates sequenced was inferred using the Neighbour Joining method. Eighteen (18) sequences of the 20 isolates in this study were included while two (2) were excluded in subsequent analysis since they had chimeras. The bacterial isolates clustered with sequences from different genera: three isolates were categorized as *Staphylococcus* spp., two (10%) as *Streptomyces* spp., two (10%) as *Chryseobacterium jejuense* strain NW50, while 11 (55%) were categorized as belonging to the genus *Bacillus*. With 100% sequence similarity, one isolate was characterized as *Paenibacillus* spp., and also *Pseudomonas monteilii* strain IHB B 2329 (**Figure 1**).

Assessment of growth enhancement effect on tissue culture banana plantlets

Inoculation of the plants with five selected isolates had a positive growth influence on tissue cultured banana plantlets treated with 5 gms, 7.5 gms and 10 gms levels of NPK fertilizer levels in height, diameter, number of green leaves and mass in dry weight compared with their respective controls. The direct effect of the inoculants on the plantlets could not be established because of the confounding fertilizer level effect. The mean plant diameter (cm) and number of green leaves in isolates plantlets treated with *Pseudomonas monteilii* and *Streptomyces* spp. were higher than their controls, as similarly observed in the mean diameter for plantlets treated with *Chryseobacterium* spp., *Bacillus megaterium* and *Bacillus flexus* strains at 10 gms fertilizer level. Plantlets treated with *Streptomyces* spp. isolate registered a significantly higher growth in all the parameters ($p < 0.05$). Height and dry weight were significantly higher in all the five strains tested ($p = 0.000$) except *Pseudomonas monteilii* in height ($p = 0.007$). Diameter was significantly higher on plants treated with *Streptomyces* spp. ($p = 0.000$) at 10 gms fertilizer level (**Figure 2**). The total accumulation of nitrogen, phosphorus, magnesium and calcium was significantly higher ($p = 0.00000684$, 0.000163 , $4.46E-09$ and $3.77E-05$ respectively) in all the treatments following inoculation with the isolates. This could be attributed to the effect of the inoculation on the uptake of these nutrients. However, the intake of potassium, zinc and iron was not influenced by inoculants since there were no differences observed between the treatments and the controls. Plant growth promoting rhizobacteria inoculation on bananas substantially increases the development and growth of tissue-cultured banana plantlets ^[19].




Figure 1. Neighbour joining phylogenetic tree showing the position of the banana rhizospheric isolates from Juja, Kenya. Eighteen strains from seven different farms in Juja that were isolated in this study are indicated in bold font. The accession numbers of the sequences are indicated in parantheses. The bootstrap values are the numbers seen at the nodes based on 1000 replications. The 16S rRNA gene sequence of *Methanoculleus thermophilus* was an outgroup used.




Figure 2. Effect of selected bacteria isolates from Juja on banana plant growth (A: Mean diameter, B: Mean height, C: Mean number of green leaves, D: Mean number of dead leaves, E: Mean mass in dry weight). An asterisk (*) means significantly different to the control (p < 0.05).

The present study shows that banana rhizospheric bacterial isolates from *Pseudomonas*, *Bacillus*, *Staphylobacterium*, *Chryseobacterium*, *Streptomyces* and *Paenibacillus* genera in Juja, Kenya harbor plant growth promoting traits and therefore these isolates can be used as biofertilizers in agriculture to promote the growth as well as production of bananas.

ACKNOWLEDGMENTS

This study was supported by a research grant from the Research, Production and Extension Division (RPE) of Jomo Kenyatta University of Agriculture and Technology-Kenya. We wish to acknowledge the departments of Horticulture and Botany (JKUAT) and their technical staff for technical support and availing their laboratory facilities for this study.

Special appreciation goes to Mr. Zachary Ndung'u (JKUAT) for his support during sample collection and green house experiments.

REFERENCES

1. Saharan BS, Nehra V. Plant Growth Promoting Rhizobacteria: A Critical Review. Life Sciences and Medicine Research 2011; LSMR-21.
2. Barmak JH et al. The Role of Plant Growth Promoting Rhizobacteria (PGPR) in Sustainable Agriculture. Advances in Environmental Biology 2011; 5: 3079-3083.
3. Mia BMA et al. Use of Plant Growth promoting Bacteria in Banana: A New Insight for Sustainable Banana Production. International Journal of Agricultural Biology 2010;12: 459-467
4. Van Loon LC Plant responses to plant growth-promoting rhizobacteria. European Journal of Plant Pathology 2007; 119: 243-254.
5. Ngamau CN et al. Isolation and identification of endophytic bacteria of bananas (*Musa* spp.) in Kenya and their potential as biofertilizers for sustainable banana production. African Journal of Microbiology Research 2013; 6: 6414-6422
6. Reena T et al. Isolation of Phosphate Solubilizing Bacteria and Fungi from Rhizospheres soil from Banana Plants and its Effect on the Growth of *Amaranthus cruentus* L. IOSR Journal of Pharmacy and Biological Sciences 2013; 5: 6-11.
7. Vessey JK Plant growth promoting rhizobacteria as biofertilizers. Plant and Soil 2003; 255: 571-586
8. Cappuccino JG et al. Microbiology: a laboratory manual (8th ed) Pearson. (2002) ISBN-13: 978-0805325782. Jaypee Brothers Medical Publishers (P) Ltd, New Delhi
9. Eckert B et al. *Azospirillum doedereinerea* species, a nitrogen- fixing bacterium associated with the C4 grass Miscanthus. International Journal for system evolution Microbiology 2001; 51: 17-26.
10. Nautiyal CS. An efficient microbiological growth medium for screening phosphate solubilizing microorganisms. FEMS Microbiol. Letters 1999; 170: 265-270
11. Glickmann E et al. A critical examination of the specificity of the Salkowski reagent for indolic compounds produced by phytopathogenic pathogenic bacteria. Applied and Environmental Microbiology 1995; 61: 793-796.
12. Embley TM et al. The molecular phylogeny and systematics of the actinomycetes. Annual Review of Microbiology 1994; 48: 257-289.
13. Altschul SF et al. (1990) Basic Local Alignment Search tool. Journal of Molecular Biology 1990; 215: 403-410.
14. Saitou N et al. The neighbor-joining method: a new method for reconstructing phylogenetic trees. Molecular Biology Evolution 1987; 4: 406-425.
15. Tamura K et al. Prospects for inferring very large phylogenies by using the neighbor-joining method. Proceedings of the National Academy of Sciences 2004; 101: 11030-11035.
16. Felsenstein J Phylogenies and the Comparative Method. The American Naturalist 1985; 125: 1-15.
17. Cole JR et al. The Ribosomal Database Project (RDP-II): Sequences and tools for high-throughput rRNA analysis. Nucleic Acids Response 2005; 33: D 294-D296.
18. Mia BMA et al. Effect of plant growth promoting rhizobacterial (PGPR) inoculation on growth and nitrogen incorporation of tissue-cultured *Musa* plantlets under nitrogen free hydroponics condition. AJCS, 2009; 4: 85-90
19. Mia MAB et al. The effect of rhizobacterial inoculation on growth and nutrient accumulation of tissue-cultured banana plantlets under low N-fertilizer regime. African Journal of Biotechnology 2009; 8: 5855-5866
20. Plank CO Plant analysis reference procedures for the southern region of the United States. (1992) Southern Cooperative Series Bulletin #368
21. Banerjee S et al. Stress induced phosphate solubilization by *Arthrobacter* sp. and *Bacillus* sp. isolated from tomato rhizosphere. Australian Journal of crop science 2010; 4: 378-383.

22. Montero-Calasanz M et al. *Chryseobacterium hispalense* sp. nov., a plant growth promoting bacterium isolated from a rainwater pond in an olive plant nursery, and emended descriptions of *Chryseobacterium defluvii*, *Chryseobacterium indologenes*, *Chryseobacterium wanjuense* and *Chryseobacterium gregarium*. International Journal of Systematic and Evolutionary Microbiology 2013; 63: 4386–4395.
23. Weller DM (1994) Current challenges in introducing beneficial microorganisms into the rhizosphere microorganisms. In: F. O’Gara (eds), et al. Molecular Ecology of Rhizosphere Microorganisms, VCH, New York. pp. 1–18.
24. Ahemad M et al. Functional aspects of Plant Growth Promoting Rhizobacteria: Recent Advancements. Insight Microbiology 2011; 1: 3:39-54
25. Jin H-J et al. *Paenibacillus sophorae* sp. nov., a nitrogen-fixing species isolated from the rhizosphere of *Sophora japonica*. International Journal of Systematic and Evolutionary Microbiology 2011; 61: 767–771
26. Naik PR et al. Genetic and Functional Diversity among *Fluorescent Pseudomonads* Isolated from the Rhizosphere of Banana. Microbial Ecology 2008; 56: 492–504
27. Cao L et al. Isolation and characterization of endophytic *streptomycete* antagonists of *fusarium* wilt pathogen from surface-sterilized banana roots. FEMS Microbiology Letters 2005; 247: 147-152.