

The Habitat Preference of Hartebeest Antelopes (*Alcelaphus buselaphus*) in Bouba-ndjidda National Park, North Region, Cameroon

Melle Ekane Maurice^{1*}, Kamgang Serge Alex¹, Ewane Divine¹, Kamah

Pascal Bumtu¹, Mbole Veronique², Esong Lionel Ebong²

¹Department of Forestry and Wildlife, University of Buea, Buea, Cameroon

²Department of Environmental Science, University of Buea, Buea, Cameroon

Research Article

Received: 11-Jun-2022, Manuscript

No. JEAES-22-66426; **Editor**

assigned: 14-Jun-2022, Pre QC No.

JEAES-22-66426 (PQ); **Reviewed:**

29-Jun-2022, QC No. JEAES-22-

66426; **Revised:** 12-Aug-2022,

Manuscript No. JEAES-22-66426

(R); **Published:** 22-Aug-2022, DOI:

10.4172/2347-7830.10.07.002

***For Correspondence:** Melle Ekane

Maurice, Department of Forestry and

Wildlife, University of Buea, Buea,

Cameroon, **Email:**

melleekane@gmail.com

Keywords: Natural coagulants; *Vicia*

villosa roth; Turbidity removal; Water

ABSTRACT

The availability of preferred habitat determines the spatial and temporal distribution of herbivores in savanna woodland ecosystems. Understanding habitat preference of a targeted wildlife species is crucial for developing effective management strategies. Hence, the objective of this study was to examine an importance of various habitats to the hartebeest population in the national park. Research data was collected on animal group activity through observation. The observed animal group activity was recorded on check -sheets alongside some ecological parameters. The survey showed a significance between the antelope-group size and behavior $X^2 = 5.441$ $df=3$, $P<0.05$. Nonetheless, the survey witnessed animal group size ranging from 1 -5 (22%) and 6-10 (78%) respectively. More so, antelope's movement recorded a significant activity 70% compared to feeding 20%, rest 7% and drinking 4% respectively. There was a significance between antelope group behavior and vegetation $X^2 = 9.723$ $df=6$, $P<0.05$. Three vegetation types were considered during this study, grassland 48%, shrub -land 45%, and forest patches 7% respectively. This study observed the hartebeest antelopes feeding on grass vegetation 93% and lower branches of shrubs 7% due to their large body sizes. Also, there was a significance between landscape and hartebeest -group behavior $X^2 = 34.371$ $df=9$, $P<0.05$. Antelope -group behavior was predominantly recorded in flat landscape areas 55% than slope 30%, hill 11%, and flood plains 4% respectively. Furthermore, landscape recorded a significance on vegetation types $X^2 = 3.332$ $df=6$, $P<0.05$.

Boubandjidda national park is rich in wildlife population such as antelopes; hence, more study is needed to be done to understand the population dynamics and behavior of the hartebeest antelopes. Unfortunately, the wildlife population in the national park does not have a rich database research reference compare to other national parks in sub Saharan Africa region.

INTRODUCTION

Effective management of multi species ungulate populations requires knowledge of the habitat requirements of the individual species and should take cognizance of the existence of interspecific competition. The extent of resource available must also be quantified to prevent environmental degradation. The relationship between animal and habitat plays a crucial role in the ecology of any species. The association between an herbivore and its habitat not only encompasses the availability of preferred food plants and their particular growth stages in its home range, it is also intimately linked to other features of the habitat [1]. In the last three decades, numerous studies have been conducted of mammalian herbivores in relation to their environment. These range from qualitative observations of the types of habitats used by particular ungulate species or assemblages to more complex quantitative investigations that attempt to explain why such habitats are used [2,3]. Studies aimed at establishing the habitat preferences and other ecological requirements of herbivorous animals are particularly important for the conservation of vulnerable species [4].

Identifying the quality and preference of different habitat types are crucial for developing conservation strategies of a targeted wildlife species [5, 6]. Herbivores are known to select habitats that provide maximum forage intake, while reducing predation risk. There are several factors that can determine the spatial and temporal distribution of herbivores in savannas ecosystems. These include the availability of resources, predation risk, fire, vegetation height and cover, human presences and livestock density [7-11]. Since a habitat type may not always have adequate resources, the tradeoffs between costs and benefits associated with searching and utilizing forage can limit herbivore selection [12]. Moreover, spatial variation in relative availability of different habitat types may result in dissimilar habitat selection among individuals of the same species [13].

In savanna grassland, where there is a cyclic rainfall, fire is used as one of the most important habitat management tool for herbivores [14]. Understanding how wildlife species respond to fire effects is crucial, particularly for endangered species that have limited range [15]. Fire effects grass height, which in turn affects habitat preference of herbivores [16]. Previous studies have identified the trends of large grazers' habitat preference in response to grass height and post-fire effect on vegetation. Herbivores could optimize their daily forage need where they are able to access the preferred grass heights [17]. There is a general consensus that grass height has a major influence on the spatial and temporal distribution of herbivores, and resource partitioning among herbivores could also occur through differential selection of grass height [18].

In African savannas, frequent burning of grass influences the habitat selection of herbivores due to impacting forage quality and reducing predation risk, and it is a key element in predicting habitat selection by specific species. Fire plays a

determinant role in the ecology and evolution of grassland eco-systems, and has historically, and still today, been used as a tool for managing grassland vegetation [19-21]. Post fire regrowth of grass influences the dry season habitat use of many herbivore species. However, there have been arguments among ecologists how burning affects habitat selection of large body sized herbivores.


Small body sized herbivores might prefer burned areas more than large body-sized herbivores due to differential preferences in relation to forage quality [22]. However, another study revealed that fire does not have relationship between body size and use of burned areas. Several studies found that decreasing fire frequency increases vegetation cover and tree densities, which in turn decreases visibility and the subsequent ability of herbivores to detect and escape from predators. As a result, herbivores may avoid areas with relatively denser vegetation cover or spend more time in those areas for vigilance rather than foraging [23, 24]. Hence, herbivores foraging in burned areas may represent either acquiring quality forage or avoiding predators. Swayne's hartebeest (*Alcelaphus buselaphus swaynei*) is a large body -sized herbivore weighing between 100 and 200 kg [25]. It was once widely distributed in Ethiopia, Somalia and Djibouti, but currently its range is confined in two protected areas: Senkele Swayne's Hartebeest Sanctuary and Maze National Park in Ethiopia and listed as endangered sub-species by IUCN Red list [26, 27]. Seasonal burning is used as a habitat management tool in the national parks, but how the Swayne's hartebeests respond to post fire effect and grass height preferences in different seasons remains untouched. Despite its small area, the Park has different habitat types. While hartebeest are known to be grazers, there may be conditions that enforce Swayne's hartebeests to utilize bush -land and forest habitats in different seasons [28].

MATERIALS AND METHODS

Description of study area

Bouba ndjidda national park is located in the northern region of Cameroon near the Chadian border, between latitude 8° 37' and 8° 37' N and longitude 14° 39' and 14° 39' E. It was created in 1932 as a wildlife reserve and became a national park in 1968. It covers an area of 220,000 ha. and has the Sudano -guinean climate characterized by two seasons, a six month rainy season from late April to mid -October and a dry season from November to April during the period which no rain is observed. The region receives between 1000 and 1,250 mm of precipitation per year and the rainiest months are August and September while the annual average temperature is 28 0 C. The region has a dense hydrographic network, unlike other rivers that are seasonal in this part of the country. Water is abundant throughout the national park and even during the dry season, thus ensuring the development of ecological interaction processes between wildlife species and habitat characteristics in the environment. The national park is also a home to 24 species of large and medium mammals such as African elephant, Lions, Giraffes, Leopards, Hyenas, Bush pig, Warthogs, Hippos, Buffaloes, Hartebeest antelopes, Eland antelopes, Kob antelopes, Reedbuck antelopes, etc. Additionally, the national park homes about 250 bird species. Considering the ornithological richness of the national park, it has been declared and area of importance for the conservation of birds (Figure 1).

Figure 1. Map of bouba ndjidda National park.



Data collection

Data collection on antelopes activity in the preferred habits were done by direct observation (focal sampling method) from appropriate visible points. This included watching an individual or group of antelopes for 10 minutes within an interval of 5 minutes. Antelopes were observed in the early morning (06:30 to 10:30) and in the late afternoon (14:00 to 18:00) when they were active [29]. Recordings were ceased whenever the animals moved out of sight. Data were collected more than ten times per month on antelope’s habitat activity. More so, observations were recorded when the antelopes were foraging (grazing/browsing/chewing/biting) on plants (tree/shrub/herb/grass) or consumed plants parts (leaf/shoot/stem /fruit) [30].


Data analysis

Before conducting the actual data analysis, data were checked for some errors (e.g. recording and consistencies). Data were analyzed using Statistical Package for Social Science (SPSS) version 20 software ($\alpha=95\%$ level of significance, $p<0.05$), descriptive statistics and Excel spreadsheet. The variables were cross -tabulated to understand their degree of association.

RESULTS AND DISCUSSION

The survey showed a significance between the antelope group size and behavior $\chi^2 = 5.441$ $df=3$, $P<0.05$ (Figure 2).


Figure 2. Antelope group sizes and behavior.



The population of hartebeest in Boubouba ndjidda national park is seemingly high compared to other antelope species. Some of the reasons, however, may be an ecological system rich in woody grassland vegetation suitable for the feeding of these antelopes. Hart ebeest is one of the largest antelope species in sub Saharan Africa that form herds of 30 to 300 animals in food -rich ecosystems. Nonetheless, the survey witnessed animal -group sizes ranging from 1 -5 (22%) and 6-10 (78%)


respectively (Figure 3).

Figure 3. Antelope group sizes.



The low group formation strength in the national park might be due to the dry season food scarcity witnessed by this study, carried out in the dry season. More so, antelopes' movement witnessed a significant activity 70% compared to feeding 20%, rest 7% and drinking 4% respectively (Figure 4).


Figure 4. Antelope behavior.



The highest behavioral activity observation. During the dry season of the year, the woodland savanna ecosystem of northern Cameroon experiences vegetation dryness due to low humidity and soil moisture content caused by the hot weather condition. This condition causes feeding problems to herbivorous antelopes such as the hart beest antelopes that predominantly depend much on grass nutrient.

There was a significance between antelope -group behavior and vegetation $\chi^2 = 9.723$ $df=6$, $P < 0.05$ (Figure 5).

Figure 5. Antelope behavior and vegetation.




Vegetation availability has been a determinant factor to the wildlife population availability in protected area management. The roles played by vegetation in the management of wildlife population are enormous. One of the most important roles is the provision of habitat for wildlife shelter; secondly, it's a food resource to herbivorous wildlife species such as hartebeest antelopes (Figure 6).

Figure 6. Vegetation cover.




Though, Boubandjidda national park is a woodland savanna ecosystem, it has rich vegetation during the raining season than the dry season. Both seasonal changes affect animal population increase, however, the wet season experiences higher increase than the dry season. The rain gives rise to healthy vegetation growth needed to feed herbivorous antelopes and other wildlife species, creating a fertile reproduction environment. Three vegetation types were considered during this study, grassland 48%, shrub -land 45%, and forest patches 7% respectively. This study observed the hartebeest antelopes feeding on grass vegetation 93% and lower branches of shrubs 7% due to their large body sizes (Figure 7).

Figure 7. Food resources.




Also, there was a significance between landscape and hartebeest -group behavior $\chi^2 = 34.371$ $df=9$, $P < 0.05$ (Figure 8).

Figure 8. Landscape and hartebeest behavior.




Antelope group behavior was predominantly recorded in flat landscape areas 55% than slope 30%, hill 11%, and flood plains 4% respectively (Figure 9).

Figure 9. Landscape.



Flat landscapes were richer in grass vegetation than other areas in the national park. The high behavioral activity recorded on the flat landscape was on animal movement in search of food. Though, open flat landscapes might expose hartebeests to predatory big cats such as lions and leopards to be sighted easily, the animal herd was always vigilant to their presence and would often trigger running whenever these cats were sighted in their territory. Landscape recorded a significance on vegetation $\chi^2 = 3.332$ $df=6$, $P<0.05$ (Figure 10). The national park is not rich in forest vegetation, but the presence of few forest patches helps to shade wildlife from the hot sun during the dry season, especially during the mid-day and afternoon periods when the sun is extremely hot. The grassland and shrub-land were major contributors in feeding these herbivorous antelopes.

Figure 10. Landscape and vegetation.



DISCUSSION

The availability of preferred habitats determines the spatial and temporal distribution of herbivores in savanna ecosystems. Understanding habitat preference of a targeted wildlife species is crucial for developing effective conservation strategies [31]. Habitat preference of large grazers in connection to grass height and post-fire effect has been debated for the last century. The effects of season, grass height and burning on the habitat preference on Swayne's hartebeest (*Alcelaphus buselaphus swaynei*) in Maze National Park were examined. Data for seasonal habitat selection were collected using both direct observation along established transect lines and pellet counting using permanently established plots. Every month, we measured grass height commonly preferred by Swayne's hartebeest in grassland habitat.

The grass height has been demonstrated to exert a major influence on bite size that in turn impacts on food intake rate achieved by grazing herbivores. Larger body sized herbivores (>100 kg body weight) are expected to graze taller grasses to meet their quantitative food requirements, while smaller body sized herbivores can achieve an adequate amount of food intake from short grass swards [32, 33]. In theory, shorter grasses are generally leafy with higher proportion of nutrients and preferred by many small body sized herbivores, while larger body sized herbivores can tolerate poorer quality food provided by the taller grasses [34]. When grass grows and matures, its nutritional quality decreases [35]. This can be demonstrated by the decrease proportion of leaves and the nitrogen content (both indicating high grass quality) in the grass with increasing grass mass in savanna ecosystem [36]. Our study showed that Swayne's hartebeests preferred open grassland habitat in Maze National Park throughout the year as observed with other wild herbivores, such as Coke's hartebeest (*Alcelaphus buselaphus cokii*) in Athi-Kapiti Plains, Kenya; hartebeests (*Alcelaphus buselaphus*) in southern border of Burkina Faso and wildebeest (*Connochaetes taurinus*) in Serengeti National Park, Tanzania [37,38]. Although the Park has a wider coverage of other habitat types, such as bush-land habitats and riverine forest, the Swayne's hartebeest rarely used them. This reflects on the fact that Swayne's hartebeest conservation is largely based on the management of the grassland habitat in Maze National Park. Our surveys detected few Swayne's hartebeest pellets in bush land habitats and riverine forests during the dry season, which likely occurred when they were walking to a water source. Swayne's hartebeests were not encountered in agricultural lands and rugged habitats except in a rare occurrence, which might have been a response to predators in the area. In the grass and habitat, the grass grows fast and reaches above one meter within a month after the wet season begins and becomes taller in the early-dry season, but decreases in height in the first few months of dry season [39]. However, Swayne's hartebeests almost abandoned the taller grass height, and consistently preferred the shorter (below 30 cm) available grass height areas in the Park. Our findings are thus in support of the previous studies in other areas, for instance, hartebeests and roan antelope (*Hippotragus equinus*) in Nazinga Game Ranch, Burkina Faso and wildebeest in Serengeti Park, Tanzania preferred short grass height. There are two speculations about short grass preferences of herbivores: (1) due to the higher nutritional quality of short grasses and (2) to avoid predation risk [40]. Even though the fear of predation may influence short grass habitat selection in some cases, in this study the predation risk is rather less due to low density of predators (mainly lions) in the Park, suggesting that the Swayne's hartebeest preference of short grass habitats is more likely the result from nutritional gain. Shorter grasses have less lignin with lower carbon to nitrogen ratios which are more palatable and digestible for grazers. Shorter grasses also have higher nutritional quality and percentage of green leaves that allow higher bite rates for herbivores foraging [41]. Grass height preferences of

Swayne's hartebeest influence their distribution in Maze National Park in different seasons. Previous studies also revealed that forage influences the distribution of herbivores [42].

CONCLUSION

The benefits of group living in wildlife are enormous, survival of sub-adults and juveniles, location and movement to new feeding sites, and mostly importantly defense against invasion. Boubandjidda national park is rich grass vegetation on flat landscape sustaining a huge population of many herbivorous wildlife species, especially antelope species like hartebeest. The huge antelope population in these areas has also been credited to the conservation authorities of the national park and the Ministry of Forestry and Wildlife. Poaching is believed to have been significantly reduced to the advantage of wildlife population increase. Secondly, the drainage system of this area is the best in the northern region of the country, an advantage to antelope-group feeding along the river banks during the dry season period of the year when rivers in other protected areas like Faro and Benue must have dried up by the

REFERENCES

1. Pienaar, et al. Habitat preference in South African antelope species and its significance in natural and artificial distribution patterns. *Koedoe*. 1974;17:185-195.
2. Williamson DT, et al. An outline of the ecology and behaviour of red lechwe (*Kobus leche leche* Gray 1850). *Afr J Ecol*. 1990;28:89-101.
3. Funston PJ, et al. A seasonal variation in the movement patterns, home range and habitat selection of buffaloes in a semi arid habitat. *Afr J Ecol*. 1994;32:100-114.
4. Ben-sharar R, et al. Resource availability and habitat preferences of three African ungulates. *Biol Conserv*. 1990;54:357-365.
5. Illius A, et al. Modelling the nutritional ecology of ungulate herbivores: evolution of body size and competitive interactions. *Oecologia* 1992;89:428-434.
6. Rapp E, et al. Micro habitat selection of herbivores in response to perceived predation risk and forage quality in Hluhluwe-I Mfolozi game reserve. Swedish University of Agricultural Sciences, Department of Wildlife, Fish, and Environmental studies, Umea, 2017;1-26.
7. Martinez-Freiria F, et al. Contemporary niche contraction affects climate change predictions for elephants and giraffes. *Divers Distribute*. 2016;22:432-444.
8. Ogutu JO, et al. Herbivore dynamics and range contraction in Kajiado County Kenya: climate and land use changes, population pressures, governance, policy and human-wildlife conflicts. *Open Ecol J*. 2014;7:9-31.
9. Sankaran M, et al. Woody cover in African savannas: the role of resources, fire and herbivore. *Glob Ecol Biogeogr*. 2008;17:236-245.
10. Laca E, et al. Effects of sward height and bulk density on bite dimensions of cattle grazing homogeneous swards. *Grass Forage Sci*. 1992;47:91-102.
11. Fritz H, et al. Habitat use by sympatric wild and domestic herbivores in African savanna woodland: the influence of cattle spatial behavior. *J Appl Ecol*. 1996;33:589-598.

12. Arsenault R, et al. Resource partitioning by grass height among grazing ungulates does not follow body size relation. *Oikos*. 2008;117:1711 –1717.
13. Klop E, et al. Resource selection by grazing herbivores on post fire regrowth in a West African woodland savanna . *Wildlife Res*. 2007;34:77 –83.
14. Klop E, et al. Savanna fires govern community structure of ungulates in Bénoué National Park, Cameroon . *J Trop Ecol*. 2008;24:39 –47.
15. Eby SL, et al. The effect of fire on habitat selection of mammalian herbivores: the role of body size and vegetation characteristics. *J Anim Ecol* 2014;83:1196 –1205.
16. Riginos C, et al. Savanna tree density, herbivores, and the herbaceous community: bottom -up vs. top-down effects. *Ecology*. 2008;89:2228 –2238.
17. Drescher M, et al. The role of grass stems as structural foraging deterrents and their effects on the foraging behavior of cattle. *Appl Anim Behav Sci*. 2006;101:10 –26.
18. Hempson GP, et al. Ecology of grazing lawns in Africa . *Biol Rev*. 2015;90:979–994.
19. Burkepile DE, et al. Habitat selection by large herbivores in a southern African savanna: the relative roles of bottom up and top down forces. *Ecosphere*. 2013;4:1–19.
20. Van Langevelde F, et al. Effects of fire and herbivore on the stability of savanna ecosystems. *Ecology* . 2003;84:337 –350.
21. Parrini F, et al. The importance of post fire regrowth for sable antelope in a southern African savanna . *Afr J Ecol*. 2010;48:526–534.
22. Wilsey BJ, et al. Variation in use of green grasses following burns among African ungulate species: the importance of body size. *Afr J Ecol*. 1996;34:32–38.
23. Hassan SN, et al. Fire history and management as determinant of patch selection by foraging herbivores in western Serengeti, Tanzania . *Int J Biodivers Sci Ecosyst Serv Manag*. 2011;7:122 –133.
24. Wagner M, et al. Impact of bush fire on grazing behavior of herbivores in Masai Mara . Master Thesis, Uppsala: Swedish University of Agricultural Sciences. 2009
25. Mamo Y, et al. Status of the Swahili Hartebeest, (*Alcelaphus buselaphus swaynei*) meta-population under land cover changes in Ethiopian protected areas . *Int J Biodivers Conserv*. 2012;4:416 –426.
26. Datiko D, et al. Population status and human impact on the endangered Swayne's hartebeest (*Alcelaphus buselaphus swaynei*) in Nechisar Plains, Nechisar National Park, Ethiopia . *Afr J Ecol*. 2011;49:311 –319.
27. Lewis J, et al. The ecology of Swayne's hartebeest . *Biol Conserv*. 1979;15:1 –12.
28. Casebeer R, et al. Food habits of wildebeest, zebra, hartebeest and cattle in Kenya Masailand . *Afr J Ecol*. 1970;8:25–36.
29. Tahani AH, et al. Diurnal Activity of Waterbuck (*Kobus ellipsiprymnus defassa*) in Dinder National Park, Sudan. *Nat Resour Environ* 2014;2:15-19
30. Wubie Bayie, et al. Population status, feeding ecology and activity pattern of common bushbuck (*Tragelaphus scriptus decula*) in Sekele Mariam Forest, West Gojjam, Ethiopia . *J Ecol Nat*. 2018;10:69 -79
31. Misganaw T, et al. The effect of season and post fire on habitat preferences of the endangered Swayne's hartebeest (*Alcelaphus buselaphus swaynei*) in Maze National Park, Ethiopia . *BMC Ecol*. 2020;20:5.

32. Bjorneraas K, et al. Habitat quality influences population distribution, individual space use and functional responses in habitat selection by a large herbivore . *Oecologia*. 2012;168:231 –243.
33. Demment MW, et al. A nutritional explanation for body size patterns of ruminant and non ruminant herbivores. *Am Nat*. 1985;125:641–672.
34. Bholá N, et al. Comparative changes in density and demography of large herbivores in the Masai Mara Reserve and its surrounding human -dominated pastoral ranches in Kenya . *Biodivers Conserv*. 2012;21:1509 –1530.
35. Thompson Hobbs N, et al. Herbivore functional response in heterogeneous environments: a contest among models . *Ecology*. 2003;84:666–681.
36. Van Langevelde F, et al. Instantaneous intake rate of herbivores as function of forage quality and mass: effects on facilitative and competitive interactions . *Ecol Model*. 2008;213:273 –284.
37. Schuette JR, et al. Diets of hartebeest and roan antelope in Burkina Faso: support of the long faced hypothesis. *J Mammal*. 1998;79:426–436.
38. Bell RHV, et al. Grazing ecosystem in the Serengeti . *Sci Am*. 1971;225:86 –93.
39. Treudt AC, et al. Herbaceous forage and selection patterns by ungulates across varying herbivore assemblages in a South African savanna . *PLoS ONE*. 2013;8: 82831.
40. Soder KJ, et al. Dietary selection by domestic grazing ruminants in temperate pastures: current state of knowledge, methodologies, and future direction . *Range Ecol Manag*. 2009;62:389 –398.
41. Bailey DW, et al. Mechanisms that result in large herbivore grazing distribution patterns. *Rangel Ecol Manag J Range Manag Archiv*. 1996;49:386 –400.
42. Masse A, et al. Habitat selection of a large herbivore at high density and without predation: trade -of between forage and cover. *J Mammal*. 2009;90:961 –970.